

Technical Comments

Comment on "Evolution of the Laminar Wake behind a Flat Plate and Its Upstream Influence"

A. F. MESSITER*

The University of Michigan, Ann Arbor, Mich.

AND

K. STEWARTSON†

The Ohio State University, Columbus, Ohio

DURING the last few years considerable effort has been expended in an attempt to understand the character of boundary-layer flow at a sharp trailing edge. The Introduction to Ref. 1 summarizes most of the published investigations of steady incompressible laminar flow past the trailing edge of a finite plate at zero incidence and high Reynolds number. In particular, asymptotic descriptions considered to be valid in the limit as $R \rightarrow \infty$ ($R = UL/\nu$; U = external flow speed, L = plate length, ν = kinematic viscosity) were given independently by Stewartson² and by Messiter.³ For the most part, Ref. 1 correctly points out the deficiencies of earlier work, but several comments about the theory derived by Stewartson and Messiter are either unjustified or incorrect.

First of all, Ref. 1 states incorrectly that the work of Ref. 4 assumed isobaric flow. The assumptions actually made were that: a) the effect of the trailing edge is confined to a distance $O(R^{-3/4}L)$; and b) the Navier-Stokes equations can be linearized about the Blasius solution in this neighborhood. The serious error lay in neglecting the discontinuity of streamline slopes in the boundary layer at $x = 1$; this was corrected in Refs. 2 and 3. Imai treated the same problem as in Ref. 4 but with the isobaric assumption. Incidentally, Ref. 4 is not without value for it still provides a useful guide to the structure of the region within a distance $O(R^{-3/4}L)$ of the trailing edge provided one adjusts the constant λ as explained in Ref. 2. Hakkinen and O'Neil,⁵ not mentioned in Ref. 1, have also discussed this region, in terms of expansions for $R^{3/4}r/L \rightarrow \infty$.

Second, it is stated in Ref. 1 that the use of linearized airfoil theory is not justified because of the large slopes of the (effective) slender body. It is certainly true that the slopes are large compared with the streamline slopes predicted by ordinary boundary-layer theory, but the slopes do remain small compared to 1. As explained in Refs. 2 and 3, $u = 1 + O(R^{-1/4})$, $v = O(R^{-1/4})$, $\partial u/\partial x = O(R^{-1/8})$, etc. In fact, Eq. (7) of Ref. 1 essentially incorporates the notion of linearization because the term $v \partial u/\partial y$ is omitted. Also, as Fig. 6 of Ref. 1 makes clear, $|c_p| \sim 0.01$ and therefore $c_p = -2(u-1)$ with error ~ 0.0001 .

Third, the main problem studied in Eqs. (4-7) of Ref. 1 contains the formulation of Refs. 2 and 3 and reduces to it as $R \rightarrow \infty$. Thus there is agreement in this limit. However, this does not provide justification for neglecting terms when $R = 10^5$. A number of terms are omitted in Ref. 1, but perhaps the most important is the pressure variation across the boundary layer. It is established in Refs. 2 and 3 that this is $O(R^{-3/8})$, just a fraction smaller than the longitudinal pressure variation $O(R^{-1/4})$. Neglecting this term is justified in the limit as $R \rightarrow \infty$, but may well lead to numerical inaccuracy at $R = 10^5$.

Fourth, the pressure plotted in Fig. 6 of Ref. 1 shows a

singularity at $x = 1$, and so dp/dx is seriously in error if x is very close to 1. The solutions to Eqs. (1-3) of Ref. 1 presumably require that the pressure be known at "the edge of Region I"; however, the pertinent details are not discussed. Furthermore, the numerical work of Ref. 1 would have benefited from making use of an asymptotic analysis of the outer region of influence of the trailing edge. Conditions at infinity are notoriously difficult to handle numerically, and the discrepancy between the results of Ref. 1 and those of Goldstein, when $x = 2$, should be regarded with caution.

Finally, Fig. 6 of Ref. 1 is in qualitative agreement with the predictions of Refs. 2 and 3, in that a favorable pressure gradient is established for $x < 1$, an adverse pressure gradient at $x = 1^+$, and a favorable pressure gradient as $(x-1)R^{3/8} \rightarrow \infty$. Fig. 6 shows that the displacement effect leads to large dc_p/dx for $\Delta x \approx 0.2$ (say) whereas Fig. 10 shows significant changes in skin friction for $\Delta x \approx 0.0005$. While the dependence on R is not clearly established by these results, the curves do show unequivocally the existence of two length scales, again qualitatively in agreement with the results of Refs. 2 and 3. The latter references do not contend that the region $O(R^{-3/4}L)$ is irrelevant to the structure of the flow, but rather that the dominant correction occurs on the larger scale, and that even for $R = 10^5$ (say) the largest effects are found in a qualitatively correct way by the asymptotic analysis given. For example, the correction to the drag coefficient on the larger scale is $O(R^{-7/8})$, whereas effects on the smaller scale lead to a change $O(R^{-5/4})$. The correction may be calculated either by integrating skin friction along the plate or by calculating a momentum deficit in the wake. For consistency the eigenfunction proposed in Eq. 5.10 of Ref. 2 is required.

References

- 1 Schneider, L. I. and Denny, V. E., "Evolution of the Laminar Wake behind a Flat Plate and Its Upstream Influence," *AIAA Journal*, Vol. 9, No. 4, April 1971, pp. 655-660.
- 2 Stewartson, K., "On the Flow Near the Trailing Edge of a Flat Plate II," *Mathematika*, Vol. 16, Pt. 1, June 1969, pp. 106-121.
- 3 Messiter, A. F., "Boundary Layer Flow Near the Trailing Edge of a Flat Plate," *SIAM Journal of Applied Mathematics*, Vol. 18, No. 1, Jan. 1970, pp. 241-257.
- 4 Stewartson, K., "On the Flow Near the Trailing Edge of a Flat Plate," *Proceedings of the Royal Society*, Vol. A306, 1968, pp. 275-290.
- 5 Hakkinen, R. J. and O'Neil, E. J., "Merging of Uniform Shear Flows at a Trailing Edge," *AIAA Journal*, Vol. 7, No. 3, March 1969, pp. 523-525.

Reply by Author to A. F. Messiter and K. Stewartson

V. E. DENNY

University of California, Los Angeles, Calif.

THE authors thank Professors Messiter and Stewartson for their interest in and critical evaluation of the results in Ref. 1. Unfortunately, a portion of their comment is based on an error in the analysis section of Ref. 1 wherein the term $v \partial u/\partial y$ was omitted from Eq. (7). As inferred implicitly in the discussion of the solution method,¹ the potential flow problem was solved "exactly".² Whether or not this was necessary is perhaps debat-

Received July 30, 1971.

* Professor, Department of Aerospace Engineering, Member AIAA.

† Visiting Professor, Department of Aeronautical and Astronautical Engineering.

Received December 20, 1971.

* Assistant Professor, Department of Energy and Kinetics.




Fig. 1 Relative slopes of the (effective) slender body.

able; however, as illustrated in Fig. 1 the normalized slopes of the (effective) slender body are somewhat larger than $O(1)$ behind the trailing edge. Thus, the authors decided, within the framework of

an "exact" treatment of the over-all problem, to forego linearized airfoil theory.

With respect to omission of cross-streamwise pressure variations in the "Region II" analysis, this was done only after it was determined that resultant errors in calculating the displacement thickness δ^* were no greater than those involved in solving the airfoil problem. (In principle, the "Region I" problem could have been solved with $y \rightarrow \infty$; i.e., $t \equiv [1 - \exp(-R^{1/2}y)] \rightarrow 1$.) In this respect, the authors reiterate that a primary objective of Ref. 1 was to demonstrate numerical techniques for solving problems of this kind.

References

¹ Schneider, L. I. and Denny, V. E., "Evolution of the Laminar Wake Behind a Flat Plate and Its Upstream Influence," *AIAA Journal*, Vol. 9, No. 4, April 1971, pp. 655-660.

² Giesing, J. P., "Solution of the Flow Field about One or More Airfoils of Arbitrary Shape in Uniform and Nonuniform Flows by the Douglas Neumann Method," LB31946, May 1968, Douglas Aircraft Co., Los Angeles, Calif.

Errata

Errata: "Vibrations of Pressurized Orthotropic Shells"

CLIVE L. DYM

Carnegie-Mellon University, Pittsburgh, Pa.

[AIAA J. 9, 1201-1203 (1971)]

EQUATIONS (4) should read

$$C_{11} \frac{\partial^2 u}{\partial x^2} + G_{12} \frac{\partial^2 u}{\partial y^2} + (C_{12} + G_{12}) \frac{\partial^2 v}{\partial x \partial y} - C_{12} \frac{1}{R} \frac{\partial w}{\partial x} = \rho \frac{\partial^2 u}{\partial t^2} \quad (4a)$$

$$(C_{12} + G_{12}) \frac{\partial^2 u}{\partial x \partial y} + G_{12} \frac{\partial^2 v}{\partial x^2} + C_{22} \frac{\partial^2 v}{\partial y^2} - C_{22} \frac{1}{R} \frac{\partial w}{\partial y} = \rho \frac{\partial^2 v}{\partial t^2} \quad (4b)$$

$$-\frac{1}{12} \left(\frac{h}{R} \right)^2 \left[C_{11} \frac{\partial^4 w}{\partial x^4} + (2C_{12} + 4G_{12}) \frac{\partial^4 w}{\partial x^2 \partial y^2} + C_{22} \frac{\partial^4 w}{\partial y^4} \right] + C_{12} \frac{1}{R} \frac{\partial u}{\partial x} + C_{22} \frac{1}{R} \left(\frac{\partial v}{\partial y} - \frac{w}{R} \right) + \tau_{xx}^0 \frac{\partial^2 w}{\partial x^2} + \tau_{yy}^0 \frac{\partial^2 w}{\partial y^2} = \rho \frac{\partial^2 w}{\partial t^2} \quad (4c)$$

The second of Eqs. (5) should read

$$v(x, y, t) = B_{mn} \sin(m\pi x/L) \sin(ny/R) \cos \omega t$$

Erratum: "A Nonlinear Method Applied to Variable-Thickness Plates"

TSAI-CHEN SOONG

Xerox Corporation, Rochester, N.Y.

[AIAA J. 10, 148-154 (1972)]

ON p. 14, the following heading and paragraph were inadvertently omitted.

Derivations of Equations

The method will be illustrated by solving the large-deflection problem of a rectangular plate of variable thickness under uniform lateral pressure. The nondimensional large-deflection plate equations of von Kármán's, corresponding to compatibility and equilibrium in the lateral direction, can be written as

$$L_f(x, y) = \nabla^4 F - t(w_{,xy}^2 - w_{,xx} w_{,yy} - \Omega(y) w_{,xx}) = 0 \quad (1)$$