

This scattering kernel for argon reflected from a silver surface is now used to calculate the effect of incident angle and velocity on reflected density as given in Eq. (1). Figure 1a gives the theoretical results for the experimental conditions of Calia and Oman³ mentioned above where θ and ϕ are in-plane and out-of-plane reflection angles as illustrated in Fig. 1 of Ref. 1. Figure 1b illustrates the effect of incident angle. The ratio of maximum density in the plane of incidence and that 30° out-of-plane is approximately 29 and 8.2 for incident angles of -50° and -20°, respectively. Figures 1c and 1d show the effect of incident energy. The ratio of maximum density in plane to that 30° out of plane for incident energies of 2.14 eV, 0.21 eV and 0.02 eV are 29, 1.78 and 1.31, respectively. The relative amount of out-of-plane scattering varies by over a factor of 20 with only an order-of-magnitude change in incident velocity, and a factor of over 3 with a change in incident angle of 30° for fixed incident speed. These results certainly disprove Oman's statement that "In Kinslow's model there can be no variation in relative amount of out-of-plane scattering with energy or incident angle,..."

Notice also that the reflected density lobe is subspecular for the lower incident energy, becomes specular and then supraspecular as incident energy is increased. Also it should be noted that at low incident energy the reflected distribution is broad and diffuse, while at higher energies it is narrow and more specular in nature.

All of the aforementioned characteristics of the proposed scattering kernel are in agreement with the experimental observations. This example refutes the statement of Oman

that "Kinslow's model would appear to require that there be no significant memory of the incident energy,..." The product scattering kernel as developed in Ref. 1 is certainly not the ultimate kernel. However, based upon comparison with experimental results from the thermal through structural scattering regime, it appears to be the best analytical model so far developed. Without question the model proposed by Oman in Eq. (1) of his comment is more general and includes the present model. However, there are doubts as to whether or not it could be developed into a useful model.

To ascertain whether or not the assumption of a product of individual kernels is justified, it is necessary to have experimental determination of the scattering kernel, not just moments such as intensity, density, or velocity as has been previously presented. I know of no results either experimental or from numerical modeling which give the basic scattering kernel. I would agree with Oman in his statement that absolute measurements are needed both in and out of the plane of incidence.

References

- ¹Kinslow, M., "A Mathematical Description of Gas-Surface Interactions Based on Reciprocity," *AIAA Journal*, Vol. 14, Oct. 1976, pp. 1358-1361.
- ²Moran, J. P., "Experiments on Scattering of Mono-Energetic Argon Beams of Heated Platinum," Ph.D. dissertation, Feb. 1968, Dept. of Aeronautics and Astronautics, MIT, Cambridge, Mass.
- ³Calia, V.S. and Oman, R.A., "Scattering Cross-Section Measurements for Epithermal AR on Ag (111) Surfaces," *Journal of Chemical Physics*, Vol. 52, 15 June 1970, pp. 6184-6188.

Errata

Prediction of Turbulent Boundary Layers at Low Reynolds Numbers

Richard H. Pletcher
Iowa State University, Ames, Iowa

AIAA J. 14 696-698 (1976)

IN the second line above Eq. (7) on page 698, $\ell/\delta < 0.089$ should be replaced by $\ell/\delta > 0.089$.

Received Dec. 20, 1976.
Index category: Boundary Layers and Convective Heat Transfer—Turbulent.

Quasilinearization and Optimal Control Problems with Control Bounds

B.P. Yeo
University of Singapore, Singapore

[AIAA J., 14, 963-966 (1976)]

EQUATION (2) should be replaced by

$$u_{\min} \leq u \leq u_{\max} \quad (2)$$

The original version of Eq. (2) resulted from a clerical error.

Received Dec. 17, 1976.
Index category: Navigation, Control, and Guidance Theory.

Quasi-Steady Gas Phase Assumption for a Burning Droplet

Josette Bellan and Martin Summerfield
Princeton University, Princeton, N. J.

[AIAA J., 14, 973-975 (1976)]

THE definition of τ_g should be

$$\tau_g = \frac{D_g}{(u_g)^2_{\text{ref}}}$$

and the convection term in Eq. (1) should read

$$\frac{u_g}{R\beta} \frac{\partial \theta_g}{\partial y}$$

Equation (2) should read

$$\frac{u_g}{R\beta} > > 1$$

and Eq. (4) should read

$$\tau_p > > \tau_d(p/334)$$

The relationship after Eq. (5) should read

$$\frac{D_g}{[R(t=0)]^2} = \frac{1}{2\tau_d[\ln(1+B)](\rho_g/\rho_d)}$$

Equation (6) should read $\tau_p > > (\tau_d/3) \times 10^{-3}$.
Equation (7) should read $\tau_p > > (\tau_d p/3) \times 10^{-3}$.

Received Nov. 3, 1976.
Index category: Combustion in Heterogeneous Media.