

- Rooks, R. E., V. Julka, M. F. Doherty, and M. F. Malone, "Structure of Distillation Regions for Multicomponent Azeotropic Mixtures," *AIChE J.*, **44**, 1382 (1998).
- Rousseau, R. W., *Crystallization, Kirk-Othmer Encyclopedia of Chemical Technology*, 4th ed., Vol. 7, Wiley Interscience, New York, p. 683 (1994).
- Safrit, B. T., and A. W. Westerberg, "Algorithm for Generating the Distillation Regions for Azeotropic Multicomponent Mixtures," *Ind. Eng. Chem. Res.*, **36**, 1827 (1997).
- Samant, K. D., and K. M. Ng, "Synthesis of Extractive Reaction Processes," *AIChE J.*, **44**, 1363 (1998).
- Takano, K., R. Gani, T. Ishikawa, and P. Kolar, "Integrated System for Design and Analysis of Industrial Processes with Electrolyte System," Paper presented in Escape '99, Budapest, Hungary (1999).
- Tare, J. P., and M. R. Chivate, "Separation of Close Boiling Isomers by Adductive and Extractive Crystallization," *AIChE Symp. Ser. No. 153*, **72**, 95 (1976).
- Thomsen, K., P. Rasmussen, and R. Gani, "Simulation and Optimization of Fractional Crystallization Processes," *Chem. Eng. Sci.*, **53**, 1551 (1998).
- Ung, S., and M. F. Doherty, "Vapor-Liquid Phase Equilibrium Systems with Multiple Chemical Reactions," *Chem. Eng. Sci.*, **50**, 23 (1995).
- Wibowo, C., and K. M. Ng, "Synthesis of Bulk Solids Processing Systems," *AIChE J.*, **45**, 1629 (1999).
- Weingaertner, D. A., S. Lynn, and D. N. Hanson, "Extractive Crystallization of Salts from Concentrated Aqueous Solution," *Ind. Eng. Chem. Res.*, **30**, 490 (1991).

Manuscript received Sept. 17, 1999, and revision received Jan. 28, 2000.

Correction

In the article titled "Corrected Horváth-Kawazoe Equations for Pore-Size Distribution" by Salil U. Rege and Ralph T. Yang (pp. 734-750, April 2000) the following corrections are made:

- On page 736, the equation for the adsorbate dispersion A_A as given by the Kirkwood-Muller formulae (Eq. 4b) was incorrectly given as:

$$A_A = \frac{3}{2} mc^2 \alpha_s \alpha_A$$

The correct formula to be used is as follows:

$$A_A = \frac{3}{2} mc^2 \alpha_A \chi_A$$

- The diameter of the nitrogen adsorbate molecule (Table 1, page 737) used in the cylindrical pore model is 3.72 Å and not 372 Å, as erroneously published.