

Synthesis of Amino Acid Conjugates and Further Derivatives of 3 α -Hydroxylup-20(29)ene-23,28-dioic acid

Tran Van Loc and Tran Van Sung

Hanoi, Institute of Chemistry, National Centre for Natural Science and Technology of Vietnam

C. Kamperdick and G. Adam*

Halle/S., Institute of Plant Biochemistry

Received May 18th, 1999, respectively November 10th, 1999


Keywords: Amino acids, Drug research, Terpenoids, Triterpenes, 3 α -Hydroxylup-20(29)ene-23,28-dioic acid

Abstract. Triterpenes of betulinic acid type exhibit many interesting biological activities. Therefore a series of new 3 α -hydroxylup-20(29)ene-23,28-dioic acid derivatives **2a–22** with putative pharmacological activities were synthesized. As starting compounds 3 α -hydroxylup-20(29)ene-23,28-dioic acid (**1a**), isolated from *Schefflera octophylla*, or its 3-*O*-acetyl derivative **1b** were used. Mono- and diesters (**2a–b** from **1a**, and **4d** from **4c**) were prepared with CH₂N₂. Oxidation of the isopropenyl side chain with OsO₄ yielded the 20,29-diols (**4a–b** from **1b**, and **19** from **17**), which were in the case of **4b** further transformed to the 29-norketones **8a–**

b. Oxidation of the isopropenyl side chain with *m*-chloroperbenzoic acid afforded the 20,29-epoxide **12** (from **1b**) and the 29-aldehydes and α -hydroxy aldehydes (**13a–c** from **2a**, **14a–c** from **2b**, and **16a–c** from **15a**). Ring A was modified by a tosylation–elimination sequence using *p*-TsCl/NaOAc, which afforded diolefin **15a** (from **2a**) with $\Delta^{2,20(29)}$ double bonds or 23-nor- $\Delta^{3,20(29)}$ diolefin **17** (from **1a**). Compounds **4b**, **4c**, and **8a** were coupled with L-methionin, L-phenylalanin, L-alanin, L-serin, and L-glutaminic acid *via* amide bonds at positions **23** and **28** to afford the amino acid conjugates **5a–7b** and **9a–11**.

Triterpenes of the betulinic acid type are getting more and more importance because of their interesting pharmacological activities. Especially they display anti-inflammatory [1], antitumor [2], leishmanicidal [3], and antimalarial [4] activities. Most interesting is the fact that betulinic acid, dihydrobetulinic acid, and platanic acid also exhibit anti-HIV activity [5]. Further investigations showed that some amides of betulinic acid are potent HIV inhibitors. The presence of an amide function within the side chain was found to be important for optimal activity. Minor modifications in the lupane backbone strongly affected activity [6, 7].

3 α -Hydroxylup-20(29)ene-23,28-dioic acid (**1a**) is a rare natural product isolated in surprisingly high yield (7%) from leaves of the Vietnamese plant *Schefflera octophylla* (Araliaceae) [8]. It is structurally closely related to betulinic acid. Therefore, we have recently synthesized a series of amino acid conjugates of **1a** with retainment of the isopropenyl side chain and of the backbone [9]. In continuation of these studies, we now report the synthesis of further derivatives of **1a** by oxidation of the isopropenyl side chain, preparation of various mono- and diesters as well as modifications at ring A. Selected derivatives were transformed to amino acid conjugates.




Comp.	R ¹	R ²	R ³	Formula	m.w.
1a	H	OH	OH	C ₃₀ H ₄₆ O ₅	486.7
1b	Ac	OH	OH	C ₃₂ H ₄₈ O ₆	528.7
2a	H	OMe	OH	C ₃₁ H ₄₈ O ₅	500.7
2b	H	OMe	OMe	C ₃₂ H ₅₀ O ₅	514.7
3	H	–NH–cyclohexyl	OH	C ₃₆ H ₅₇ NO ₄	567.9

Oxidation of the Isopropenyl Side Chain

A 20,29-dihydroxy functionality was generated by reaction of the 3 α -acetyl derivative **1b** [8] with OsO₄ leading to the two diastereomeric diols **4a** and **4b**. Subsequent oxidation with KIO₄ yielded the 29-norketone


8a. Alkaline hydrolysis of **4b** and **8a** gave the corresponding 3 α -hydroxy compounds **4c** and **8b**, respectively. Oxidation of the isopropenyl side chain of the 3 α -acetoxy compound **1b** with *m*-chloroperbenzoic acid furnished the 20,29-epoxide **12** as pure diastereomer (configuration at C-20 not determined). On the other hand the 3 α -hydroxy compounds **2a**, **2b**, and **15a** (synthesis see below) afforded as main products their corresponding 29-aldehydes as mixtures of the C-20 epimers **13a/b**, **14a/b**, and **16a/b**, respectively, separated by chromatography in the case of the pair **16a/b**. As minor products the corresponding α -hydroxy aldehydes **13c**, **14c**, and **16c**, respectively, were obtained as only one isomer.



4a-d, 5a,b, 6a,b, 7a,b


Comp.	R ¹	R ²	R ³	R ⁴	R ⁵	Formula	m.w.
4a ^{a)}	Ac	OH	OH	H	H	C ₃₂ H ₅₀ O ₈	562.7
4b ^{a)}	Ac	OH	OH	H	H	C ₃₂ H ₅₀ O ₈	562.7
4c	H	OH	OH	H	H	C ₃₀ H ₄₈ O ₇	520.7
4d	H	OMe	OMe	H	H	C ₃₂ H ₅₂ O ₇	548.8
5a	Ac	L-Met-OMe	L-Met-OMe	R ⁴ -R ⁵ = -CO-	H	C ₄₅ H ₇₀ N ₂ O ₁₁ S ₂	879.2
5b	H	L-Met-OH	L-Met-OH	H	H	C ₄₀ H ₆₆ N ₂ O ₉ S ₂	783.1
6a	Ac	L-Phe-OEt	L-Phe-OEt	R ⁴ -R ⁵ = -CO-	H	C ₅₅ H ₇₄ N ₂ O ₁₁	939.2
6b	H	L-Phe-OH	L-Phe-OH	H	H	C ₄₈ H ₆₆ N ₂ O ₉	815.1
7a	CO-CO-L-Ala-OMe	L-Ala-OMe	L-Ala-OMe	R ⁴ -R ⁵ = -CO-	H	C ₄₅ H ₆₇ N ₃ O ₁₄	874.0
7b	H	L-Ala-OH	L-Ala-OH	H	H	C ₃₆ H ₅₈ N ₂ O ₉	662.9

^{a)} Configuration at C-20 not assigned



8a-c, 9a,b, 10, 11

Comp	R ¹	R ²	R ³	Formula	m.w.
8a	Ac	OH	OH	C ₃₁ H ₄₆ O ₇	530.7
8b	H	OH	OH	C ₂₉ H ₄₄ O ₆	488.7
8c	H	OMe	OMe	C ₃₁ H ₄₈ O ₆	516.7
9a	Ac	L-Ala-OMe	L-Ala-OMe	C ₃₉ H ₆₀ N ₂ O ₉	700.9
9b	H	L-Ala-OH	L-Ala-OH	C ₃₅ H ₅₄ N ₂ O ₈	630.8
10	Ac	L-Ser-OMe	L-Ser-OMe	C ₃₉ H ₆₀ N ₂ O ₁₁	732.9
11	Ac	L-Glu(OMe) ₂	L-Glu(OMe) ₂	C ₄₅ H ₆₈ N ₂ O ₁₃	845.0




12, 13a-c, 14a-c

Comp.	R ¹	R ²	R ³	R ⁴	Formula	m.w.
12 ^{a)}	Ac	OH	OH	1,2-epoxyisopropyl	C ₃₂ H ₄₈ O ₇	544.7
13a/b ^{b)}	H	OMe	OH	-C(H,CH ₃)-CHO	C ₃₁ H ₄₆ O ₆	516.7
13c ^{a)}	H	OMe	OH	-C(OH,CH ₃)-CHO	C ₃₁ H ₄₈ O ₇	532.7
14a/b ^{b)}	H	OMe	OMe	-C(H,CH ₃)-CHO	C ₃₂ H ₅₀ O ₆	530.7
14c ^{a)}	H	OMe	OMe	-C(OH,CH ₃)-CHO	C ₃₂ H ₅₀ O ₇	546.7

^{a)} Configuration at C-20 not determined ^{b)} Mixture of C-20 epimers

Mono- and Diesters

The monomethyl ester **2a** and dimethylester **2b** [8] of **1a** were prepared by reaction with equimolar or excess amounts of CH₂N₂. The location of the ester group in **2a** was determined by comparison of the carbon signals of **2a** and **1a** in CD₃OD, which showed identical signals ($\Delta\delta < 0.1$ ppm) for all carbons except for C-3 ($\Delta\delta + 0.4$), C-4 ($\Delta\delta + 0.7$), C-5 ($\Delta\delta + 0.2$) and C-6 ($\Delta\delta + 0.2$), thus proving **2a** to be the 23-monomethyl ester of **1a**. From the 20,29-dihydroxy derivative **4c**, too, the diester **4d** was prepared, which was further oxidized with KIO₄ to give 29-norketone **8c**.



Comp.	R ¹	R ²	R ³	Formula	m.w.
15a	OMe	OH	isopropenyl	C ₃₁ H ₄₆ O ₄	482.7
15b	OH	OH	isopropenyl	C ₃₀ H ₄₄ O ₄	468.7
16a ^{a)}	OMe	OH	–C(H,CH ₃)–CHO	C ₃₁ H ₄₆ O ₅	498.7
16b ^{a)}	OMe	OH	–C(H,CH ₃)–CHO	C ₃₁ H ₄₆ O ₅	498.7
16c ^{a)}	OMe	OH	–C(OH,CH ₃)–CHO	C ₃₁ H ₄₆ O ₆	514.7


^{a)} Configuration at C-20 not assigned

Modifications of Ring A

Starting from 23-methylester **2a**, a double bond was introduced at ring A *via* a tosylation–elimination sequence using *p*-TsCl and NaOAc. The obtained $\Delta^{2,20(29)}$ diolefin **15a** was hydrolyzed to give the free acid **15b**. The presence of the newly introduced Δ^2 double bond was revealed by the carbon spectra, which show two new olefinic methin carbons at δ 125.7 and 130.4 and a significant downfield shift of C-1 (δ 40.7).

Starting from the free diacid **1a**, the tosylation–elimination sequence interestingly yielded the 23-nor- $\Delta^{3,20(29)}$ diolefin **17** with decarboxylation of the 23-carboxy group. OsO₄-Hydroxylation of **17** afforded a mixture of 3 α ,4 α -diol **18** and 3 α ,4 α ,20 ξ ,29-tetrol **19**, the latter one isolated as pure diastereomer. Oxidative diol cleavage of **18** with KIO₄ yielded the 3,4-*seco* compound **20**.


The 23-nor-3-ketone **21** was prepared by oxidation of **1a** with Jones reagent.



Comp.	Formula	m.w.
17	C ₂₉ H ₄₄ O ₂	424.7


Synthesis of Amino Acid Conjugates and Further Nitrogen Containing Compounds

The diacids **4b**, **4c**, and **8a** were transformed to the diamino acid esters **5a** and **6a** (from **4b**), **7a** (from **4c**) and **9a**, **10**, and **11** (from **8a**) by activation of the carboxyl group with oxalyl chloride followed by reaction with amino acid ester hydrochlorides in the presence of triethylamine [9]. Hereby, the 20,29-dihydroxy functions in **4b** and **4c** were transformed to oxo-1,3-dioxolan side chains (compounds **5a**, **6a**, and **7a**) at C-19.




Comp.	R	Formula	m.w.
18	isopropenyl	C ₂₉ H ₄₆ O ₄	458.7
19 ^{a)}	1,2-dihydroxyisopropyl	C ₂₉ H ₄₈ O ₆	492.7

^{a)} Configuration at C-20 not assigned



Comp.	Formula	m.w.
20	C ₂₉ H ₄₄ O ₄	456.7




Comp.	Formula	m.w.
21	C ₂₉ H ₄₄ O ₃	440.7

Because of the unprotected 3 α -hydroxy group **4c** gave the 3-*O*-oxamoyl compound **7a** with three conjugated amino acids. Subsequent alkaline hydrolysis of the esters yielded the corresponding diamino acids **5b**, **6b**, **7b**, and **9b** with simultaneous regeneration of the 3 α -hydroxy and the 20,29-dihydroxy functionalities. In the

case of the methyl esters of the diserin and diglutamic acid conjugates **10** and **11**, their corresponding free acid conjugates could not be obtained pure, probably due to further intra- and intermolecular reactions.

Upon reaction of diacid **1a** with dicyclohexylcarbodiimide, the 28-carboxylic function remained untouched, leading to a mixture of cyclohexyl amide **3** and imino-oxo-1,3-dioxane **22**.



The structures of all new compounds were confirmed by MS and NMR spectroscopy (¹H, ¹³C, APT). The compounds are subjected to an extensive biological screening.

We thank Dr. J. Schmidt for the mass spectra and Dr. H. Ripperger for discussions. We are indebted to the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie, Bonn, and the Volkswagen-Stiftung, Hannover, for financial support.

Experimental

m.p. were measured with a Zeiss heating stage microscope (Boëtius) and are uncorrected. EI MS were recorded at 70 eV on a AMD 402 spectrometer. ESI MS were measured with a Finnigan TSQ 700 spectrometer. ¹H and ¹³C NMR spectra were taken on a Varian Gemini 300 spectrometer (300/75 MHz). Chemical shifts are given in ppm (δ -scale), coupling constants (*J*) in Hz. ¹H-chemical shifts were referenced to TMS, ¹³C-chemical shifts were referenced to CDCl₃ (77.0), CD₃OD (49.0) or C₅D₅N (123.5). IR-spectra were taken on a Bruker IFS28 spectrometer. Optical rotations were measured on a Jasco DIP 1000 Polarimeter, cell length 100 mm. Flash-chromatography was carried out on silica gel 60, 40–63 μ m (Merck) using a glass column equipment for normal pressure and 0.5 bar N₂.

OsO₄-Dihydroxylation of **1b** and **17** (General Procedures)

A solution of 529 mg (1 mmol) 3 α -acetoxy-20(29)-ene-23,28-dioic acid (**1b**), 176 mg (1.5 mmol) *N*-methylmorpholine-*N*-oxide and 20 mg (0.08 mmol) of OsO₄ in 20 ml acetone/H₂O

(8:2) was stirred at *r.t.* for 48 h [10]. 10 mg (0.1 mmol) NaHSO₃ and 15 ml H₂O were added, the solution filtered and the filtrate neutralized with 1N H₂SO₄. After evaporation of the organic solvent under reduced pressure, the aqueous layer was extracted with EtOAc, dried with Na₂SO₄, and concentrated to dryness. The residue was chromatographed on silica gel with CHCl₃/EtOAc (95:5) and yielded the diastereomeric compounds **4a** (27%) and **4b** (48%). Analogously, **18** (30%) and **19** (15%) were prepared from **17**.

Oxidation of **4b**, **4d**, and **18** with KIO₄

A solution of **4b**, **4d** or **18** (1 mmol) and 690 mg (3 mmol) KIO₄ in 20 ml MeOH/H₂O (8:2) was stirred at *r.t.* for 20 h. The mixture was concentrated under reduced pressure and purified by flash chromatography (silica gel, CHCl₃/MeOH 95:5) to afford **8a** (96%), **8c** (94%) or **20** (90%), respectively.

Oxidation of **1b**, **2a**, **2b**, and **15a** with *m*-Chloroperbenzoic Acid

A solution of 529 mg (1 mmol, dissolved in Et₂O) **1b** and 432 mg (2.5 mmol) *m*-chloroperbenzoic acid in 10 ml CH₂Cl₂ was stirred at 25 °C for 24 h. The resulting reaction mixture was washed twice with H₂O, dried with Na₂SO₄ and concentrated under reduced pressure. Flash chromatography (silica gel, *n*-hexane/EtOAc 75:25) afforded **12** (82%). Analogously, **13a/b** (50%) and **13c** (20%) were prepared from **2a**. **14a/b** (60%), and **14c** (20%) were obtained from **2b**. **15a** afforded **16a** (35%), **16b** (25%), and **16c** (15%).

Alkaline Hydrolysis of Compounds **4b**, **8a**, and **15a**

Compound **4b**, **8a** or **15a** (1 mmol) was hydrolyzed with 5% KOH in MeOH (10 ml) by stirring at *r.t.* for 20 h. H₂O (30 ml) was added, the mixture acidified to pH 4 with HCl (5%) and extracted with EtOAc. The organic layer was washed with H₂O, dried with Na₂SO₄ and the solvent was evaporated under reduced pressure to afford **4c** (92%), **8b** (95%) or **15b** (90%).

Preparation of Methyl Esters **2a**, **2b**, and **4d**

487 mg (1 mmol) of compound **1a** in MeOH (20 ml) were stirred for 1 h at 25 °C with CH₂N₂ (1 mmol, dissolved in Et₂O). The solvent was removed under reduced pressure and the residue purified by flash chromatography (*n*-hexane/EtOAc 8:2) to yield **2a** (72%). Analogously, **2b** (98%) and **4d** (92%) were prepared from **1a** and **4c**, respectively, but using 4 mmol CH₂N₂.

Two-Step Tosylation/Elimination of **1a** and **2a**

To a solution of **1a** or **2a** (1 mmol) in pyridine (5 ml) 477 mg (2.5 mmol) *p*-toluenesulfonylchloride were added and the solution stirred for 24 h at *r.t.* After addition of H₂O (15 ml) a white solid precipitated, which was filtered off and used in the next step without purification. It was dissolved in DMF (30 ml), NaOAc was added and the mixture heated at 120 °C for 6 h. After evaporation of the solvent, the residue was treated with H₂O (20 ml) and extracted with EtOAc. The organic layer was dried with Na₂SO₄, filtered and concentrated under

reduced pressure. The product was chromatographed on silica gel (*n*-hexane/EtOAc 8:2) to give **17** (92%) or **15a** (82%).

Jones Oxidation of **1a**

487 mg (1 mmol) of **1a** in acetone (20 ml) were stirred with 1 ml Jones reagent (2.8 g K₂Cr₂O₇ in 2.3 ml conc. H₂SO₄ and 10 ml H₂O) in an ice bath for 1 minute and then for 30 minutes at *r.t.* After addition of ice, the mixture was extracted with CHCl₃. The organic layer was washed with H₂O and dilute NaHCO₃ solution, dried with Na₂SO₄ and concentrated. The residue was purified by flash chromatography (*n*-hexane/EtOAc 7:3) to afford **21** (76%).

Preparation of the Amino Acid Conjugates **5a**, **5b**, **6a**, **6b**, **7a**, **7b**, **9a**, **9b**, **10**, and **11**

A solution of 563 mg (1 mmol) of **4b** and 1.02 g oxalyl chloride (8 mmol) in benzene (20 ml) was stirred at *r.t.* for 24 h and concentrated under reduced pressure. A solution of the residue, amino acid methyl or ethylester hydrochloride (5 mmol) and 1.01 g (10 mmol) triethylamine in CH₂Cl₂ (30 ml) was stirred at *r.t.* for 20 h. The resulting reaction mixture was washed twice with H₂O, dried with Na₂SO₄ and concentrated under reduced pressure. Flash chromatography (silica gel, *n*-hexane/EtOAc) furnished the ester derivatives **5a** (86%) and **6a** (90%). The esters were hydrolyzed (same procedure as described for hydrolysis of **4b**) to give **5b** (88%) and **6b** (90%).

Analogously, **7a** (68%) and **7b** (86%) were prepared from **4c**. **9a** (96%), **9b** (94%), **10** (98%), and **11** (87%) were prepared from **8a**.

Reaction of **1a** with Dicyclohexylcarbodiimide

A solution of 487 mg (1 mmol) of **1a** in tetrahydrofuran (50 ml) was cooled to –5 °C. Then a concentrated solution of 206 mg (1 mmol) dicyclohexylcarbodiimide was added. After 10 h at –5 °C and 15 h at *r.t.*, CH₂Cl₂ (20 ml) was added and the solution washed with 5% HCl, 5% NaHCO₃ and H₂O and dried with Na₂SO₄. The solvent was evaporated under reduced pressure and the residue chromatographed on silica gel with CHCl₃/EtOAc to afford **3** (18%) and **22** (30%).

Physical Data of the Compounds Prepared [11, 12]

3 α -Hydroxy-lup-20(29)-ene-23,28-dioic acid 23-methyl ester (2a**)**

m.p. 125–126 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ –1.6 (CHCl₃, *c* = 1.0). – ¹H NMR (CDCl₃): δ /ppm = 3.75 (br s, H-3), 3.70 (s, OMe), 1.16 (s, H₃-24). – ¹³C NMR (CD₃OD): δ /ppm = 180.1 (C-28), 178.7 (C-23), 74.2 (C-3), 57.5 (C-17), 52.0 (OMe), 48.5 (C-19), 45.2 (C-5), 22.5 (C-6). – EIMS: *m/z* (%) = 500 [M]⁺ (22), 482 (8), 468 (8), 454 (14), 248 (37), 233 (79), 201 (55), 187 (80), 173 (99), 119 (100), 107 (81), 95 (77).

3 α -Hydroxy-lup-20(29)-ene-23,28-dioic acid dimethyl ester (2b**)**

m.p. 83–84 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ –4.0 (CHCl₃, *c* = 1.0). – ¹H NMR (CDCl₃): δ /ppm = 4.70 (m, *J* = 2.2 Hz, H-29^B), 4.60 (q, *J* = 1.3 Hz, H-29^A), 3.73 (br s, H-3), 3.70, 3.66 (each s, OMe), 3.00 (m, H-19), 1.16 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 178.1 (C-23), 176.6 (C-28), 72.4 (C-3),

56.6 (C-17), 51.8, 51.1 (OMe) 24.9 (C-2). – EIMS: *m/z* (%) = 514 [M]⁺ (56), 496 (10), 482 (11), 454 (26), 264 (75), 249 (77), 233 (59), 203 (57), 189 (100), 175 (66), 107 (53), 93 (51), 81 (49).

3 α -Hydroxy-lup-20(29)-ene-23,28-dioic acid 23-cyclohexylamide (3**)**

m.p. 186–187 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ +16.6 (CHCl₃, *c* = 1.0). – ¹H NMR (CDCl₃): δ /ppm = 3.79 (m, CHN), 3.64 (br s, H-3), 4.74 (d, *J* = 1.7 Hz, H-29^B), 4.61 (br s, H-29^A), 3.01 (td, *J* = 10.4 and 4.1 Hz, H-19), 1.69 (s, H₃-30), 1.13 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 181.7 (C-28), 177.8 (C-23), 71.8 (C-3), 56.3 (C-17), 46.9 (C-19), 48.5, 32.5, 32.1, 24.9, 24.7 (cyclohexyl), 19.2 (C-30). – EIMS: *m/z* (%) = 567 [M]⁺ (38), 549 (31), 534 (15), 424 (25), 259 (16), 194 (100), 155 (52).

3 α -Acetoxy-20 ξ ,29-dihydroxylupane-23,28-dioic acid (4a**)**

m.p. 206–208 °C (MeOH/CHCl₃). – $[\alpha]_D^{24}$ –36.5 (MeOH, *c* = 2.0). – IR: ν /cm^{–1} (KBr) = 3 446 (m, br), 2 947 (s), 2 872 (w), 1 718 (s), 1 457 (m), 1 378 (m), 1 260 (s), 1 025 (m) 988 (w). – ¹H NMR (CD₃OD): δ /ppm = 4.92 (br s, H-3), 3.45 (d, *J* = 11.0 Hz, H-29^B), 3.36 (d, *J* = 11.0 Hz, H-29^A), 1.04 (s, H₃-30). – ¹³C NMR (CD₃OD): δ /ppm = 180.4 (C-28), 179.0 (C-23), 77.0 (C-3), 76.1 (C-20), 71.1 (C-29), 59.5 (C-17), 45.6 (C-19), 23.3 (C-2), 19.5 (C-30). – Negative ESI MS: *m/z* (%) = 561.6 (100) [M–H][–].

C-20 Epimer **4b**

m.p. 197–198 °C (MeOH/CHCl₃). – $[\alpha]_D^{24}$ –7.1 (MeOH, *c* = 2.0). – IR: ν /cm^{–1} (KBr) = 3 446 (m, br), 2 947 (s), 2 878 (w), 1 718 (s), 1 457 (m), 1 378 (m), 1 259 (s), 1 027 (m), 989 (w). – ¹H NMR (CD₃OD): δ /ppm = 3.41 (s, H₂-29), 1.15 (s, H₃-30). (H-3 hidden under H₂O at δ 4.92). – ¹³C NMR (CD₃OD): δ /ppm = 180.6 (C-28), 179.0 (C-23), 76.1 (C-20), 69.5 (C-29), 60.1 (C-17), 46.6 (C-19), 24.8 (C-30), 23.3 (C-2). – Negative ESI MS: *m/z* (%) = 561.5 (100) [M–H][–].

3 α , 20 ξ ,29-Trihydroxylupane-23,28-dioic acid (4c**)**

m.p. 234–235 °C (MeOH). – $[\alpha]_D^{24}$ +6.5 (MeOH, *c* = 1.0). – ¹H NMR (CD₃OD): δ /ppm = 3.72 (br s, H-3), 3.41 (s, H₂-29), 1.15 (s, H₃-24, H₃-30). – Negative ESI-MS: *m/z* (%) = 519.4 [M–H][–] (100).

3 α ,20 ξ ,29-Trihydroxylupane-23,28-dioic acid 23,28-dimethyl ester (4d**)**

m.p. 130–131 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ +174.6 (MeOH, *c* = 0.5). – IR: ν /cm^{–1} (CHCl₃) = 3 689 (m), 2 951 (m), 2 871 (w), 1 717 (s), 1 602 (m), 1 457 (m), 1 434 (m), 1 379 (w), 1 351 (w), 1 269 (m), 1 142 (w), 1 126 (w), 1 105 (w), 994 (w). – ¹H NMR (CDCl₃): δ /ppm = 3.75 (br s, H-3), 3.70 (s, OMe), 3.66 (s, OMe), 3.61 (d, *J* = 10.9 Hz, H-29^B), 3.40 (d, *J* = 10.9 Hz, H-29^A), 1.19, 1.17, 1.01, 0.89, 0.87 (each s, CH₃). – ¹³C NMR (CDCl₃): δ /ppm = 178.2 (C-28), 176.9 (C-23), 75.0 (C-20), 72.5 (C-3), 67.7 (C-29), 59.1 (C-17), 51.9, 51.3 (both OMe), 24.9 (C-2), 24.5 (C-30). – Positive ESI MS: *m/z* (%) = 571.5 [M+Na]⁺ (11), 531.4 [M+H–H₂O]⁺ (53), 513.4 (73), 471.4 (56), 453.4 (100).

Dioxolan derivative **5a**

m.p. 120–121 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ –18.9 (MeOH, *c* = 2.0). – IR: ν /cm^{–1} (CHCl₃) = 3 693 (w), 3 438 (w), 2 954

(s), 2870 (w), 1795 (s), 1734 (s), 1661 (s), 1500 (s), 1438 (m), 1374 (m), 1301 (w), 1109 (w), 1070 (m), 1023 (w), 994 (w). – ^1H NMR (CDCl_3): δ/ppm = 4.86 (br s, H-3), 4.30 (d, J = 8.8 Hz, H-29^B), 4.12 (d, J = 8.5 Hz, H-29^A), 2.77 (td, H-19), 2.08 (s, CO–CH₃), 1.52 (s, H₃-30). – ^{13}C NMR (CDCl_3): δ/ppm = 175.6, 174.6 (C-23, C-28), 154.5 (O–CO–O), 85.9 (C-20), 75.0 (C-3), 72.1 (C-29), 57.6 (C-17), 45.8 (C-19), 26.7 (C-30). – EIMS: m/z (%) = 878 [M]⁺ (7), 818 (16), 744 (24), 628 (69).

3 α ,20 ξ ,29-Trihydroxylupane-23,28-dioic acid 23,28-dime-thionin amide (5b)

m.p. 165–166 °C (EtOAc). – $[\alpha]_{\text{D}}^{24}$ –54.1 (MeOH, c = 1.0). – IR: ν/cm^{-1} (KBr) = 3420 (m, br), 2941 (s), 2870 (w), 1734 (s), 1636 (s), 1507 (s), 1448 (m), 1387 (w), 1203 (m, br), 1042 (w), 960 (w). – ^1H NMR (CD_3OD): δ/ppm = 3.74 (br s, H-3), 3.40 (s, H₂-29), 1.154, 1.145 (each s, H₃-24, H₃-30). – ^{13}C NMR (CD_3OD): δ/ppm = 175.9, 175.8 (C-23, C-28), 76.1 (C-20), 73.6 (C-3), 69.6 (C-29), 59.6 (C-17), 46.4 (C-19), 28.7 (C-15), 24.7 (C-30). – Positive ESI MS: m/z (%) = 783 [$\text{M} + \text{H}$]⁺ (100), 381 (14).

Dioxolan Derivative 6a

m.p. 110–112 °C ($\text{Me}_2\text{CO}/n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ +76.9 (MeOH, c = 2.0). – IR: ν/cm^{-1} (CHCl_3) = 3447 (m), 2951 (m), 2868 (w), 1795 (s), 1733 (s), 1662 (s), 1497 (s), 1374 (m), 1246 (m), 1109 (m), 1069 (m), 1024 (w). – ^1H NMR (CDCl_3): δ/ppm = 4.79 (br s, H-3), 4.26 (d, J = 8.6 Hz, H-29^B), 4.10 (d, J = 8.8 Hz, H-29^A), 2.72 (t, J = 8.8 Hz, H-19), 1.88 (s, CO–CH₃), 1.50 (s, H₃-30), 1.13, 0.94, 0.86, 0.69 (each s, 3H, CH₃). – ^{13}C NMR (CDCl_3): δ/ppm = 175.3, 174.5 (C-23, C-28), 154.6 (O–CO–O), 86.0 (C-20), 75.3 (C-3), 72.1 (C-29), 57.5 (C-17), 46.0 (C-19), 29.0, 27.9 (C-21, C-15), 26.8 (C-30). – Positive ESI MS: m/z (%) = 961.8 [$\text{M} + \text{Na}$]⁺ (70), 941.0 (55), 880.6 (100).

3 α ,20 ξ ,29-Trihydroxylupane-23,28-dioic acid 23,28-di(phenylalanin)amide (6b)

m.p. 168–169 °C (EtOAc/ $n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ +169.5 (MeOH, c = 1.0). – IR: ν/cm^{-1} (KBr) = 3434 (s), 2938 (s), 2870 (w), 1734 (s), 1636 (s), 1507 (s), 1456 (m), 1387 (m), 1207 (m, br), 1032 (m), 919 (w), 737 (w), 701 (m). – ^1H NMR (CD_3OD): δ/ppm = 3.65 (br s, H-3), 3.35 (s, H₂-29), 2.38 (t, H-19), 1.10, 1.05, 0.91, 0.81, 0.41 (each s, CH₃). – ^{13}C NMR (CD_3OD): δ/ppm = 175.5, 175.2 (C-23, C-28), 76.1 (C-20), 73.5 (C-3), 69.5 (C-29), 59.4 (C-17), 46.3 (C-19), 24.6 (C-30). – Positive ESI MS: m/z (%) = 854 [$\text{M} + \text{K}$]⁺ (20), 815 [$\text{M} + \text{H}$]⁺ (100).

3-O-Oxamoyl Derivative 7a

m.p. 128–130 °C ($\text{Me}_2\text{CO}/n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ –46.4 (MeOH, c = 0.5). – IR: ν/cm^{-1} (CHCl_3) = 3674 (w), 3437 (m), 3408 (w), 2954 (s), 2872 (m), 1795 (s), 1742 (s), 1706 (s), 1654 (s), 1506 (s), 1452 (s), 1378 (m), 1342 (m), 1279 (m), 1069 (m), 984 (m). – ^1H NMR (CDCl_3): δ/ppm = 4.87 (br s, H-3), 4.28 (d, J = 8.8 Hz, H-29^B), 4.11 (d, J = 8.3 Hz, H-29^A), 2.77 (tm, J = 8.9 Hz, H-19), 1.50 (s, H₃-30). – ^{13}C NMR (CDCl_3): δ/ppm = 175.1, 174.1 (C-23, C-28), 158.3, 156.2 (O–CO–CO) 154.5 (O–CO–O), 86.1 (C-20), 78.0 (C-3), 72.5 (C-29), 57.4 (C-17), 46.5 (C-19), 26.4 (C-30). – Positive ESI MS: m/z (%) = 897.5 [$\text{M} + \text{Na} + \text{H}$]⁺ (32), 874.6 [$\text{M} + \text{H}$]⁺ (100), 699.5 (61).

3 α ,20 ξ ,29-Trihydroxylupane-23,28-dioic acid 23,28-dialanin amide (7b)

m.p. 195–196 °C (EtOAc/ $n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ –45.9 (MeOH, c = 0.5). – ^1H NMR (CD_3OD): δ/ppm = 3.76 (br s, H-3), 3.40 (s, H₂-29), 2.58 (t, J = 11.7 Hz, H-19), 1.15 (s, H₃-30). – ^{13}C NMR (CD_3OD): δ/ppm = 176.9, 176.5 (C-23, C-28), 76.1 (C-20), 73.8 (C-3), 69.6 (C-29), 59.4 (C-17), 46.4 (C-19), 24.7 (C-30). – Positive ESI MS: m/z (%) = 663.6 [$\text{M} + \text{H}$]⁺ (100), 645.5 (33), 554.7 (64).

3 α -Acetoxy-20-oxo-29-norlupane-23,28-dioic acid (8a)

m.p. 155–157 °C (EtOAc). – $[\alpha]_{\text{D}}^{24}$ +61.8 (MeOH, c = 1.0). – IR: ν/cm^{-1} (CHCl_3) = 3674 (w), 3511 (w), 2950 (s), 2869 (m), 1733 (s), 1700 (s), 1602 (w), 1452 (m), 1375 (m), 1357 (m), 1248 (s), 1172 (m), 1130 (w), 1027 (m), 990 (w). – ^1H NMR (CDCl_3): δ/ppm = 4.95 (br s, H-3), 3.23 (td, J = 10.7 and 4.4 Hz, H-19), 2.19 (s, H₃-30). – ^{13}C NMR (CDCl_3): δ/ppm = 212.2 (C-20), 182.7 (C-28), 181.7 (C-23), 75.3 (C-3), 56.3 (C-17), 30.0 (C-30). – EIMS: m/z (%) = 530 [M]⁺ (35), 512 (16), 484 (20), 470 (46), 452 (84), 426 (100), 383 (54), 261 (53), 219 (88), 189 (69), 175 (74).

3 α -Hydroxy-20-oxo-29-norlupane-23,28-dioic acid (8b)

m.p. 231–232 °C (EtOAc). – $[\alpha]_{\text{D}}^{24}$ –38.6 (MeOH, c = 1.0). – ^1H NMR (CD_3OD): δ/ppm = 3.71 (br s, H-3), 3.27 (dt, H-19), 2.18 (s, H₃-30). – ^{13}C NMR (CD_3OD): δ/ppm = 215.5 (C-20), 180.5 (C-23), 179.6 (C-28), 73.8 (C-3), 57.3 (C-17), 50.6 (C-19), 29.8 (C-30). – EIMS: m/z (%) = 488 [M]⁺ (14), 470 (21), 452 (18), 442 (44), 426 (78), 424 (45), 383 (31), 261 (70), 219 (51), 175 (100), 147 (58).

3 α -Hydroxylupane-20-oxo-29-norlupane-23,28-dioic acid 23,28-dimethyl ester (8c)

m.p. 133–134 °C ($\text{Me}_2\text{CO}/n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ +97.2 (MeOH, c = 0.5). – ^1H NMR (CDCl_3): δ/ppm = 3.74 (br s, H-3), 3.70 (s, OMe), 3.67 (s, OMe), 3.25 (td, J = 11.2 and 4.1 Hz, H-19), 2.25 (m, 1H), 2.18 (s, H₃-30), 1.16 (s, H₃-24). – ^{13}C NMR (CDCl_3): δ/ppm = 212.3 (C-20), 178.1 (C-23), 176.5 (C-28), 72.5 (C-3), 56.4 (C-17), 51.4, 51.3 (both OMe), 49.4 (C-19), 30.0 (C-30). – EIMS: m/z (%) = 516 [M]⁺ (100), 498 (36), 484 (33), 466 (63), 456 (32), 438 (34), 275 (29), 233 (73), 189 (42), 147 (36), 57 (41).

3 α -Acetoxy-20-oxo-29-norlupane-23,28-dioic acid 23,28-di(alanin methyl ester)amide (9a)

m.p. 139–141 °C ($\text{Me}_2\text{CO}/n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ –55.0 (CHCl_3 , c = 2.0). – IR: ν/cm^{-1} (CHCl_3) = 3691 (w), 3446 (m), 2953 (s), 2868 (m), 1735 (s), 1706 (w), 1656 (m), 1602 (w), 1502 (m), 1450 (m), 1374 (w), 1250–1233 (m), 1172 (w), 985 (w). – ^1H NMR (CDCl_3): δ/ppm = 4.84 (br s, H-3), 3.41 (td, J = 11.1 and 4.0 Hz, H-19), 2.17 (s, H₃-30). – ^{13}C NMR (CDCl_3): δ/ppm = 212.7 (C-20), 175.7, 174.1 (C-23, C-28), 75.0 (C-3), 55.4 (C-17), 49.8 (C-19), 30.2 (C-30). – Positive ESI MS: m/z (%) = 723.6 [$\text{M} + \text{Na}$]⁺ (33), 641.6 [$\text{M} + \text{H} - \text{CH}_3\text{COOH}$]⁺ (100), 510.4 (44).

3 α -Hydroxy-20-oxo-29-norlupane 23,28-dioic acid 23,28-dialanin amide (9b)

m.p. 210–212 °C (EtOAc/ $n\text{-hexane}$). – $[\alpha]_{\text{D}}^{24}$ +21.8 (MeOH, c = 2.0). – IR: ν/cm^{-1} (KBr) = 3420 (m, br), 2944 (s), 2869 (w), 1734 (s), 1636 (s), 1521 (s), 1457 (m), 1379 (w), 1201 (m), 1066 (w). – ^1H NMR (CD_3OD): δ/ppm = 3.75 (br s, H-

3), 2.16 (s, H₃-30). – ¹³C NMR (CD₃OD): δ /ppm = 215.9 (C-20), 179.6, 178.8 (C-23, C-28), 73.7 (C-3), 56.7 (C-17), 49.2 (C-19), 29.8 (C-30). – Positive ESI MS: m/z (%) = 653 [M+Na]⁺ (12), 631 [M+H]⁺ (100).

3 α -Acetoxy-20-oxo-29-norlupane-23,28-dioic acid 23,28-di (serin methyl ester)amide (10)

m.p. 138–139 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ – 77.4 (MeOH, *c* = 2.0). – IR: ν /cm^{–1} (CHCl₃) = 3 691 (w), 3 442 (w), 2 955 (m), 2 870 (w), 1 735 (s), 1 706 (w), 1 659 (m), 1 602 (w), 1 501 (m), 1 438 (m), 1 375 (m), 1 356 (m), 1 240 (s). – ¹H NMR (CDCl₃): δ /ppm = 4.86 (br s, H-3), 3.37 (td, *J* = 11.0 and 3.9 Hz, H-19), 2.17 (s, H₃-30), 2.08 (s, CO–CH₃). – ¹³C NMR (CDCl₃): δ /ppm = 213.0 (C-20), 176.6, 175.3 (C-23, C-28), 74.8 (C-3), 55.6 (C-17), 49.7 (C-19), 30.1 (C-30). – EIMS: m/z (%) = 732 [M]⁺ (7), 714 (10), 672 (61), 654 (20), 526 (40), 201 (39), 120 (32), 57 (100).

3 α -Acetoxy-20-oxo-29-norlupane-23,28-dioic acid 23,28-di (glutaminic acid dimethyl ester) amide (11)

m.p. 124–125 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ – 42.4 (CHCl₃, *c* = 1.0). – IR: ν /cm^{–1} (CHCl₃) = 3 687 (w), 3 436 (w), 2 953 (m), 2 868 (w), 1 743 (s), 1 661 (m), 1 602 (w), 1 502 (m), 1 438 (m), 1 373 (w), 1 356 (m), 1 245 (m), 1 173 (m), 1 023 (w), 990 (w). – ¹H NMR (CDCl₃): δ /ppm = 4.84 (br s, H-3), 3.38 (td, *J* = 10.9 and 3.6 Hz, H-19), 2.16 (s, H₃-30), 2.03 (s, CO–CH₃). – ¹³C NMR (CDCl₃): δ /ppm = 212.7 (C-20), 176.4, 174.9 (C-23, C-28), 75.2 (C-3), 55.6 (C-17), 49.6 (C-19), 30.2 (C-30). – Positive ESI MS: m/z (%) = 867 [M+Na]⁺ (26), 845 [M+H]⁺ (100), 785 (52).

3 α -Acetoxy-20 ξ ,29-epoxylupane-23,28-dioic acid (12)

m.p. 180–182 °C (CHCl₃). – $[\alpha]_D^{24}$ – 29.2 (CHCl₃, *c* = 1.0). – IR: ν /cm^{–1} (CHCl₃) = 2 693 (w), 3 510 (m), 2 951 (s), 2 871 (m), 1 733 (s), 1 700 (s), 1 602 (w), 1 453 (m), 1 376 (m), 1 249 (s), 1 129 (w), 1 105 (w), 1 023 (w), 986 (w), 941 (w). – ¹H NMR (CDCl₃): δ /ppm = 4.95 (br s, H-3), 2.65 (br s, H₂-29), 1.24, 1.25 (each s, H₃-24, H₃-30). – ¹³C NMR (CDCl₃): δ /ppm = 182.4 (C-28), 181.6 (C-23), 77.2 (C-20), 75.4 (C-3), 60.2 (C-29), 56.6 (C-17), 45.3 (C-19), 26.9 (C-21), 18.2 (C-30). – Positive ESI MS: m/z (%) = 583 [M+K]⁺ (31), 567 [M+Na]⁺ (49), 562 [M+NH₄]⁺ (100), 545 [M+H]⁺ (18), 526 (12), 439 (18).

3 α -Hydroxy-29-oxolupane-23,28-dioic acid 23-methyl ester (C-20 epimers 13a/b)

Solid. – $[\alpha]_D^{24}$ – 13.4 (CHCl₃, *c* = 0.65). – ¹H NMR (CDCl₃): δ /ppm = 9.85 (d, *J* = 1.7 Hz, H-29)/9.64 (s, H-29), 3.76 (brs, H-3), 3.71 (s, OMe), 1.12 (d, *J* = 6.9 Hz, H₃-30)/1.02 (d, *J* = 6.9 Hz, H₃-30). – ¹³C NMR (CDCl₃): δ /ppm = 206.7/204.7 (C-29), 181.6/181.5 (C-28), 178.2 (C-23), 72.5 (C-3), 24.9 (C-2), 56.6/56.3 (C-17), 52.0 (OMe), 49.2/48.1 (C-19), 42.6/42.7 (C-14), 42.4/36.9 (C-20), 27.3/24.3 (C-21), 14.4/6.8 (C-30). – EIMS: m/z (%) = 516 [M]⁺ (11), 498 (6), 486 (14), 459 (14), 436 (16), 414 (27), 233 (100), 173 (88), 147 (64), 133 (64), 121 (84), 107 (81), 81 (82).

3 α ,20 ξ -Dihydroxy-29-oxolupane-23,28-dioic acid 23-methyl ester (13c)

m.p. 230–232 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ – 3.8 (CHCl₃, *c* = 1.0). – ¹H NMR (CDCl₃): δ /ppm = 9.53 (s, H-29), 3.75 (br s, H-3), 3.71 (s, OMe), 2.45 (t, *J* = 9.2 Hz, H-19), 1.27 (s,

H₃-30). 1.16 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 204.3 (C-29), 182.0 (C-28), 178.1 (C-23), 80.0 (C-20), 72.5 (C-3), 58.8 (C-17), 45.0 (C-19), 24.9 (C-2), 22.5 (C-30). – EIMS: m/z (%) = 532 [M]⁺ (17), 514 (6), 503 (18), 487 (100), 460 (16), 409 (16), 381 (19), 233 (63), 173 (55), 147 (75), 121 (68), 95 (64), 81 (68). – HRMS: 532.3382 [M]⁺ (C₃₁H₄₈O₇ requires 532.3400).

3 α -Hydroxy-29-oxolupane-23,28-dioic acid dimethyl ester (C-20 epimers 14a/b)

Solid. – $[\alpha]_D^{24}$ – 10.7 (CHCl₃, *c* = 0.85). – ¹H NMR (CDCl₃): δ /ppm = 9.85 (d, *J* = 1.7 Hz, H-29)/9.64 (s, H-29), 3.75 (br s, H-3), 3.704/3.665 (each s, OMe), 3.689/3.669 (each s, OMe), 2.98 (m, H-19), 1.00/1.11 (each d, *J* = 6.8 Hz, H₃-30). 1.17 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 206.8/204.7 (C-29), 178.12/178.08 (C-23), 176.47/176.42 (C-28), 72.4 (C-3), 56.75/56.53 (C-17), 51.9 (OMe), 51.32/51.27 (OMe), 42.46/36.87 (C-20), 49.96/49.89 (C-18), 49.2/48.2 (C-19), 24.9 (C-2), 14.4/6.7 (C-30). – EIMS: m/z (%) = 530 [M]⁺ (8), 500 (37), 473 (16), 440 (14), 264 (42), 251 (32), 233 (100), 189 (52), 175 (61), 161 (68), 119 (79), 105 (61), 81 (63).

3 α ,20 ξ -Dihydroxy-29-oxolupane-23,28-dioic acid dimethyl ester (14c)

m.p. 116–117 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ + 62.6 (CHCl₃, *c* = 0.3). – ¹H NMR (CDCl₃): δ /ppm = 9.53 (s, H-29), 3.75 (t, *J* = 2.8 Hz, H-3), 3.70, 3.66 (each s, OMe), 2.47 (t, *J* = 8.5 Hz, H-19), 1.27 (s, H₃-30). 1.17 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 204.4 (C-29), 178.1 (C-23), 176.7 (C-28), 80.0 (C-20), 72.5 (C-3), 51.9, 51.4 (both OMe), 24.9 (C-2), 22.4 (C-30). – EIMS: m/z (%) = 546 [M]⁺ (9), 528 (6), 517 (24), 501 (91), 473 (34), 423 (16), 329 (15), 233 (40), 175 (32), 161 (48), 147 (56), 119 (38), 83 (100).

Lup-2,20(29)-diene-23,28-dioic acid 23-methyl ester (15a)

m.p. 238–240 °C (MeOH). – $[\alpha]_D^{24}$ + 28.6 (CHCl₃, *c* = 1.0). – IR: ν /cm^{–1} (CHCl₃) = 2 950 (s), 2 871 (w), 1 717 (s), 1 697 (m), 1 646 (w), 1 602 (w), 1 457 (m), 1 435 (m), 1 377 (m), 1 146 (w), 1 103 (m), 1 084 (m), 891 (m). – ¹H NMR (CDCl₃): δ /ppm = 5.64 (ddd, *J* = 9.9, 6.1 and 1.6 Hz, H-2), 5.49 (dd, *J* = 9.9 and 2.2 Hz, H-3), 1.20 (s, H₃-24). – ¹³C NMR (CDCl₃): δ /ppm = 182.7 (C-28), 177.6 (C-23), 130.4 (C-3), 125.7 (C-2), 56.4 (C-17), 40.7 (C-1), 18.1 (C-24). – EIMS: m/z (%) = 482 [M]⁺ (34), 467 (8), 456 (11), 436 (16), 423 (34), 377 (20), 248 (40), 246 (39), 233 (39), 203 (27), 189 (86), 119 (100), 105 (54). – HRMS: 482.3400 [M]⁺ (C₃₁H₄₆O₄ requires 482.3396).

Lup-2,20(29)-diene-23,28-dioic acid (15b)

m.p. 290–292 °C (EtOAc). – $[\alpha]_D^{24}$ – 22.9 (MeOH, *c* = 1.0). – ¹H NMR (pyridine-d₅): δ /ppm = 5.92 (dd, *J* = 10.0 and 2.1 Hz, H-3), 5.75 (ddd, *J* = 10.4 and 4.5 Hz, H-2), 3.54 (td, *J* = 10.7 and 4.2 Hz, H-19), 2.76 (td, 1H, *J* = 11.8 and 3.0 Hz), 2.59 (dm, 1H, *J* = 12.9 Hz), 2.49 (dm, 1H, *J* = 8.3 Hz), 1.78 (s, H₃-30), 1.48 (s, H₃-24). – ¹³C NMR (pyridine-d₅): δ /ppm = 179.4, 178.9 (C-28, C-23), 132.3 (C-3), 125.3 (C-2), 56.6 (C-17), 48.5 (C-4), 41.3 (C-1), 18.2 (C-24). – EIMS: m/z (%) = 468 [M]⁺ (6), 440 (11), 422 (14), 248 (24), 235 (30), 203 (33), 189 (68), 173 (42), 105 (44), 97 (53), 81 (56) 69 (65), 57 (100).

[20 ξ]-29-Oxo-lup-2-ene-23,28-dioic acid 23-methyl ester (16a)

m.p. 194–196 °C (CHCl₃/*n*-hexane). – $[\alpha]_D^{24}$ –43.5 (CHCl₃, *c* = 1.0). – IR: ν/cm^{-1} (CHCl₃) = 2950 (s), 2872 (m), 1716 (s), 1700 (m), 1602 (w), 1456 (m), 1435 (m), 1378 (m), 1146 (m), 1102 (m), 1083 (m), 997 (w). – ¹H NMR (CDCl₃): δ/ppm = 9.84 (d, *J* = 1.9 Hz, H-29), 5.65 (ddd, *J* = 9.9, 4.3 and 1.4 Hz, H-2), 5.51 (dd, *J* = 10.2 and 2.2 Hz, H-3), 3.67 (s, OMe), 2.57 (qm, *J* = 7.1 and 2.3 Hz, H-20), 2.45 (m, H-19), 1.20 (s, H₃-24), 1.13 (d, *J* = 7.2 Hz, H₃-30). – ¹³C NMR (CDCl₃): δ/ppm = 206.8 (C-29), 182.1 (C-28), 177.5 (C-23), 130.5 (C-3), 125.5 (C-2), 56.6 (C-17), 52.2 (OMe), 48.8 (C-19), 48.2 (C-4), 42.4 (C-20), 40.7 (C-1), 27.4 (C-21), 18.1 (C-24), 14.4 (C-30). – EIMS: *m/z* (%) = 498 [M]⁺ (4), 485 (9), 469 (23), 442 (17), 233 (24), 219 (37), 189 (42), 173 (84), 161 (52), 147 (74), 133 (71), 119 (100), 105 (89), 93 (87), 67 (73), 55 (96).

C-20 Epimer 16b

m.p. 197–198 °C (CHCl₃/*n*-hexane). – $[\alpha]_D^{24}$ +39.1 (CHCl₃, *c* = 1.0). – IR: ν/cm^{-1} (CHCl₃) = 2950 (s), 2870 (m), 1717 (s), 1700 (m), 1602 (w), 1453 (m), 1434 (w), 1378 (m), 1146 (w), 1103 (w), 1083 (w), 997 (w). – ¹H NMR (CDCl₃): δ/ppm = 9.65 (s, H-29), 5.65 (ddd, *J* = 10.5, 4.8 and 1.4 Hz, H-2), 5.51 (dd, *J* = 10.2 and 2.2 Hz, H-3), 3.67 (s, OMe), 2.99 (tm, *J* = 10.7 Hz, H-19), 2.65 (qd, *J* = 6.8 and 2.9 Hz, H-20), 1.21 (s, H₃-24), 1.02 (d, *J* = 6.9 Hz, H₃-30). – ¹³C NMR (CDCl₃): δ/ppm = 204.7 (C-29), 182.1 (C-28), 177.6 (C-23), 130.5 (C-3), 125.5 (C-2), 56.4 (C-17), 52.2 (OMe), 48.3 (C-4), 36.9 (C-20), 40.7 (C-1), 24.3 (C-21), 18.1 (C-24), 6.8 (C-30). – EIMS: *m/z* (%) = 498 [M]⁺ (11), 439 (14), 396 (24), 233 (38), 187 (36), 173 (99), 148 (54), 133 (42), 119 (100), 105 (51), 95 (56).

20 ξ -Hydroxy-29-oxo-lup-2-ene-23,28-dioic acid 23-methyl ester (16c)

m.p. 218–219 °C (CHCl₃/*n*-hexane). – $[\alpha]_D^{24}$ –14.6 (CHCl₃, *c* = 0.25). – IR: ν/cm^{-1} (CHCl₃) = 3691 (w), 3510 (m), 2951 (s), 2872 (m), 1718 (s), 1602 (w), 1454 (m), 1435 (w), 1378 (w), 1146 (w), 1104 (w), 1083 (w), 831 (w). – ¹H NMR (CDCl₃): δ/ppm = 9.54 (s, H-29), 5.65 (dd, *J* = 9.9 and 4.7 Hz, H-2), 5.50 (dd, *J* = 10.2 and 2.2 Hz, H-3), 3.67 (s, OMe), 2.47 (t, *J* = 9.1 Hz, H-19), 1.19 (s, H₃-24), 1.28 (s, H₃-30). – ¹³C NMR (CDCl₃): δ/ppm = 204.3 (C-29), 182.3 (C-28), 177.5 (C-23), 130.5 (C-3), 125.6 (C-2), 80.0 (C-20), 58.9 (C-17), 52.2 (OMe), 48.2 (C-4), 43.9 (C-19), 40.8 (C-1), 29.9 (C-21), 22.5 (C-30), 18.1 (C-24). – EIMS: *m/z* (%) = 514 [M]⁺ (4), 485 (38), 469 (36), 395 (21), 231 (22), 173 (63), 147 (48), 119 (100), 105 (64), 95 (60), 81 (48).

23-Nor-lup-3,20(29)-dien-28-oic acid (17)

m.p. 208–210 °C (EtOAc). – $[\alpha]_D^{24}$ +37.7 (MeOH, *c* = 0.5). – ¹H NMR (pyridine-d₅): δ/ppm = 5.29 (br s, H-3), 3.54 (td, H-19), 1.80 (s, H₃-24), 1.65 (s, H₃-30), 1.09, 1.05, 0.73 (each s, Me). – ¹³C NMR (pyridine-d₅): δ/ppm = 178.9 (C-28), 135.1 (C-4), 120.6 (C-3), 56.6 (C-17), 47.7 (C-19), 21.6 (C-24). – EIMS: *m/z* (%) = 424 [M]⁺ (100), 409 (32), 369 (12), 259 (47), 248 (23), 201 (16), 189 (28), 175 (56). – HRMS: 424.3318 [M]⁺ (C₂₉H₄₄O₂ requires 424.3341).

3 α ,4 α -Dihydroxy-23-norlup-20(29)-en-28-oic acid (18)

m.p. 258–259 °C (EtOAc/*n*-hexane). – $[\alpha]_D^{24}$ +35.9 (MeOH,

c = 1.0). – IR: ν/cm^{-1} (CHCl₃) = 3679 (w), 3517 (w), 2948 (s), 2871 (m), 1793 (w), 1737 (w), 1697 (s), 1641 (w), 1602 (w), 1452 (m), 1376 (m), 1129 (m), 1104 (m), 1067 (w), 890 (m). – ¹H NMR (CDCl₃): δ/ppm = 3.56 (br s, H-3), 3.01 (m, H-19), 1.69 (s, H₃-30), 1.11 (s, H₃-24), 1.00, 0.94, 0.80 (each s, Me). – ¹³C NMR (CDCl₃): δ/ppm = 182.1 (C-28), 73.9 (C-3), 73.6 (C-4), 56.4 (C-17), 49.2 (C-19), 29.6 (C-12), 25.4 (C-2), 21.7 (C-24), 19.3 (C-30), 17.2 (C-6). – EIMS: *m/z* (%) = 458 [M]⁺ (14), 440 (28), 422 (15), 248 (51), 234 (33), 203 (32), 191 (56), 187 (50), 173 (65), 81 (54), 43 (100).

3 α ,4 α ,20 ξ ,29-Tetrahydroxy-23-norlupan-28-oic acid (19)

m.p. 234–236 °C (MeOH). – $[\alpha]_D^{24}$ +14.7 (MeOH, *c* = 0.5). – ¹H NMR (pyridine-d₅): δ/ppm = 4.04 (d, *J* = 10.5 Hz, H-29^B), 3.95 (d, *J* = 10.5 Hz, H-29^A), 3.85 (br s, H-3), 1.62, 1.26, 1.12, 0.99, 0.82 (each s, Me). – ¹³C NMR (pyridine-d₅): δ/ppm = 179.4 (C-28), 75.0 (C-20), 74.1 (C-3), 72.7 (C-4), 69.2 (C-29), 59.3 (C-17), 48.6 (C-19), 28.4 (C-12), 26.5 (C-2), 25.8 (C-30), 22.1 (C-24), 17.7 (C-6). – EIMS: *m/z* (%) = 592 [M]⁺ (1), 474 (13), 461 (59), 456 (38), 438 (28), 219 (29), 205 (33), 191 (96), 175 (64), 173 (100), 81 (45), 55 (51).

3,4-Dioxo-23-nor-3,4-seco-lup-20(29)-en-28-oic acid (20)

m.p. 94–96 °C (MeOH/*n*-hexane). – $[\alpha]_D^{24}$ +35.4 (MeOH, *c* = 1.0). – ¹H NMR (CDCl₃): δ/ppm = 9.76 (t, *J* = 1.6 Hz, H-3), 3.01 (td, *J* = 10.9 and 4.4 Hz, H-19), 2.57 (td, 1H, *J* = 12.1 and 3.3 Hz), 2.11 (s, H₃-24), 1.70 (s, H₃-30), 1.01, 0.99, 0.97 (each s, Me). – ¹³C NMR (CDCl₃): δ/ppm = 212.3 (C-4), 202.3 (C-3), 181.9 (C-28), 56.7 (C-17), 56.3 (C-9), 46.8 (C-19), 40.8 (C-5), 31.0 (C-24), 30.4, 29.6, 25.2 (C-15, C-12, C-2), 19.8, 19.3 (C-30, C-25). – EIMS: *m/z* (%) = 456 [M]⁺ (24), 438 (31), 412 (59), 395 (24), 369 (91), 259 (66), 189 (67), 175 (58), 107 (63), 93 (73), 81 (87), 55 (100). – HRMS: 456.3246 [M]⁺ (C₂₉H₄₄O₄ requires 456.3240).

3-Oxo-23-nor-lup-20(29)-en-28-oic acid (21)

m.p. 202–204 °C (EtOAc). – $[\alpha]_D^{24}$ +44.5 (MeOH, *c* = 0.5). – IR: ν/cm^{-1} (CHCl₃) = 2950 (s), 2875 (m), 1698 (s), 1646 (w), 1456 (m), 1377 (m), 1179 (m), 1134 (m), 890 (m). – ¹H NMR (CDCl₃): δ/ppm = 3.02 (td, *J* = 10.7 and 4.3 Hz, H-19), 2.44 (ddd, 1H, *J* = 15.1, 13.2 and 6.9 Hz), 1.70 (s, H₃-30), 0.978 (d, *J* = 6.6 Hz, H₃-24), 1.02, 1.00, 0.976 (each s, Me). – ¹³C NMR (CDCl₃): δ/ppm = 214.0 (C-3), 182.6 (C-28), 56.3 (C-17), 53.3 (C-4), 46.9 (C-19), 40.3 (C-2), 19.3 (C-30), 15.8, 14.5, 13.5, 11.6 (C-24, C-25, C-26, C-27). – EIMS: *m/z* (%) = 440 [M]⁺ (31), 425 (13), 394 (24), 248 (56), 235 (41), 219 (35), 203 (54), 189 (100), 177 (36), 107 (39), 95 (37), 81 (44).

2-Imino-1,3-dioxane derivative (22)

m.p. 187–188 °C (Me₂CO/*n*-hexane). – $[\alpha]_D^{24}$ +82.3 (CHCl₃, *c* = 1.0). – ¹H NMR (CD₃OD): δ/ppm = 4.71 (br s, H-29^B), 4.59 (br s, H-29^A), 4.41 (m, CH–N), 4.40 (br s, H-3), 3.01 (td, *J* = 10.7 and 4.1 Hz, H-19), 1.69 (s, H₃-30), 1.16 (s, H₃-24), 0.98, 0.97, 0.95 (each s, Me). – ¹³C NMR (CD₃OD): δ/ppm = 180.0 (C-28), 175.2 (C-23), 151.9 (C=N), 79.5 (C-3), 57.4 (C-17), 46.2 (C-4), 48.5 (C-19), 56.2, 29.6, 27.5, 27.4, 26.6 (cyclohexyl), 19.5 (C-30). – EIMS: *m/z* (%) = 593 [M]⁺

(56), 575 (19), 547 (100), 512 (29), 372 (22), 359 (86), 248 (35), 190 (69). – HRMS: 593.4086 [M]⁺ (C₃₇H₅₅NO₅ requires 593.4080).

References

- [1] H. Safayhi, E.-R. Sailer, *Planta Med.* **1997**, 63, 487
- [2] E. Pisha, H. Chai, I.-S. Lee, T. E. Chagwedera, N. R. Farnsworth, G. A. Cordell, C. W. Beecher, H. H. Fing, A. D. Kinghorn, D. M. Brown, M. C. Wani, M. E. Wall, T. J. Hieken, T. K. Gupta, J. M. Pezzuto, *Nature Med.* **1995**, 1, 1046
- [3] M. Sauvain, N. Kunesch, J. Poisson, J. L. Gantier, P. Gayral, J. P. Dedet, *Phytother. Res.* **1996**, 10, 1
- [4] G. Bringmann, W. Saeb, L. A. Assi, G. François, A. S. S. Narayanan, K. Peters, E.-M. Peters, *Planta Med.* **1997**, 63, 255
- [5] T. Fujioka, Y. Kashiwada, R. Kilkuskie, L. M. Cosentino, L. M. Ballas, J. B. Jiang, W. P. Janzen, I.-S. Chen, K.-H. Lee, *J. Nat. Prod.* **1994**, 57, 243
- [6] F. Soler, C. Poujade, M. Evers, J.-C. Carry, Y. Hénin, A. Bousseau, T. Huet, R. Pauwels, E. De Clercq, J.-F. Mayaux, J.-B. Le Pecq, N. Dereu, *J. Med. Chem.* **1996**, 39, 1069
- [7] M. Evers, C. Poujade, F. Soler, Y. Ribeill, C. James, Y. Lelièvre, J.-C. Gueguen, D. Reisdorf, I. Morize, R. Pauwels, E. De Clercq, Y. Hénin, A. Bousseau, J.-F. Mayaux, J.-B. Le Pecq, N. Dereu, *J. Med. Chem.* **1996**, 39, 1056
- [8] G. Adam, H. V. Lischewski, H. V. Phiet, A. Preiss, J. Schmidt, T. V. Sung, *Phytochemistry* **1982**, 21, 1385
- [9] T. V. Loc, H. Ripperger, C. Kamperdick, T. V. Sung, G. Adam, *Pharmazie* **1998**, 53, 677
- [10] V. VanRheenen, R. C. Kelly, D. Y. Cha, *Tetrahedron Lett.* **1976**, 23, 1973
- [11] Elemental analyses were within 0.3% of theory for all compounds.
- [12] The backbones of all compounds are nearly identical, so that the proton and carbon shifts are to a large extent very similar to those of the starting materials **1a** and **1b**, the full data of which are given in [9]. Therefore, only those carbon and proton shifts are given, that show significant differences because of structural variations. Full spectra are available from the corresponding author.

Address for correspondence :
Prof. Dr. G. Adam
Institute of Plant Biochemistry
Natural Products Chemistry
Weinberg 3
D-06120 Halle (Saale)
Fax : Internat. code (0)345/5582-102
e-Mail : gadam@ipb.uni-Halle.de