

A Straightforward Synthesis of Enantiomerically Pure *trans*-2,5-Bis(alkyloxymethyl)pyrrolidines by 1,3-Dipolar Cycloaddition Reactions of Azomethine Ylides ¹⁾

Claudia Wittland and Nikolaus Risch*

Paderborn, Universität-GH, Fachbereich für Chemie und Chemietechnik

Received February 15th, 2000

Keywords: Azomethine ylides, Domino reactions, Synthetic methods, Iminium salts, Pyrrolidines

Abstract. A new efficient method for the stereoselective preparation of *trans*-2,5-bis(alkyloxymethyl)pyrrolidines (**7**) using easily available starting materials is described. The main step of this synthesis is the stereoselective formation of the

pyrrolidine ring by the 1,3-dipolar cycloaddition reaction of an *in situ* generated azomethine ylide. Both enantiomers of the C₂-symmetric auxiliaries are prepared separately in a short reaction sequence.


2,5-Disubstituted pyrrolidine derivatives with a C₂ axis of symmetry are versatile chiral auxiliaries in asymmetric synthesis [1–3]. *Trans*-2,5-bis(alkyloxymethyl)pyrrolidines **7** especially are very powerful auxiliaries and the development of new efficient syntheses for these substances is of current interest. The corresponding diacid is a natural product which has been isolated from the red alga *schizymenia dubyi* [4].

The first synthesis of enantiomerically pure *trans*-2,5-bis(alkyloxymethyl)pyrrolidines **7** by optical resolution was published by Katsuki *et al.* [5] This method, starting from 2,5-dibromoadipate and dibenzylamine, was optimized by Ghazes *et al.* [6]. Later on, Yamamoto *et al.* [7] and Durst *et al.* [8] improved this synthesis by using chiral auxiliaries. Additional preparations have been reported in the last few years which started from optically active *D*-mannitol [9, 10], (*S*)-*O*-benzylglycidol [11], $\Delta^{4,5}$ oxazolidin-2-one [12, 13] or L-serine [14]. Most of these methods are based on long reaction sequences or require starting materials which are expensive or difficult to obtain.

During our investigations on iminium salts and 1,3-dipolar cycloaddition reactions of azomethine ylides [15] we developed an efficient one-pot reaction for the stereoselective preparation of highly substituted pyrrolidines [16, 17]. The *in situ* generation of azomethine ylides from secondary benzylic amines and aldehydes *via* iminium salts, followed by the cycloaddition with different dipolarophiles gives rise to a wide variety of 2,5-dicarboxylated and 2,5-diphenyl-substituted pyrrolidines. Using optically pure 1-phenylethylamine as a chiral template, enantiomerically pure 2,3,4,5-tetrasubstituted proline derivatives have already been synthesized by this method [17].

We now describe a new and short reaction sequence for the synthesis of enantiomerically pure *trans*-2,5-


bis(alkyloxymethyl)pyrrolidines **7** which includes the 1,3-dipolar cycloaddition reaction of an azomethine ylide to phenylvinylsulfone [18, 19] as an ethene equivalent. This method allows the stereoselective formation of the pyrrolidine ring by a simple one-pot reaction using easily available and inexpensive starting materials.



Scheme 1 Mixture of cycloadducts **3** with d.r. = 38 : 35 : 17 : 10

¹⁾ Presented in part at the Fourth Conference on Iminium Salts, Stimpfach-Rechenberg (Germany), September 14–16, 1999

The one-pot reaction of ethyl *N*-((*R*)-1-phenylethyl)glycinecarboxylate ((*R*)-**1**) and ethyl glyoxalate with phenylvinylsulfone as the dipolarophile in refluxing toluene, with azeotropic removal of water, yielded a diastereomeric mixture of the pyrrolidines **3** with d.r. = 38:35:17:10 (Scheme 1). These cycloadducts are formed by an *exo*- and an *endo*-attack of the dipolarophile on the *in situ* generated *anti*-azomethine ylide **2**. Using an excess of amine (*R*)-**1** and glyoxalate to reach a total conversion of the sulfone, we also obtained small amounts of oxazolidine as a by-product, which results from the cycloaddition of ethyl glyoxalate to the ylide **2**. The cycloaddition products were purified by column chromatography to yield 67% of the mixture of pyrrolidines **3**. These diastereomers were not separated, because it is more efficient to carry out the following reaction steps with the diastereomeric mixture of the cycloadducts **3**.



Scheme 2 a: LiAlH₄, 91%; b: NaH, MeI, 95%; c: NaH, Bzl-Br, 84%; d: 1. Na-Hg (10%), Na₂HPO₄, THF/MeOH; 2. chromatography; e: R = Me: H₂, 10% Pd/C, quant.; R = Bzl: H₂, 20% Pd(OH)₂/C, 90%.

The reduction of the pyrrolidines **3** with LiAlH₄ in THF led to the diols **4** in 91% yield. These products were transformed into the ethers **5** without prior purification. At this step of the synthesis different protecting groups of the hydroxy function can be introduced into the molecule. We prepared the methyl ethers **5a** by the reaction of the diols **4** with sodium hydride and methyl iodide in THF. The use of benzyl bromide gave rise to the dibenzylic ethers **5b**. In the latter case column chromatography was necessary to remove the excess of benzyl bromide, but the desulfonation that followed was also carried out successfully if the benzyl bromide was removed *in vacuo*, as far as possible.

The reductive desulfonation of **5a** with 10% Na-Hg in THF/MeOH, using Na₂HPO₄ as buffer, yielded the 2,5-disubstituted *N*-protected pyrrolidines **6a**. The two diastereomers were separated by column chromatography to give 48% of the major diastereomer (2*R*,5*R*)-**6a** and 14% of the minor diastereomer (2*S*,5*S*)-**6a**. The purity of the isolated fractions was determined by GC using a chiral column (HYDRODEX β-PM). The absolute configuration was proved by the determination of the optical rotation and comparison with the values reported in literature [7]. Using the same procedure the benzylic ethers **5b** were desulfonated to yield 39% of (2*R*,5*R*)-**6b** and 10% of (2*S*,5*S*)-**6b**.

The debenzylation of the *N*-protected auxiliaries **6** by heterogeneous catalysed hydrogenolysis offered the enantiomerically pure pyrrolidines **7**. **6a** was debenzylated by a standard method using 10% Pd/C as catalyst. Selective *N*-debzylolation of the benzylic ethers **6b** was achieved with Pearlman's catalyst (20% Pd(OH)₂/C) [9, 20].

In conclusion, we have developed an efficient reaction sequence for the synthesis of *trans*-2,5-bis(alkyloxymethyl)pyrrolidines **7**, which is also suitable for the preparation of larger amounts of these important auxiliaries. The 1,3-dipolar cycloaddition reaction of the *in situ* generated azomethine ylide **2** allows the stereoselective formation of the pyrrolidine ring by a simple one-pot reaction. As the separation of the diastereomeric cycloadducts was carried out just before the last step of the synthesis, both enantiomers of the auxiliaries **7** were obtained in a single reaction sequence.

By starting the synthesis with (*S*)-**1** as the amine we obtained the auxiliaries (2*S*,5*S*)-**7** as major products and (2*R*,5*R*)-**7** as minor products.

We would like to thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for supporting this research.

Experimental

Ethyl *N*-(1-phenylethyl)glycine carboxylate (**1**) [21] was syn-

thesized according to literature procedures. Anhydrous THF was freshly distilled from sodium under argon. Petroleum ether refers to the fraction with *b.p.* 50–70 °C. Column chromatography on silica gel was performed with Merck Kieselgel 60 (0.040–0.063 mm). ^1H and ^{13}C NMR spectra were recorded on a Bruker ARX 200 or a Bruker AMX 300 spectrometer, using TMS as internal standard. IR spectra were recorded on a Nicolet 510 P FT-IR spectrometer. GC/MS data were obtained from a Finnigan MAT Magnum System 240 and MS data from a VG Fisons Autospec. Specific optical rotations were measured with a Perkin Elmer 241 polarimeter using a 1.00 dm cell. Elemental analyses were performed on a Perkin Elmer 240 Elemental Analyzer. Microanalyses were in satisfactory agreement with the calculated values: $\text{C} \pm 0.31$, $\text{H} \pm 0.23$, $\text{N} \pm 0.22$.

2,5-Diethyl 1-(1-Phenylethyl)-3-phenylsulfonylpyrrolidine-2,5-dicarboxylate (**3**)

A solution of amine **1** (1.55 g, 7.5 mmol), ethyl glyoxalate (2.0 mL, 10.0 mmol; 50% in toluene) and phenyl vinyl sulfone (0.84 g, 5.0 mmol) in toluene (20 mL) was refluxed for 2 d under Dean-Stark conditions. After removal of the solvent under reduced pressure the crude product was purified by column chromatography (silica gel; petroleum ether/EtOAc, 80:20) to yield 1.54 g (67%) of the diastereomers **3** as a yellowish oil. The diastereomeric mixture is used for the next reaction step without further purification.

Diastereomeric Mixture of **3**

IR (film): $\nu/\text{cm}^{-1} = 2979, 1733, 1447, 1150$. – ^1H NMR (300 MHz, CDCl_3): $\delta/\text{ppm} = 1.04\text{--}1.41$ (m, $\text{CO}_2\text{CH}_2\text{CH}_3$, NCHCH_3), $2.43\text{--}3.03$ (m, 4-H), $3.67\text{--}4.33$ (m, 2-H, 3-H, 5-H, $\text{CO}_2\text{CH}_2\text{CH}_3$, NCHCH_3), $7.19\text{--}7.28$ (m, H_{arom}), $7.45\text{--}7.66$ (m, H_{arom}), $7.79\text{--}7.91$ (m, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): $\delta/\text{ppm} = 13.31\text{--}13.97$ ($\text{CO}_2\text{CH}_2\text{CH}_3$), $21.78\text{--}22.95$ (NCHCH_3), $29.10\text{--}30.53$ (C-4), $57.47\text{--}66.71$ (C-2, C-3, C-5, $\text{CO}_2\text{CH}_2\text{CH}_3$, NCHCH_3), $126.60\text{--}144.02$ (C_{arom}), $169.66\text{--}175.17$ (C=O). MS (70 eV): m/z (%) = 386 (16, $\text{M}^+ - \text{CO}_2\text{C}_2\text{H}_5$), 318 (6, $\text{M}^+ - \text{SO}_2\text{Ph}$), 282 (18), 216 (11), 140 (19), 105 (100).

$\text{C}_{24}\text{H}_{28}\text{NO}_6\text{S}$ Calcd.: C 62.86 H 6.16 N 3.06
(458.16) Found: C 63.11 H 5.97 N 3.27.

2,5-Bis(hydroxymethyl)-1-(1-phenylethyl)-3-phenylsulfonylpyrrolidine (**4**)

Under argon a solution of the diastereomeric mixture of **3** (1.54 g, 3.27 mmol) in anhydrous THF (40 mL) was cooled to 0 °C, LiAlH_4 (0.23 g, 6 mmol) was added, and the mixture was stirred for 3 h at r.t. H_2O was added carefully at 0 °C to destroy the excess of LiAlH_4 . The solid was separated by filtration and washed with EtOAc several times. The filtrate was dried (MgSO_4), and concentrated under reduced pressure to yield 1.26 g (quant.) of the diastereomers **4** as a yellow viscous oil.

Major Diastereomers **4**

^1H NMR (300 MHz, CDCl_3): $\delta/\text{ppm} = 1.42$ (d, 3H, NCHCH_3), 1.49 (d, 3H, NCHCH_3), $2.24\text{--}2.36$ (m, 4H, 4-H), 3.10 (br. s, OH), $3.28\text{--}3.86$ (m, 14H, 2-H, 3-H, 5-H, CH_2OH),

4.14 (q, 1H, NCHCH_3), 4.22 (q, 1H, NCHCH_3), $7.18\text{--}7.37$ (m, 10H, H_{arom}), $7.48\text{--}7.67$ (m, 6H, H_{arom}), $7.82\text{--}7.92$ (m, 4H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): $\delta/\text{ppm} = 17.42, 18.81$ (NCHCH_3), $27.56, 28.75$ (C-4), $53.96, 54.38$ (NCHCH_3), 59.30 (CH), 60.53 (CH_2), 61.21 (CH), 61.50 (CH_2), 62.05 (CH_2), 62.45 (CH), 62.51 (CH_2), $63.79, 64.57$ (CH), $126.45, 126.88, 127.27, 127.33, 128.35, 128.46, 128.57, 129.11, 129.20, 133.66, 133.73, 137.77, 137.89, 142.20, 145.23$ (C_{arom}).

$\text{C}_{20}\text{H}_{25}\text{NO}_4\text{S}$ Calcd.: C 63.97 H 6.71 N 3.73
(375.15) Found: C 64.25 H 6.34 N 3.54.

Mixture of all Diastereomers **4**

IR (Film): $\nu/\text{cm}^{-1} = 3373, 2937, 1446, 1032$. – MS (70 eV): m/z (%) = 344 (11, $\text{M}^+ - \text{CH}_2\text{OH}$), 240 (6), 220 (6), 202 (4), 105 (100).

2,5-Bis(methoxymethyl)-1-(1-phenylethyl)-3-phenylsulfonylpyrrolidine (**5a**)

A solution of the diastereomeric mixture of **4** (1.25 g, 3.3 mmol) in anhydrous THF (20 mL) and MeI (2.55 g, 17.9 mmol) was added slowly to a suspension of NaH (0.52 g, 17.9 mmol; 80%, washed with anhydrous *n*-Hexane) in anhydrous THF (10 mL) at 0 °C. The mixture was stirred for 2 h at r.t. and cooled to 0 °C before H_2O was added carefully. The solid was separated by filtration and the filtrate extracted with Et_2O (3×30 mL). The organic layer was washed with brine (2×30 mL) and dried (MgSO_4). The solvent was removed under reduced pressure to yield 1.25 g (95%) of the diastereomers **5a** as a yellowish oil.

Major Diastereomers **5a**

^1H NMR (300 MHz, CDCl_3): $\delta/\text{ppm} = 1.33$ (d, 3H, NCHCH_3), 1.39 (d, 3H, NCHCH_3), $1.98\text{--}2.42$ (m, 4H, 4-H), $3.01\text{--}3.72$ (m, 14H, 2-H, 3-H, 5-H, CH_2O), 3.06 (s, 6H, OCH_3), 3.14 (s, 3H, OCH_3), 3.15 (s, 3H, OCH_3), $4.01\text{--}4.15$ (m, 2H, NCHCH_3), $7.14\text{--}7.32$ (m, 10H, H_{arom}), $7.48\text{--}7.60$ (m, 6H, H_{arom}), $7.81\text{--}7.89$ (m, 4H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): $\delta/\text{ppm} = 18.20, 20.44$ (NCHCH_3), $30.00, 30.93$ (C-4), $55.29, 58.73, 59.04, 59.17, 59.25, 59.37, 59.96, 61.88, 64.95, 65.98$ (CH, CH_3), $72.91, 73.11, 75.05, 76.44$ (CH_2O), $126.75, 127.14, 127.25, 127.94, 128.39, 128.43, 129.19, 129.33, 129.43, 133.81, 138.36, 143.05, 145.84$ (C_{arom}).

$\text{C}_{22}\text{H}_{29}\text{NO}_4\text{S}$ Calcd.: C 65.48 H 7.25 N 3.47
(403.18) Found: C 65.24 H 7.45 N 3.69.

Mixture of all Diastereomers **5a**

IR (film): $\nu/\text{cm}^{-1} = 2859, 1450, 1182$. – MS (70 eV): m/z (%) = 358 (29, $\text{M}^+ - \text{CH}_2\text{OCH}_3$), 254 (18), 216 (5), 105 (100).

2,5-Bis(benzyloxymethyl)-1-(1-phenylethyl)-3-phenylsulfonylpyrrolidine (**5b**)

5b was prepared in accordance to the preparation of **5a** starting from the diastereomeric mixture of **4** (0.43 g, 1.2 mmol), benzyl bromide (0.59 g, 3.5 mmol) and NaH (0.19 g, 6.5 mmol; 80%, washed with anhydrous *n*-Hexane) to yield 0.51 g (84%) of the diastereomers **5b**. For the preparation of **6b** it is sufficient to remove the excess of benzyl bromide *in vacuo* as far as possible. Otherwise the crude product can be purified by

column chromatography (silica gel; petroleum ether/EtOAc, 70:30).

Mixture of all Diastereomers **5b**

IR (film): ν/cm^{-1} = 2857, 1452, 1186, 1029. – ^1H NMR (300 MHz, CDCl_3): δ/ppm = 1.22–1.36 (m, NCHCH_3), 2.15–2.60 (m, 4-H), 3.02–3.79 (m), 3.98–4.15 (m), 4.19–4.61 (m), 7.16–7.85 (m, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 17.75, 20.72 (NCHCH_3), 29.69, 30.51 (C-4), 54.67, 55.15 (NCHCH_3), 58.44, 59.08, 59.68, 61.60, 64.72, 65.98 (C-2, C-3, C-5), 69.85, 70.02, 72.53, 72.84, 73.00, 73.88 (CH_2O , OCH_2Ph), 126.27–133.48, 137.62–133.24 (C_{arom}). – MS (70 eV): m/z (%) = 434 (52, M^+ – $\text{PhCH}_2\text{OCH}_2$), 330 (31), 188 (22), 158 (22), 105 (83), 91 (100).

$\text{C}_{34}\text{H}_{37}\text{NO}_4\text{S}$ Calcd.: C 73.48 H 6.72 N 2.52
(555.24) Found: C 73.36 H 6.95 N 2.34.

2,5-Bis(methoxymethyl)-1-(1-phenylethyl)-pyrrolidine (**6a**) [7]

Under argon Na_2HPO_4 (0.72 g, 6.1 mmol) was added to a solution of the sulfone **5a** (1.1 g, 2.75 mmol) in anhydrous THF (15 mL) and anhydrous MeOH (10 mL). The suspension was cooled to 0 °C and 10% Na–Hg (3.25 g, 14 mmol Na) was added. After stirring at r.t. over night the liquid was decanted and the solid washed with Et_2O several times. The combined organic layers were washed with H_2O (2×30 mL) and dried (MgSO_4). The solvent was removed under reduced pressure and the crude product purified by column chromatography (silica gel; petroleum ether/EtOAc, 90:10) to yield 0.35 g (48%) of (2*R*,5*R*)-**6a**, 0.1 g (14%) of (2*S*,2*S*)-**6a** and 0.12 g (18%) of a mixture of both diastereomers **6a**.

(1*R*,2*R*,5*R*)-2,5-Bis(methoxymethyl)-1-(1-phenylethyl)-pyrrolidine ((**2R,5R**)-**6a**)

$[\alpha]_{\text{D}}^{20}$ = + 68.1° (c = 1.4, CH_2Cl_2). – IR (film): ν/cm^{-1} = 2924, 2875, 1451, 1120. – ^1H NMR (200 MHz, CDCl_3): δ/ppm = 1.36 (d, J = 6.7 Hz, 3H, NCHCH_3), 1.66–2.08 (m, 4H, 3-H, 4-H), 3.09–3.37 (m, 6H, 2-H, 5-H, $2\text{CH}_2\text{O}$), 3.23 (s, 6H, 2OCH_3), 4.02 (q, J = 6.7 Hz, 1H, NCHCH_3), 7.13–7.37 (m, 5H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 24.75 (NCHCH_3), 27.15 (C-3, C-4), 57.66 (NCHCH_3), 58.69, 58.78 (C-2, C-5, 2OCH_3), 75.19 ($2\text{CH}_2\text{O}$), 126.20, 126.69, 127.94, 146.83 (C_{arom}). – GC/MS (80 eV, 220 °C): m/z (%) = 263 (40, M^+), 231 (12), 218 (100), 114 (83), 105 (74).

$\text{C}_{16}\text{H}_{25}\text{NO}_2$ Calcd.: C 72.95 H 9.57 N 5.32
(263.19) Found: C 72.65 H 9.77 N 5.49.

(1*R*,2*S*,5*S*)-2,5-Bis(methoxymethyl)-1-(1-phenylethyl)-pyrrolidine ((**2S,5S**)-**6a**)

$[\alpha]_{\text{D}}^{20}$ = – 42.7° (c = 1.2, CH_2Cl_2). – IR (film): ν/cm^{-1} = 2929, 1436, 1031. – ^1H NMR (200 MHz, CDCl_3): δ/ppm = 1.49 (d, J = 6.5 Hz, 3H, NCHCH_3), 1.59–2.98 (m, 4H, 3-H, 4-H), 2.87–3.05 (m, 4H), 3.16 (s, 6H, 2OCH_3), 3.17–3.36 (m, 2H), 3.95 (q, J = 6.5 Hz, 1H, NCHCH_3), 7.23–7.38 (m, 5H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 21.01 (NCHCH_3), 26.86 (C-3, C-4), 57.49 (NCHCH_3), 58.60, 59.22 (C-2, C-5, 2OCH_3), 73.75 ($2\text{CH}_2\text{O}$), 126.86, 127.89, 145.09 (C_{arom}). – GC/MS (80 eV, 220 °C): m/z (%) = 263 (42, M^+), 231 (12), 217 (100), 114 (80), 105 (72).

$\text{C}_{16}\text{H}_{25}\text{NO}_2$ Calcd.: C 72.95 H 9.57 N 5.32
(263.19) Found: C 73.11 H 9.42 N 5.13.

2,5-Bis(benzyloxymethyl)-1-(1-phenylethyl)-pyrrolidine (**6b**)

6b was prepared in accordance to the preparation of **6a** starting from **5b** (0.51 g, 0.9 mmol) to yield 0.15 g (39%) of the major diastereomer (2*R*,5*R*)-**6b** and 0.04 g (10%) of the minor diastereomer (2*S*,2*S*)-**6b** as colourless viscous oils.

(1*R*,2*R*,5*R*)-2,5-Bis(benzyloxymethyl)-1-(1-phenylethyl)-pyrrolidine ((**2R,5R**)-**6b**)

$[\alpha]_{\text{D}}^{20}$ = + 58.7° (c = 1.1, CHCl_3). – IR (film): ν/cm^{-1} = 2855, 1454, 1099, 1029. – ^1H NMR (300 MHz, CDCl_3): δ/ppm = 1.29 (d, J = 6.6 Hz, 3H, NCHCH_3), 1.81–1.88 (m, 2H, 3-H, 4-H), 2.05–2.10 (m, 2H, 3-H, 4-H), 3.24–3.26 (m, 4H, $2\text{CH}_2\text{O}$), 3.35–3.43 (m, 2H, 2-H, 5-H), 3.98 (q, J = 6.6 Hz, 1H, NCHCH_3), 4.43 (s, 4H, OCH_2Ph), 7.25–7.35 (m, 15H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 25.02 (NCHCH_3), 27.55 (C-3, C-4), 57.80 (NCHCH_3), 59.07 (C-2, C-5), 72.82, 72.97 ($2\text{CH}_2\text{O}$, $2\text{OCH}_2\text{Ph}$), 126.20, 126.73, 127.28, 127.35, 128.00, 128.17, 138.56, 146.91 (C_{arom}).

$\text{C}_{28}\text{H}_{33}\text{NO}_2$ Calcd.: C 80.91 H 8.00 N 3.37
(416.25) Found: C 81.03 H 8.20 N 3.57.

(1*R*,2*S*,5*S*)-2,5-Bis(benzyloxymethyl)-1-(1-phenylethyl)-pyrrolidine ((**2S,5S**)-**6b**)

$[\alpha]_{\text{D}}^{20}$ = – 34.7° (c = 1.0, CHCl_3). – IR (film): ν/cm^{-1} = 2855, 1454, 1099, 1029. – ^1H NMR (300 MHz, CDCl_3): δ/ppm = 1.44 (d, J = 6.5 Hz, 3H, NCHCH_3), 1.77–1.83 (m, 2H, 3-H, 4-H), 1.91–1.98 (m, 2H, 3-H, 4-H), 3.01–3.14 (m, 4H, $2\text{CH}_2\text{O}$), 3.23–3.28 (m, 2H, 2-H, 5-H), 3.95 (q, J = 6.5 Hz, 1H, NCHCH_3), 4.32 (s, 4H, OCH_2Ph), 7.20–7.34 (m, 15H, H_{arom}). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 20.88 (NCHCH_3), 27.05 (C-3, C-4), 57.29 (NCHCH_3), 59.27 (C-2, C-5), 71.53, 72.87 ($2\text{CH}_2\text{O}$, $2\text{OCH}_2\text{Ph}$), 126.85, 127.21, 127.31, 127.86, 127.91, 128.09, 128.18, 138.42, 144.99 (C_{arom}).

$\text{C}_{28}\text{H}_{33}\text{NO}_2$ Calcd.: C 80.91 H 8.00 N 3.37
(416.25) Found: C 80.60 H 8.23 N 3.58.

(2*R*,5*R*)-2,5-Bis(methoxymethyl)pyrrolidine ((**2R,5R**)-**7a**) [7]

A solution of (2*R*,5*R*)-**6a** (0.3 g, 1.14 mmol) in anhydrous EtOH (20 mL) was stirred at r.t. in the presence of 10% Pd/C (0.06 g) and hydrogen was bubbled through the mixture until the debenzylation was completed (TLC control). After removal of the catalyst by filtration through Celite, the filtrate was evaporated to yield 0.18 g (quant.) of (2*R*,5*R*)-**7a** as a colourless oil. – $[\alpha]_{\text{D}}^{20}$ = – 5.2° (c = 1.3, Ethanol) [Lit.[7] $[\alpha]_{\text{D}}^{20}$ = – 7.6° (c = 3.0, Ethanol)]. – IR (film): ν/cm^{-1} = 3382, 2928, 2884, 2691, 1456, 1120. – ^1H NMR (200 MHz, CDCl_3): δ/ppm = 1.41–1.54 (m, 2H, 3-H, 4-H), 1.86–1.97 (m, 2H, 3-H, 4-H), 2.92 (br. s, 1H, NH), 3.25–3.47 (m, 6H, 2-H, 5-H, $2\text{CH}_2\text{O}$), 3.35 (s, 6H, 2OCH_3). – ^{13}C NMR (75 MHz, CDCl_3): δ/ppm = 27.56 (C-3, C-4), 56.73, 58.74 (C-2, C-5, 2OCH_3), 75.55 ($2\text{CH}_2\text{O}$). – GC/MS (70 eV, 220 °C): m/z (%) = 159 (57, M^+), 113 (100), 82 (48).

$\text{C}_8\text{H}_{17}\text{NO}_2$ C 60.33 H 10.77 N 8.80
(159.13) C 60.62 H 10.61 N 8.95.

(2*S*,5*S*)-**7a** was obtained from (2*S*,5*S*)-**6a**

$[\alpha]_{\text{D}}^{20}$ = + 5.0° (c = 1.1, Ethanol) [Lit.[7] $[\alpha]_{\text{D}}^{20}$ = + 7.8° (c = 3.0, Ethanol)]

$\text{C}_8\text{H}_{17}\text{NO}_2$ C 60.33 H 10.77 N 8.80
(159.13) C 60.59 H 10.92 N 8.97.

(2R,5R)-Bis(benzyloxymethyl)pyrrolidine (**7b**) [9]

(*2R,5R*)-**6b** (0.09 g, 0.22 mmol) was debenzylated according to the preparation of (*2R,5R*)-**7a**, using 20% Pd(OH)₂/C (0.02 g) as catalyst, to yield 0.06 g (90%) of (*2R,5R*)-**7b** as a colourless oil. – $[\alpha]_D^{25} = -2.8^\circ$ ($c = 1.3$, CHCl₃) [Lit.[9] $[\alpha]_D^{25} = -3.27^\circ$ ($c = 1$, MeOH)]. – IR (film): $\nu/\text{cm}^{-1} = 3385, 2930, 1454, 1122$. – ¹H NMR (300 MHz, CDCl₃): $\delta/\text{ppm} = 1.73\text{--}1.82$ (m, 2H, 3-H, 4-H), 2.03–2.11 (m, 2H, 3-H, 4-H), 3.60–3.74 (m, 4H, 2CH₂O), 3.77–3.89 (m, 2H, 2-H, 5-H), AB-signal ($\delta_A = 4.52$, $\delta_B = 4.61$, $J = 12.0$ Hz, 2OCH₂Ph), 7.25–7.30 (m, 10H, H_{arom}). – ¹³C NMR (75 MHz, CDCl₃): $\delta/\text{ppm} = 27.07$ (C-3, C-4), 58.47 (C-2, C-5), 69.84, 73.07 (2CH₂O, 2OCH₂Ph), 127.52, 127.72, 128.19, 137.65 (C_{arom}). – GC/MS (70 eV, 220 °C): m/z (%) = 311 (2, M⁺), 190 (55), 91 (100).

C ₂₀ H ₂₅ NO ₂	Calcd.: C 77.12	H 8.10	N 4.50
(311.19)	Found: C 77.38	H 8.27	N 4.71.

(*2S,5S*)-**7b** was obtained from (*2S,5S*)-**6b**; $[\alpha]_D^{20} = +2.9^\circ$ ($c = 1.1$, CHCl₃).

C ₂₀ H ₂₅ NO ₂	Calcd.: C 77.12	H 8.10	N 4.50
(311.19)	Found: C 77.35	H 7.95	N 4.69.

References

- [1] M. Pichon, B. Figadère, *Tetrahedron: Asymm.* **1996**, 7, 927
- [2] J. K. Whitesell, *Chem. Rev.* **1989**, 89, 1581
- [3] J. Seyden-Penne, *Chiral Auxiliaries and Ligands in Asymmetric Synthesis* Wiley, New York 1995
- [4] G. Impellizzeri, S. Mangiafico, G. Oriente, M. Piateli, S. Scinto, *Phytochemistry* **1975**, 14, 1549
- [5] Y. Kawanami, Y. Ito, T. Kitagawa, Y. Taniguchi, T. Katsuki, M. Yamaguchi, *Tetrahedron Lett.* **1984**, 25, 857
- [6] L. Y. Chen, L. Ghosez, *Tetrahedron Lett.* **1990**, 31, 4467
- [7] Y. Yamamoto, J. Hoshino, Y. Fujimoto, J. Ohmoto, S. Sawada, *Synthesis* **1993**, 298
- [8] K. Koh, R. N. Ben, T. Durst, *Tetrahedron Lett.* **1994**, 35, 375
- [9] M. Marzi, P. Minetti, D. Misiti, *Tetrahedron* **1992**, 48, 10127
- [10] M. Marzi, D. Misiti, *Tetrahedron Lett.* **1989**, 30, 6075
- [11] S. Takano, M. Moriya, Y. Iwabuchi, K. Ogasawara, *Tetrahedron Lett.* **1989**, 30, 3805
- [12] Y. Yuasa, J. Ando, S. Shibuya, *J. Chem. Soc., Perkin Trans. 1* **1996**, 465
- [13] Y. Yuasa, J. Ando, S. Shibuya, *J. Chem. Soc., Chem. Commun.* **1994**, 455
- [14] N. A. Sasaki, I. Sagnard, *Tetrahedron* **1994**, 50, 7093
- [15] R. Huisgen, In *1,3-Dipolar Cycloadditions - Introduction, Survey, Mechanism*, A. Padwa, Ed., Wiley, New York 1984, Vol.1, p. 1
- [16] C. Wittland, M. Arend, N. Risch, *Synthesis* **1996**, 367
- [17] C. Wittland, U. Flörke, N. Risch, *Synthesis* **1997**, 1291
- [18] R. V. C. Carr, R. V. Williams, L. A. Paquette, *J. Org. Chem.* **1983**, 48, 4976
- [19] S. Cossu, O. De Lucchi, R. Durr, F. Fabris, *Synth. Commun.* **1996**, 26, 211
- [20] R. C. Bernotas, R. V. Cube, *Synth. Commun.* **1990**, 20, 1209
- [21] A. J. Speziale, E. G. Jaworski, *J. Org. Chem.* **1960**, 25, 728

Address for correspondence:
 Prof. Dr. Nikolaus Risch
 Universität-GH Paderborn
 Fachbereich für Chemie und Chemietechnik
 Warburger Str. 100
 D-33098 Paderborn
 Fax: Internat. code (0) 5251 603245
 e-Mail: nr@chemie.uni-paderborn.de