

Efficient Synthesis of 2-Substituted Indoles Based on Palladium(II) Acetate/Tri-*tert*-butylphosphine-Catalyzed Alkynylation/Amination of 1,2-Dihalobenzenes

Zhen-Yu Tang, Qiao-Sheng Hu*

Department of Chemistry, College of Staten Island and the Graduate Center of the City University of New York, Staten Island, New York 10314, U.S.A.
Fax: (+1)-718-982-3910, e-mail: qiaohu@mail.csi.cuny.edu

Received: January 22, 2006; Accepted: March 18, 2006


Supporting Information for this article is available on the WWW under <http://asc.wiley-vch.de/home/>.

Abstract: A simple, efficient palladium(II) acetate/tri-*tert*-butylphosphine-catalyzed indole synthesis from *o*-alkynylhalobenzenes, readily available from 1,2-dihalobenzenes, has been achieved in good to excellent yields. A one-pot, Pd(OAc)₂/*t*-Bu₃P-catalyzed sequential reaction from 1-chloro-2-iodobenzene, phenylacetylene and amines has also been achieved in good yields.

Keywords: alkynylation; amination; 1,2-dihalobenzenes; indoles; palladium; phosphane ligands

Many natural products and biologically active compounds contain indole skeleton as their structural subunits.^[1] Such a prevalence stimulated the development of a number of methods for indole synthesis in past decades.^[1–3] Among the methods developed, the palladium-catalyzed transformations of *o*-haloanilines *via* alkynylation to form *o*-alkynylanilines followed by intramolecular hydroamination have been studied intensively and represent a fruitful strategy to access 2-substituted indoles.^[2,3] As part of our program directed to develop new sequential and tandem reactions based on powerful cross-coupling reactions,^[4,5] we became interested in developing sequential/tandem reactions for the effi-

cient synthesis of heterocycles including indoles from readily available starting materials. Based on our analysis of possible routes to the key intermediate *o*-alkynylanilines for the synthesis of 2-substituted indoles, we reasoned that readily available 1,2-dihalobenzenes could be suitable starting materials to access *o*-alkynylanilines *via* an alkynylation followed by an amination sequence (Scheme 1). This possible route appeared to be very attractive considering: (a) due to the electronic nature of alkynyl groups, *o*-alkynylhalobenzenes are expected to be more reactive than strong deactivating amino group-containing *o*-haloanilines, and are thus expected to behave similarly like other *o*-substituted halobenzenes for the Buchwald–Hartwig aminations^[6] and (b) *o*-alkynylhalobenzenes, especially *o*-alkynylchlorobenzenes, have been well-established to be readily accessible from the monoalkynylation of 1,2-dihalobenzenes.^[7,8] Based on the fact that enormous progress has been achieved in Pd(0)-catalyzed amination reactions in recent years and different types of aryl halides including aryl chlorides have been demonstrated as suitable substrates,^[6,9] we believed that the amination of *o*-alkynylhalobenzenes including *o*-alkynylchlorobenzenes could be possible and the strategy of alkynylation followed by amination of 1,2-dihalobenzenes would provide us an efficient and general method to access 2-substituted indoles.^[10] Herein our results are reported.









Scheme 1. Possible routes for the preparation of 2-substituted indoles.

Although the monoalkynylation of 1,2-dihalobenzenes with terminal alkynes to form *o*-alkynylhalobenzenes has been well-established,^[7,8] we were surprised to find that the amination of *o*-alkynylhalobenzenes for the synthesis of 2-substituted indoles was rarely studied.^[10,11] We thus began our study to identify palladium catalyst systems that could efficiently catalyze the amination reaction of *o*-alkynylhalobenzenes. 1-Chloro-2-(2-phenylethynyl)benzene was chosen as the model substrate because of (a) its high yield, selective formation from readily available 1-bromo-2-chlorobenzene (81%) and 1-chloro-2-iodobenzene (95%),^[8] and (b) the consideration that catalyst systems that work for 1-chloro-2-(2-phenylethynyl)benzene which contains a very inert C–Cl bond should also work for other more reactive *o*-alkynylhalobenzenes. A number of commercially available ligands that have been demonstrated as suitable for the amination of aryl halides were screened for the Pd(0)-catalyzed amination of 1-chloro-2-(2-phenylethynyl)benzene with benzylamine, which yielded 1-benzyl-2-phenylindole as the final product after intramolecular hydroamination reaction (Table 1). We found that under the conditions of KO-*t*-Bu as base and toluene as solvent at 110 °C, bidentate bis(2-diphenylphosphinophenyl) ether (DPEphos) and 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (XantPhos) were ineffective ligands (Table 1, entries 1 and 2).^[12] Buchwald's aryldialkylphosphines proved to be excellent ligands for the reaction (Table 1, entries 3 and 4).^[13] An N-heterocyclic carbene, *in situ* generated from 1,3-bis(2,4,6-trimethylphenyl)imidazolium chloride, was also tested and a moderate yield was observed (Table 1, entry 5).^[9c, d] We found that the best result was obtained when (*t*-Bu)₃P was employed as ligand (Table 1, entry 6). Other monodentate ligands such as (Cy)₃P and (*sec*-Bu)₃P were also tested and no desired cyclization product was observed (Table 1, entry 7).

With Pd(0)/P(*t*-Bu)₃ as the catalyst system,^[14] several *o*-alkynylhaloarenes, which were prepared from 1-bromo-2-iodobenzene, 1-bromo-2-chlorobenzene and 1-chloro-2-iodobenzene by following reported methods,^[7,8] were tested for the amination followed by the intramolecular hydroamination and our results are listed in Table 2. We found that 1-bromo-2-(2-phenylethynyl)benzene and 1-chloro-2-(2-phenylethynyl)benzene reacted smoothly with both alkylamines and anilines to give 2-substituted indoles in excellent isolated yields (Table 2, entries 1–6). We also found that although *o*-chlorooctynylbenzene and *o*-chlorohexynylbenzene also reacted with benzylamine and octylamine to give 2-substituted indoles with satisfactory results, they appeared to be less reactive than 1-chloro-2-(2-phenylethynyl)benzene (Table 2, entries 7–10). In the reaction of aniline with *o*-chlorohexynylbenzene, the uncyclized product 2-(1-hexynyl)-*N*-phenylaniline was isolated as the main product (Table 2, entry 11). Our results suggested that alkyl groups, which are more electron-donating

Table 1. Ligand screening.^[a]

			
Entry	Ligands	Reaction Condition ^[b]	Yields [%] ^[c]
1		A	0
2		A	0
3		A	88
4		A	93
5		A	59
6	(<i>t</i> -Bu) ₃ P	A or B	91 – 95
7	(Cy) ₃ P and (<i>sec</i> -Bu) ₃ P	A	0

^[a] Alkyne (1 mmol), amine (1.2 mmol), 3% Pd(OAc)₂, 6% ligands.

^[b] Reaction conditions: A: *t*-BuOK (3 equivs.), toluene (2 mL), 110–120 °C. B: K₃PO₄ (3 equivs.), DMA (2 mL), 130 °C.

^[c] Yields of isolated products.

than the phenyl group, likely through the triple bond, retarded the amination reaction and also slowed down the intramolecular hydroamination process.

With Pd(0)/P(*t*-Bu)₃ as the catalyst system, we have also carried out the indole synthesis in a sequential fashion from 1-chloro-2-iodobenzene, alkynes and amines.^[10] We found that conditions for the first alkynylation step played an important role for the successful sequential reactions. When Et₃N was employed as base for the first alkynylation step, no cyclization product was obtained. Fortunately, when inorganic bases such as Cs₂CO₃ or *t*-BuOK were used, 2-substituted indoles were obtained in good yields for phenylacetylene as the alkyne source (Table 3). However, with 1-octyne as the alkyne source, lower yields of 2-substituted indoles were observed, along with significant amounts of uncyclized *N*-benzyl-2-(1-octynyl)aniline. Such lower yield observation was probably due, at least in part, to the electron-donating alkyl group which might make

Table 2. Pd(0)-catalyzed synthesis of 2-substituted indoles from *o*-alkynylhalobenzenes.^[a]

Entry	X	R ¹	R ² NH ₂	Method ^[b]	Yield [%] ^[c]
1	Br		PhCH ₂ NH ₂	A	95
2	Br		PhNH ₂	A	95
3	Cl		PhCH ₂ NH ₂	B	95
4	Cl		PhNH ₂	B	99
5	Cl		4-MeO-PhNH ₂	B	95 ^[d]
6	Cl			A	99
7	Cl		PhCH ₂ NH ₂	A	85
8	Cl			A	74
9	Cl		PhCH ₂ NH ₂	A	85
10	Cl			A	90
11	Cl		PhNH ₂	A	23 ^[e]

^[a] 3% Pd(OAc)₂, 6% (*t*-Bu)₃P, 1 mmol alkyne, 1.2 mmol amine.^[b] Method A: *t*-BuOK (3 equivs.), toluene (2 mL), 110–12 °C, 14 h; Method B: K₃PO₄ (3 equivs.), DMA (2 mL), 130 °C, 14 h.^[c] Yields of isolated products.^[d] 5% Pd(OAc)₂ and 10% (*t*-Bu)₃P were used.^[e] 2-(1-Hexynyl)-*N*-phenylaniline was obtained in 55% yield.

the amination and hydroamination less effective. We have also attempted to carry out the reaction by loading 1-chloro-2-iodobenzene, phenylacetylene and benzylamine at the same time, but no desired indole product was observed.

In summary, based on our analysis that readily available *o*-alkynylhalobenzenes would be more reactive than strong deactivating amino group-containing *o*-haloanilines and would behave similarly like other *o*-substituted halobenzenes for the Buchwald–Hartwig aminations, we have successfully developed a simple, efficient Pd(OAc)₂/*t*-Bu)₃P-catalyzed indole synthesis from *o*-alkynylhalobenzenes including *o*-alkynylchlorobenzenes. A one-pot, Pd(OAc)₂/*t*-Bu)₃P-catalyzed sequential reaction from 1-chloro-2-iodobenzene, phenylacety-

lene and amines has also been developed. Further investigations on Pd(0)-catalyzed sequential/tandem reactions of 1,2-dihalobenzenes for the synthesis of other types of heterocyclic compounds are currently underway.

Experimental Section

General Procedures for Pd(0)/(*t*-Bu)₃P-Catalyzed Indole Synthesis from *o*-Alkynylchlorobenzene

Method A: Under an N₂ atmosphere, to a mixture of *o*-alkynylchlorobenzene (1 mmol), *t*-BuOK (3 mmol), Pd(OAc)₂ (0.03 mmol) and (*t*-Bu)₃P (0.06 mmol) were added an amine

Table 3. Indole synthesis in an one-pot, sequential fashion.

1). $R \equiv$ (1.2 equivs.),
3% Pd(OAc)₂/6% (*t*-Bu)₃P /
10% CuI, *t*-BuOK or Cs₂CO₃ (5 equivs.),
toluene, 110 °C, 1 hour

2). R'-NH₂ (1.2 equivs.), 110 °C, 12 hours

Entry	R≡	R'-NH ₂	Yields [%] ^[a]
1	Ph≡		64
2	Ph≡		71
3	Ph≡		80
4	Ph≡		47
5			23 ^[b]

^[a] Yield of isolated product.

^[b] *N*-Benzyl-2-(1-octynyl)aniline was observed in 14% yield.

(1.2 mmol) and toluene (2 mL). The resulting mixture was heated to 110–120 °C for 14 hours. After quenching with water, the reaction mixture was extracted with ethyl acetate and the organic layer was washed with brine. Rota-evaporation and flash chromatography on silica gel (hexane: ethyl acetate = 5:95 to 15:85) gave 2-substituted indoles as the product.

Method B: Under an N₂ atmosphere, to a mixture of *o*-alkynylchlorobenzene (1 mmol), K₃PO₄ (3 mmol), Pd(OAc)₂ (0.03 mmol) and (*t*-Bu)₃P (0.06 mmol) were added an amine (1.2 mmol) and dimethylacetamide (2 mL). The resulting mixture was heated to 130 °C for 14 hours. After quenching with water, the reaction mixture was extracted with ethyl acetate and the organic layer was washed with brine. Rota-evaporation and flash chromatography on silica gel (hexane: ethyl acetate = 5:95 to 15:85) gave 2-substituted indoles as the product.

General Procedure for One-Pot, Sequential Preparation of 2-Substituted Indoles

Under an N₂ atmosphere, to a mixture of 1-iodo-2-chlorobenzene (1 mmol), Pd(OAc)₂ (0.03 mmol), CuI (0.1 mmol), *t*-BuOK (3 mmol), and (*t*-Bu)₃P (0.06 mmol) were added an alkyne (1.2 mmol) and toluene (2 mL). After the resulting mixture was heated to 110 °C for 1 hour, an amine (1.2 mmol) was added and the mixture was heated at 110 °C for another 12 h. After quenching with water, the reaction mixture was extracted with ethyl acetate and the organic layer was washed with brine. Rota-evaporation and flash chromatography on silica gel (hexane: ethyl acetate = 5:95 to 15:85) gave 2-substituted indoles as the product.

For detailed procedures and characterization data, see Supporting Information.

Acknowledgements

We gratefully thank the NIH (GM69704) for their generous funding. Partial support from the PSC-CUNY Research Award Programs, and GRTI (The Graduate Research and Technology Initiative) grants are also gratefully acknowledged. ZYT thanks the CUNY for Mario Capelloni Dissertation Fellowships. We also thank Frontier Scientific for its generous gifts of palladium(II) acetate and Combiphos Catalysts, Inc for its generous gifts of tri-*tert*-butylphosphine.

References and Notes

- [1] a) T. L. Gilchrist, *Heterocyclic Chemistry*; Addison-Wesley Longman Limited, Singapore, **1997**; b) J. A. Joule, K. Mills, G. F. Smith, *Heterocyclic Chemistry*; Stanley Thornes Ltd., Cheltenham, **1995**.
- [2] For recent reviews, see: a) S. Cacchi, G. Fabrizi, *Chem. Rev.* **2005**, *105*, 2873–2920; b) G. Leni, R. C. Larock, *Chem. Rev.* **2004**, *104*, 2285–2309; c) D. A. Horton, G. T. Bourne, M. L. Smythe, *Chem. Rev.* **2003**, *103*, 893–930; d) H.-J. Knölker, K. R. Reddy, *Chem. Rev.* **2002**, *102*, 4303–4427; e) R. C. Larock, *J. Organomet. Chem.* **1999**, *576*, 111–124.
- [3] Selected recent examples, see: a) M. Nazaré, C. Schneider, A. Lindenschmidt, D. W. Will, *Angew. Chem. Int. Ed.* **2004**, *43*, 4526–4528; b) L.-P. Sun, X.-H. Huang, W.-M. Dai, *Tetrahedron* **2004**, *60*, 10983–10992; c) L. Ackermann, L. T. Kaspar, C. J. Gschrei, *Chem. Commun.* **2004**, 2824–2825; d) K. B. Hong, C. W. Lee, E. K. Yum, *Tetrahedron Lett.* **2004**, *45*, 693–697; e) C. Koradin, W. Dohle, A. L. Rodriguez, B. Schmid, P. Knochel, *Tetrahedron* **2003**, *59*, 1571–1587; f) B. C. Van Esseveldt, F. L. van Delft, R. de Felder, F. P. J. T. Rutjes, *Org. Lett.* **2003**, *5*, 1717–1720; g) J. C. Torres, R. A. Pilli, M. D. Vargas, F. A. Violante, S. J. Garden, A. C. Pinto, *Tetrahedron* **2002**, *58*, 4487–4492; h) T. Iwaki, A. Yasuhara, T. Sakamoto, *J. Chem. Soc. Perkin Trans. 1* **1999**, 1505–1510; i) C.-Y. Chen, D. R. Lieberman, R. D. Larsen, T. R. Verhoeven, P. J. Reider, *J. Org. Chem.* **1997**, *62*, 2676–2677.
- [4] a) C.-G. Dong, Q.-S. Hu, *J. Am. Chem. Soc.* **2005**, *127*, 10006–10007; b) C.-G. Dong, Q.-S. Hu, *Angew. Chem. Int. Ed.* **2006**, *45*, 2289–2292.
- [5] For recent general reviews on tandem reactions, see: a) L. F. Tietze, N. Rackelmann, *Pure Appl. Chem.* **2004**, *76*, 1967–1983; b) K. C. Nicolaou, T. Montagnon, S. A. Snyder, *Chem. Commun.* **2003**, 551–564; c) P. J. Parsons, C. S. Penkett, A. J. Shell, *Chem. Rev.* **1996**, *96*, 195–206; d) L. F. Tietze, *Chem. Rev.* **1996**, *96*, 115–136.
- [6] For recent reviews, see: a) B. H. Yang, S. L. Buchwald, *J. Organomet. Chem.* **1999**, *576*, 125–146; b) J. P. Wolfe, S. Wagaw, J.-F. Marcoux, S. L. Buchwald, *Acc. Chem. Res.* **1998**, *31*, 805–818.
- [7] a) J. A. Marsden, M. M. Haley, in: *Metal-Catalyzed Cross-Coupling Reactions*, 2nd edn., (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, **2004**, 317–394; b) K. Sonogashira, in: *Comprehensive Organic Syn-*

- thesis, (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, Vol. 3, 521–549.
- [8] a) B. Liang, M. Dai, J. Chen, Z. Yang, *J. Org. Chem.* **2005**, *70*, 391–393; b) I. Bytschkov, H. Siebeneicher, S. Doye, *Eur. J. Org. Chem.* **2003**, 2888–2902; c) M. J. Mio, L. C. Kopel, J. B. Braun, T. L. Gadzikwa, K. L. Hull, R. G. Brisbois, C. J. Markworth, P. A. Grieco, *Org. Lett.* **2002**, *4*, 3199–3202; d) H. Li, J. L. Petersen, K. K. Wang, *J. Org. Chem.* **2001**, *66*, 7804–7810; e) H.-R. Zhang, K. K. Wang, *J. Org. Chem.* **1999**, *64*, 7996–7999.
- [9] Selected recent examples: for Pd(0)/P(*t*-Bu)₃-catalyzed amination of aryl chlorides, see: a) T. Yamamoto, M. Nishiyama, Y. Koie, *Tetrahedron Lett.* **1998**, *39*, 2367–2370; b) J. F. Hartwig, M. Kawatsura, S. I. Hauck, K. H. Shaughnessy, L. M. Alcazar-Roman, *J. Org. Chem.* **1999**, *64*, 5575–5580; for N-heterocyclic carbenes as ligands for C–N bond formation, see: c) S. R. Stauffer, S. Lee, J. P. Stambuli, S. I. Hauck, J. F. Hartwig, *Org. Lett.* **2000**, *2*, 1423–1426; d) J. Huang, G. Grasa, S. P. Nolan, *Org. Lett.* **1999**, *1*, 1307–1309.
- [10] Near the end of our work, Ackermann reported a Pd(0)-catalyzed indole synthesis from *o*-alkynylhaloarenes and dihaloarenes using highly steric hindered N-heterocyclic carbenes as ligands: a) L. Ackermann, *Org. Lett.* **2005**, *7*, 439–442; b) L. T. Kasper, L. Ackermann, *Tetrahedron* **2005**, *61*, 11311–11316.
- [11] A two-step, one-pot approach consisting of a hydroamination and a subsequent intramolecular palladium-catalyzed amination reaction has also been reported: H. Siebenreicher, I. Bytschkov, S. Doye, *Angew. Chem. Int. Ed.* **2003**, *42*, 3042–3044.
- [12] Recent examples of DPEphos and XantPhos as ligands for C–N bond formation, see: a) G. B. W. L. Ligthart, H. Ohkawa, R. P. Sijbesma, E. W. Meijer, *J. Org. Chem.* **2006**, *71*, 375–378; b) A. Y. Lebedev, A. S. Kharatulyari, A. Z. Voskoboynikov, *J. Org. Chem.* **2005**, *70*, 596–602; c) S. Cacchi, G. Fabrizi, A. Goggiamani, E. Licandro, S. Maiorana, D. Perdicchia, *Org. Lett.* **2005**, *7*, 1497–1500; d) A. Kuwahara, K. Nakano, K. Nozaki, *J. Org. Chem.* **2005**, *70*, 413–419; e) Y. Chen, X. P. Zhang, *J. Org. Chem.* **2003**, *68*, 4432–4438; f) J. Yin, S. L. Buchwald, *J. Am. Chem. Soc.* **2002**, *124*, 6043–6048; g) J. Yin, S. L. Buchwald, *Org. Lett.* **2000**, *2*, 1101–1104.
- [13] For examples: a) A. Ghosh, J. E. Sieser, M. Riou, W. Cai, L. Rivera-Ruiz, *Org. Lett.* **2003**, *5*, 2207–2210; b) J. P. Wolfe, H. Tomori, J. P. Sadighi, J. Yin, S. L. Buchwald, *J. Org. Chem.* **2000**, *65*, 1158–1174.
- [14] Pd(OAc)₂/(*t*-Bu)₃P catalyst system appears to be more active than that of Pd(OAc)₂/1,3-bis(2,6-diisopropylphenyl)imidazolium chloride system for the preparation of 2-substituted indoles from *o*-alkynylhalobenzenes and 1,2-dihalobenzenes because comparable yields were observed from a 3% Pd(OAc)₂/(*t*-Bu)₃P catalyst loading or a 5–10% Pd(OAc)₂/1,3-bis(2,6-diisopropylphenyl)imidazolium chloride catalyst loading (ref.^[10]). In addition, tri-*tert*-butylphosphine is cheaper than the sterically hindered 1,3-bis(2,6-diisopropylphenyl)imidazolium chloride: For example, the prices from the 2004–2006 Strem's catalog: tri-*tert*-phosphine: \$ 37.0/g; 1,3-bis(2,6-diisopropylphenyl)imidazolium chloride: \$ 58.0/500 mg.