

Ferric Chloride Hexahydrate-Catalyzed Highly Regio- and Stereoselective Conjugate Addition Reaction of 2,3-Allenates with Grignard Reagents: An Efficient Synthesis of β,γ -Alkenates

Guobi Chai,^a Zhan Lu,^a Chunling Fu,^a and Shengming Ma^{a,*}

^a Laboratory of Molecular Recognition and Synthesis, Department of Chemistry, Zhejiang University, Hangzhou 310027, Zhejiang, People's Republic of China
Fax: (+86)-21-6260-9305; e-mail: masm@mail.sioc.ac.cn

Received: February 10, 2009; Revised: May 2, 2009; Published online: July 7, 2009

 Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.200900091>.

Abstract: Ferric chloride hexahydrate was shown to be an efficient catalyst for the conjugate addition of 2,3-allenates with alkyl-, aryl-, or vinyl-Grignard reagents to synthesize polysubstituted β,γ -unsaturated alkenates with high regio- and stereoselectivity. The *in situ* formed magnesium dienolate may readily react with different electrophilic reagents under dif-

ferent reaction conditions with or without a catalyst to construct an allylic quaternary carbon at the α -position of the ester group and a stereocontrollable retention of the carbon-carbon double bond.

Keywords: allenes; esters; Grignard reagents; iron; Michael addition

Introduction

Substituted β,γ -unsaturated alkenates bearing a non-conjugated double bond are widely found in biologically active natural products such as actinarin A methyl ester,^[1a] 4-hydroxycinnamate derivatives,^[1b] cytochalasin Q,^[1c] and, thus, they have caught the attention of many synthetic organic and pharmaceutical chemists.^[1] It is not so easy to synthesize this type of compounds due to the possible migration of the C=C double bond to the conjugated α,β -position and the stereoselectivity referring to the formation of the C=C double bond.^[2] On the other hand, allenes, especially those with electron-withdrawing groups, have become of great interest in synthetic organic chemistry owing to the high reactivity of the allene moiety.^[3] In the past decades, the conjugate addition of organometallic reagents to 2,3-allenates has proven to be an efficient way to synthesize β,γ -unsaturated carbonyl compounds *via* the α -protonation of the *in situ* formed metallic dienolates:^[4] addition reactions of lithium dialkylcuprates or dialkenylcuprate with simple 2,3-allenates^[4a] or 2,3-allenates with functional groups such as hydroxy,^[4b] methoxycarbonyl,^[4c] or ethylthio groups^[4d] afford β,γ -unsaturated alkenates with good regio- and stereoselectivity; the re-

action of an organolithium with 2,3-allenates in the presence of CuCN·2LiCl (0.5 equiv.)^[4e,f] also affords β,γ -unsaturated alkenates with good regio- and stereoselectivity.

However, these reactions are limited to the use of stoichiometric or substoichiometric organocuprate reagents; further improvements in this area are obviously desirable. Particularly, in the view of industrial applications, an environmentally friendly and cheap catalytic system is important. In this context, iron is one of the best choices.^[5,6] In our previous work, we utilized the anhydrous iron salt Fe(acac)₃ to catalyze the conjugate addition reaction of 2,3-allenates with Grignard reagents, affording β,γ -unsaturated alkenates with high regio- and stereoselectivity.^[7]

Compared with Fe(acac)₃, FeCl₃·6H₂O is surely more readily available and extremely cheap. However, there are very limited reports concerning its applications as a catalyst.^[8] In this context, we present our recent surprising observation on the use of simple FeCl₃·6H₂O as an alternative catalyst for the conjugate addition reaction of 2,3-allenates with various Grignard reagents to afford β,γ -unsaturated alkenates. Furthermore, highly stereocontrollable α -acylation and allylation reactions of the *in situ* generated magnesium dienolate intermediate were achieved by


variations of the electrophiles and reaction conditions.^[9,10]

Results and Discussion

Synthesis of 2,3-Allenates 3

2,3-Allenates **3a** and **3e–3o** can be prepared from the corresponding Wittig reagents with acyl chlorides **2** in the presence of Et₃N (Scheme 1).^[11]

2-Alkyl-4-phenyl-2,3-butadienoates **3b–3d** can be prepared from the esterification of corresponding 2,3-allenoic acids **4** (Scheme 2).^[12]



Scheme 2. Synthesis of 2,3-allenoates **3b–3d**.


FeCl₃·6H₂O-Catalyzed Highly Regio- and Stereoselective Conjugate Addition Reaction of 2,3-Allenates with Grignard Reagents

Initially, we accidentally attempted the reaction of ethyl 4-phenyl-2-propyl-2,3-butadienoate **3a** with methylmagnesium chloride under the catalysis of 5 mol% of FeCl₃·6H₂O and the same conditions utilized in our previous work.^[7] To our surprise, ethyl 3-methyl-4-phenyl-2-propyl-3(*Z*)-butenoate **Z-5a** was afforded in 93% yield highly selectively (entry 1, Table 1). The stereochemistry was established by NOE analysis (Figure 1).


Next, 2 mol% of FeCl₃·6H₂O was used and a similar result was obtained (entry 2, Table 1). However, when we reduced the loading of FeCl₃·6H₂O further to 0.5 mol%, the reaction was quite slow, resulting in a very low yield of **Z-5a** (entry 3, Table 1). Only 13%

of **Z-5a** was formed when no catalyst was used with **3a** being recovered in 84% yield (entry 4, Table 1). In all cases, the formation of the *E* isomer of the conjugate addition product was not observed; the α,β-unsaturated conjugated isomer 3-methyl-4-phenyl-2-propyl-2-butenate **6a** was formed to the extent of ≤1.2% (entries 1–4, Table 1). Obviously, compared with the results from Fe(acac)₃, although the formation of α,β-unsaturated enoate **6a** is slightly higher,^[7] FeCl₃·6H₂O is more efficient and high yielding (compare entries 1 and 2 with entries 5 and 6, Table 1).

With the optimized reaction conditions in hand, we explored the substrate scope of the FeCl₃·6H₂O-catalyzed conjugate addition reaction of 2,3-allenoates with alkyl-Grignard reagents as shown in Table 2. R¹



Scheme 1. Synthesis of 2,3-allenoates **3a** and **3e–3o**.


Table 1. Effect of catalyst on the Fe(III)-catalyzed conjugate addition of 2,3-allenoate **3a** with methylmagnesium chloride.^[a]

Entry	FeCl ₃ ·6H ₂ O [mol%]	Yield of Z-5a [%] ^[b]	Recovery of 3a [%] ^[b]	Yield of 6a [%] ^[b]
1	5	93	—	0.7
2	2	98	—	0.7
3	0.5	46	46	1.2
4	—	13	84	0.9
5	5 ^[c]	81	—	0.8

^[a] The reaction was conducted using 0.4 mmol of the 2,3-allenoate, 3 equiv. of the Grignard reagent (3M solution in THF) in 5 mL of toluene at -78°C .

^[b] Determined by ^1H NMR spectroscopy using 1,3,5-trimethylbenzene as the internal standard.


^[c] Fe(acac)₃ was used as catalyst instead of FeCl₃·6H₂O.^[7]

**Figure 1.** NOE study of **Z-5a**.

could be aryl or alkyl groups; R^2 could be H, Bn or alkyl groups; R^3 could be alkyl, Bn or allyl groups. The reaction may proceed with primary (entries 1–12, Table 2), secondary, and tertiary Grignard reagents (entries 13–15, Table 2). However, in some cases such as substrates **3g**, **3i** and **3j**, the reaction requires 5 mol% of FeCl₃·6H₂O for a complete conversion of the starting 2,3-allenoates (entries 7, 9 and 10, Table 2).

The highly regio- and stereoselective addition reaction of 2,3-allenoates with vinylmagnesium halides (entries 1–3, Table 3) or phenylmagnesium chloride (entries 4–7, Table 3) can also be performed to afford the conjugated dienes or 3-phenyl-substituted β,γ -alkenoates. 2-Thienylmagnesium bromide may also be reacted with **3e** to obtain the corresponding product **E-5w** in 63% yield (entry 8, Table 3).


Treatment of the conjugate addition reaction mixture with D₂O afforded **8** in 70% yield with 96% D incorporation, indicating the intermediacy of the magnesium 1,3(*Z*)-dienolate **Z-7a** [Eq. (1)].



Stereocontrollable α -Acylation and Allylation Reactions of the *in situ* Generated Magnesium Dienolate Intermediates with Electrophiles

As a result, the magnesium 1,3-dienolate **7a** may undergo a coupling reaction with acetyl chloride to afford **Z-9a** in 65% yield with an *Z/E* ratio of $>99\%$ (conditions A, entry 1, Table 4). Similarly,^[10] the reaction of the *in situ* generated magnesium dienolate **7a** with allyl acetate at -78°C followed by warming up to room temperature afforded α -acylated 3(*E*)-alkenoate **E-9a** in 50% yield with a reversed stereoselectivity (*E/Z* = 93/7) (conditions B, entry 1, Table 4). The stereochemistry was also established by NOE analysis. It should be noted that the formation of the α -allylated product was not observed based on the ^1H NMR analysis of the crude reaction mixture. The *in situ* generated dienolates from the reactions of **3f**, **3m** and **3o** also reacted with acetyl chloride or allyl acetate affording the corresponding products **Z-9** or **E-9** with high stereoselectivity, respectively (conditions A and B, entries 3, 5 and 7, Table 4). In the *Z*-acylation reactions of the *in situ* generated magnesium dienolate **7**, FeCl₃·6H₂O and Fe(acac)₃ both exhibited very high stereoselectivity ($>99/1$) (conditions A, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 4). However, in the *E*-acylation reactions of the *in situ* generated magnesium dienolate **7**, the results obtained with Fe(acac)₃ are better than those with FeCl₃·6H₂O (conditions B, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 4).

Furthermore, we were again quite happy to observe that under the catalysis of Pd(PPh₃)₄, the reaction of magnesium dienolate **7a** with allyl acetate afforded the α -allylated 3(*Z*)-alkenoate **Z-10a** exclusively in 75% yield with high stereoselectivity (*Z/E* = 98/2) (conditions C, entry 1, Table 5). The configuration of the C=C double bond in this compound was established again by NOE analysis. Likewise, under the catalysis of CuI, the reaction with allyl bromide afforded the opposite stereoisomer **E-10a** in 52% yield

Table 2. FeCl₃·6H₂O-catalyzed conjugate addition of alkylmagnesium chlorides to 2,3-allenoates.^[a]



Entry	R ¹	R ²	R ³	R ⁴	Time [h]	Yield of 5 [%] ^[b]
1	Ph	<i>n</i> -Pr	Et (3a)	Me	2.5	88 (<i>Z</i> - 5a)
2	Ph	<i>n</i> -Pr	Me (3b)	Me	2.4	89 (<i>Z</i> - 5b)
3	Ph	<i>n</i> -Pr	Bn (3c)	Me	3.5	78 (<i>Z</i> - 5c)
4	Ph	Me	allyl (3d)	Me	3	72 (<i>Z</i> - 5d)
5	Ph	Me	Et (3e)	Me	3.5	62 (<i>Z</i> - 5e)
6	<i>p</i> -BrC ₆ H ₄	Me	Et (3f)	Me	3	87 (<i>Z</i> - 5f)
7 ^[c]	<i>p</i> -BrC ₆ H ₄	<i>n</i> -Pr	Et (3g)	Me	2.4	99 (<i>Z</i> - 5g)
8	<i>p</i> -MeOC ₆ H ₄	Me	Et (3h)	Me	1	67 (<i>Z</i> - 5h)
9 ^[c]	<i>n</i> -C ₇ H ₁₅	<i>n</i> -Pr	Et (3i)	Me	3	85 (<i>Z</i> - 5i)
10 ^[c]	<i>n</i> -C ₄ H ₉	Bn	Et (3j)	Me	3	89 (<i>Z</i> - 5j)
11	<i>n</i> -C ₄ H ₉	Me	Et (3k)	Me	2	76 (<i>Z</i> - 5k)
12	<i>n</i> -C ₄ H ₉	H	Et (3l)	<i>n</i> -C ₅ H ₁₁ ^[d]	3.5	61 (<i>Z</i> - 5l)
13	Ph	Me	Et (3e)	<i>c</i> -Hex	2	79 (<i>Z</i> - 5m)
14	<i>n</i> -C ₇ H ₁₅	<i>n</i> -Pr	Et (3i)	<i>s</i> -Bu ^[d]	1.5	85 (<i>Z</i> - 5n)
15	Ph	Me	Et (3e)	<i>t</i> -Bu	1.5	81 (<i>E</i> - 5o)

^[a] The reaction was conducted using 0.4 mmol of 2,3-allenoates, 3 equiv. of the Grignard reagents (solution in THF), and 2 mol% of FeCl₃·6H₂O in 5 mL of toluene at –78 °C, unless otherwise stated.

^[b] Isolated yield.

^[c] 5 mol% of FeCl₃·6H₂O was used.

^[d] Alkylmagnesium bromides were used.

Table 3. FeCl₃·6H₂O-catalyzed conjugate addition of aryl or vinyl magnesium halides with 2,3-allenoates.^[a]


Entry	R ¹	R ²	R ⁴	Time [h]	Yield of 5 [%] ^[b]
1 ^[c]	Ph	<i>n</i> -Pr (3a)	vinyl	1.5	84 (<i>Z</i> - 5p)
2	<i>p</i> -ClC ₆ H ₄	Me (3m)	vinyl ^[d]	3	85 (<i>Z</i> - 5q)
3	<i>n</i> -C ₄ H ₉	Me (3k)	vinyl	3	75 (<i>Z</i> - 5r)
4	<i>p</i> -MeOC ₆ H ₄	Me (3h)	Ph	3	81 (<i>E</i> - 5s)
5	<i>n</i> -C ₄ H ₉	H (3l)	Ph	2.5	64 (<i>E</i> - 5t)
6	<i>n</i> -C ₇ H ₁₅	H (3n)	Ph	3	60 (<i>E</i> - 5u)
7	<i>n</i> -C ₄ H ₉	Me (3k)	Ph	3	70 (<i>E</i> - 5v)
8 ^[e]	Ph	Me (3e)	2-thienyl ^[d]	11	63 (<i>E</i> - 5w)

^[a] The reaction was conducted using 0.4 mmol of 2,3-allenoates, 3 equiv. of the Grignard reagents (solution in THF), and 2 mol% of FeCl₃·6H₂O in 5 mL of toluene at –78 °C. Freshly prepared vinyl-Grignard reagent was needed.

^[b] Isolated yield.


^[c] 5 mol% of FeCl₃·6H₂O was used.

^[d] Magnesium bromide was used.

^[e] This reaction was conducted at room temperature as no reaction occurred at –78 °C.

(*E/Z* = 96/4) (conditions D, entry 1, Table 5). The configuration of the C=C double bond in this compound was also established by NOE analysis. The *in situ* generated dienolates from the reaction of **3f**, **3m** and **3o** also reacted with allyl acetate under the catalysis of

Pd(PPh₃)₄ or allyl bromide under the catalysis of CuI affording the corresponding products *Z*-**10** or *E*-**10** with high stereoselectivity, respectively (conditions C and D, entries 3, 5 and 7, Table 5). In the α -allylation of magnesium dienolate intermediates with allyl ace-

Table 4. Reaction of *in situ* generated magnesium dienolate **7** with acetyl chloride or allyl acetate.^[a]

Entry	R^1	Conditions A		Conditions B	
		Yield of <i>Z</i> - 9 [%] ^[b]	Ratio (<i>Z</i> / <i>E</i>) ^[c]	Yield of <i>E</i> - 9 [%] ^[b]	Ratio [<i>E</i> / <i>Z</i>] ^[c]
1	Ph (3e)	65 (<i>Z</i> - 9a)	> 99/1	50 (<i>E</i> - 9a)	93/7
2 ^[d]	Ph (3e)	68 (<i>Z</i> - 9a)	> 99/1	50 (<i>E</i> - 9a)	95/5
3	<i>p</i> -BrC ₆ H ₄ (3f)	70 (<i>Z</i> - 9b)	> 99/1	45 (<i>E</i> - 9b)	95/5
4 ^[d]	<i>p</i> -BrC ₆ H ₄ (3f)	87 (<i>Z</i> - 9b)	> 99/1	62 (<i>E</i> - 9b)	97/3
5	<i>p</i> -ClC ₆ H ₄ (3m)	82 (<i>Z</i> - 9c)	> 99/1	55 (<i>E</i> - 9c)	96/4
6 ^[d]	<i>p</i> -ClC ₆ H ₄ (3m)	84 (<i>Z</i> - 9c)	> 99/1	50 (<i>E</i> - 9c)	98/2
7	<i>p</i> -FC ₆ H ₄ (3o)	77 (<i>Z</i> - 9d)	> 99/1	50 (<i>E</i> - 9d)	94/6
8 ^[d]	<i>p</i> -FC ₆ H ₄ (3o)	76 (<i>Z</i> - 9d)	> 99/1	69 (<i>E</i> - 9d)	98/2

^[a] **Conditions A:** 2 mmol of acetyl chloride were added to 0.4 mmol of the *in situ* generated magnesium dienolate at $-78^\circ C$. **Conditions B:** 2 mmol of allyl acetate were added to 0.4 mmol of the *in situ* generated magnesium dienolate at $-78^\circ C$ followed by warming up to room temperature naturally. The formation of 6–31% hydrolysis product was observed by 1H NMR analysis of the crude reaction mixture.

^[b] Isolated yield.

^[c] The ratio was determined by 1H NMR analysis of the crude products.

^[d] $Fe(acac)_3$ was used instead of $FeCl_3 \cdot 6H_2O$.

tate under the catalysis of $Pd(PPh_3)_4$, the amounts of α -acylated product 3(*E*)-alkenoate *E*-**9** are in the order of $\leq 0.9\%$, if any, based on the 1H NMR analysis of the crude reaction mixture. In all cases, $FeCl_3 \cdot 6H_2O$ exhibited better or similar stereoselectivity (conditions C and D, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 5).

A possible mechanism has been proposed for the inversion of the configuration of the C=C bond for the formation of *E*-**9** and *E*-**10** (Scheme 3). The dienolate intermediate *Z*-**7** may isomerize to the delocalized π -allylic intermediate, i.e., *anti*-allylic-MgCl, which may isomerize to the more stable *syn*-allylic metallic intermediate due to the steric effect between the coordinated metal and the methyl/ R^1 groups.^[10,13]

The coordination of the C=C bond in the allylic carboxylate or allylic bromide with the metal or the presence of the more bulky Cu^+ instead of Mg^{2+} would further increase this steric effect leading to the isomerization of the *in situ* generated intermediate *Z*-**7** to the thermodynamically more stable *E*-**7**. Subsequent reaction with the allylic carboxylate or allylic bromide leads to the formation of *E*-**9** or *E*-**10**.^[14]

Conclusions

In summary, we have developed an efficient $FeCl_3 \cdot 6H_2O$ -catalyzed method to synthesize trisubstituted β,γ -unsaturated *Z*-alkenoates with high regio-



**Scheme 3.** Possible mechanism for the formation of *E*-**9** and *E*-**10**.

Table 5. Transition metal-catalyzed allylation reactions of *in situ* generated dienolate **7** with different allyl reagents.^[a]


Entry	R ¹	Conditions C		Conditions D	
		Yield of Z-10 [%] ^[b]	Ratio (<i>Z/E</i>) ^[c]	Yield of E-10 (%) ^[b]	Ratio (<i>E/Z</i>) ^[c]
1	Ph (3e)	76 (Z-10a)	98/2	52 (E-10a)	96/4
2 ^[d]	Ph (3e)	67 (Z-10a)	96/4	51 (E-10a)	96/4
3	<i>p</i> -BrC ₆ H ₄ (3f)	70 (Z-10b)	97/3	71 (E-10b)	> 99/1
4 ^[d]	<i>p</i> -BrC ₆ H ₄ (3f)	80 (Z-10b)	93/7	65 (E-10b)	97/3
5	<i>p</i> -ClC ₆ H ₄ (3m)	72 (Z-10c)	97/3	63 (E-10c)	97/3
6 ^[d]	<i>p</i> -ClC ₆ H ₄ (3m)	66 (Z-10c)	96/4	71 (E-10c)	97/3
7	<i>p</i> -FC ₆ H ₄ (3o)	69 (Z-10d)	96/4	69 (E-10d)	98/2
8 ^[d]	<i>p</i> -FC ₆ H ₄ (3o)	74 (Z-10d)	94/6	72 (E-10d)	97/3

^[a] Conditions C: 5 mol% of Pd(PPh₃)₄ and 2 mmol of allyl acetate were added to 0.4 mmol of the *in situ* generated magnesium dienolate at −78 °C followed by warming up to room temperature naturally. Conditions D: 10 mol% of CuI and 2 mmol of allyl bromide were added to 0.4 mmol of the *in situ* generated magnesium dienolate at room temperature.

^[b] Isolated yield (see the experimental section for the details). The formation of 1–13% hydrolysis product was observed by ¹H NMR analysis of the crude reaction mixture.

^[c] The ratio was determined by ¹H NMR analysis of the crude products.

^[d] Fe(acac)₃ was used instead of FeCl₃·6H₂O.

and stereoselectivity. Primary, secondary and tertiary alkyl, vinylic or aromatic groups may be introduced by using the readily available Grignard reagents to the 3-position of 2,3-allenonates. Good to excellent yields were obtained with a catalytic loading of FeCl₃·6H₂O. Compared with the results of Fe(acac)₃, FeCl₃·6H₂O is more efficient and high yielding with slightly higher yields for the formation of α,β-unsaturated enoate (compare entries 1 and 2 with entries 5 and 6, Table 1). Due to the cheap and easy handling nature of the catalyst, this catalytic system will have promising applications in organic synthesis. Furthermore, stereocontrollable α-acylation and allylation reactions of the *in situ* generated magnesium dienolate intermediates **7** were achieved with the use of different electrophiles and conditions: 2-acylated 3(*Z*)- and 3(*E*)-alkenones **9** were formed when we used acetyl chloride or allyl acetate as the electrophiles; 2-allylated 3(*Z*)- and 3(*E*)-alkenones **10** were formed when we used allyl acetate in the presence of Pd(PPh₃)₄ or allyl bromide in the presence of CuI as the electrophiles, respectively. In the *Z*-acylation reactions of the *in situ* generated magnesium dienolates **7**, FeCl₃·6H₂O and Fe(acac)₃ both exhibited very high stereoselectivity

(> 99/1) (conditions A, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 4) while in the *E*-acylation reactions of the *in situ* generated magnesium dienolate **7**, the results obtained with Fe(acac)₃ are better than those with FeCl₃·6H₂O (conditions B, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 4). On the other hand, FeCl₃·6H₂O exhibited better or similar stereoselectivity in the allylation reactions of the *in situ* generated magnesium dienolate **7** (conditions C and D, compare entries 1, 3, 5 and 7 with 2, 4, 6 and 8, Table 5). Further studies in this area are being conducted in our laboratory.

Experimental Section

Ethyl 3-Methyl-4-phenyl-2-propyl-3(*Z*)-butenoate (**Z-5a**); Typical Procedure for the Synthesis of Compounds **5**

To a dried Schlenk tube were added FeCl₃·6H₂O (2.2 mg, 2 mol%, 0.008 mmol), **3a** (91.7 mg, 0.4 mmol), and toluene (5 mL) sequentially under a nitrogen atmosphere at room temperature. A solution of CH₃MgCl in THF (0.4 mL, 3 M,

1.2 mmol, 3 equiv.) was then added with a syringe to the reaction mixture within 3 min at -78°C . After the reaction was over as monitored by TLC, the reaction mixture was quenched slowly with a saturated aqueous solution of NH_4Cl (1 mL) at -78°C , followed by warming up to room temperature naturally and the addition of water (5 mL). After extraction with diethyl ether (30×3 mL), the organic layer was washed sequentially with diluted HCl (5%, aqueous), a saturated aqueous solution of NaHCO_3 , and brine, and dried over anhydrous Na_2SO_4 . Filtration, evaporation, and column chromatography on silica gel (eluent: petroleum ether/ethyl acetate = 100/1) afforded the desired product **Z-5a** as a liquid;^[7] yield: 86.2 mg (88%); ^1H NMR (300 MHz, CDCl_3): δ = 7.36–7.24 (m, 4H), 7.24–7.15 (m, 1H), 6.49 (s, 1H), 4.17 (q, J = 7.2 Hz, 2H), 3.70 (t, J = 7.7 Hz, 1H), 1.84 (s, 3H), 1.80–1.65 (m, 1H), 1.65–1.25 (m, 1H), 1.28 (t, J = 7.2 Hz, 3H), 1.23–1.01 (m, 2H), 0.76 (t, J = 7.4 Hz, 3H).

Ethyl 2-Acetyl-2,3-dimethyl-4-phenyl-3(Z)-butenoate (**Z-9a**); Typical Procedure for the Synthesis of Compounds **Z-9**

To a dried Schlenk tube were added $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (2.6 mg, 2.4 mol%, 0.010 mmol), **3e** (81.8 mg, 0.4 mmol), and toluene (5 mL) sequentially under a nitrogen atmosphere at room temperature. Then a solution of CH_3MgCl in THF (0.4 mL, 3M, 1.2 mmol, 3 equiv.) was added with a syringe to the reaction mixture within 3 min at -78°C . After the reaction was over as monitored by TLC, acetyl chloride (0.14 mL, d = 1.104 g mL^{-1} , 154.6 mg, 2 mmol, 5 equiv.) was slowly added dropwise at -78°C . After the reaction was over (1.5 h) as monitored by TLC, the reaction mixture was quenched slowly with a saturated aqueous solution of NH_4Cl (1 mL) at -78°C , followed by warming up to room temperature naturally. After extraction with diethyl ether (30×3 mL), the organic layer was washed sequentially with diluted HCl (5%, aqueous), a saturated aqueous solution of NaHCO_3 , and brine, and dried over anhydrous Na_2SO_4 . Filtration, evaporation, and column chromatography on silica gel (eluent: petroleum ether/ethyl acetate = 100/1 ~ 20/1) afforded **Z-9a** as a liquid;^[10] yield: 68.7 mg (65%); ^1H NMR (300 MHz, CDCl_3): δ = 7.28–7.08 (m, 5H), 6.65 (s, 1H), 3.90–3.68 (m, 2H), 2.04 (s, 3H), 1.91 (s, 3H), 1.50 (s, 3H), 1.15 (t, J = 7.1 Hz, 3H).

Ethyl 2-Acetyl-2,3-dimethyl-4-phenyl-3(E)-butenoate (**E-9a**); Typical Procedure for the Synthesis of Compounds **E-9**

To a dried Schlenk tube were added $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (2.2 mg, 2 mol%, 0.008 mmol), **3e** (80.9 mg, 0.4 mmol), and toluene (5 mL) sequentially under a nitrogen atmosphere at room temperature. Then a solution of CH_3MgCl in THF (0.4 mL, 3M, 1.2 mmol, 3 equiv.) was added with a syringe to the reaction mixture within 3 min at -78°C . After the reaction was over as monitored by TLC, allyl acetate (0.22 mL, d = 0.928 g mL^{-1} , 204.2 mg, 2 mmol, 5 equiv.) was slowly added dropwise at -78°C followed by warming up to room temperature naturally. After the reaction was over as monitored by TLC, the reaction mixture was quenched with a saturated solution of NH_4Cl (1 mL) at 0°C , followed by warming up to room temperature naturally. After extraction with diethyl

ether (30×3 mL), the organic layer was washed sequentially with diluted HCl (5%, aqueous), a saturated aqueous solution of NaHCO_3 , and brine, and dried over anhydrous Na_2SO_4 . Filtration, evaporation, and column chromatography on silica gel (eluent: petroleum ether/ethyl acetate = 100/1 ~ 20/1) afforded **E-9a** as a liquid;^[10] yield: 51.8 mg (50%); ^1H NMR (300 MHz, CDCl_3): δ = 7.35 (t, J = 7.5 Hz, 2H), 7.25 (t, J = 7.5 Hz, 3H), 6.41 (s, 1H), 4.26 (q, J = 7.1 Hz, 2H), 2.30 (s, 3H), 1.88 (d, J = 1.2 Hz, 3H), 1.61 (s, 3H), 1.31 (t, J = 7.1 Hz, 3H).

Ethyl 2-Allyl-2,3-dimethyl-4-phenyl-3(Z)-butenoate (**Z-10a**); Typical Procedure for the Synthesis of Compounds **Z-10**

To a dried Schlenk tube were added $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (1.9 mg, 1.8 mol%, 0.008 mmol), **3e** (81.8 mg, 0.4 mmol), and toluene (5 mL) sequentially under a nitrogen atmosphere at room temperature. Then a solution of CH_3MgCl in THF (0.4 mL, 3M, 1.2 mmol, 3 equiv.) was added with a syringe to the reaction mixture within 3 min at -78°C . After the reaction was over (3 h) as monitored by TLC, to the resulting reaction mixture were added sequentially $\text{Pd}(\text{PPh}_3)_4$ (22.4 mg, 0.02 mmol, 5 mol%) and allyl acetate (0.22 mL, d = 0.928 g mL^{-1} , 204.2 mg, 2 mmol, 5 equiv., dropwise) at -78°C , followed by warming up to room temperature naturally. After the reaction was over as monitored by TLC, it was quenched with a saturated aqueous solution of NH_4Cl (1 mL) at 0°C , followed by warming up to room temperature naturally. After extraction with diethyl ether (30×3 mL), the organic layer was washed sequentially with diluted HCl (5%, aqueous), a saturated aqueous solution of NaHCO_3 , and brine, and dried over anhydrous Na_2SO_4 . Filtration, evaporation, and column chromatography on silica gel (eluent: petroleum ether/ CH_2Cl_2 = 8/1 ~ petroleum ether/ethyl acetate = 100/1) afforded **Z-10a** as a liquid;^[10] yield: 79.1 mg (76%); ^1H NMR (300 MHz, CDCl_3): δ = 7.29–7.22 (m, 2H), 7.21–7.14 (m, 1H), 7.13–7.08 (m, 2H), 6.49 (s, 1H), 5.77–5.61 (m, 1H), 5.07–4.97 (m, 2H), 3.61 (dq, J_1 = 10.8 Hz, J_2 = 7.2 Hz, 1H), 3.50 (dq, J_1 = 10.8 Hz, J_2 = 7.2 Hz, 1H), 2.52–2.35 (m, 2H), 1.92 (d, J = 1.8 Hz, 3H), 1.28 (s, 3H), 1.06 (t, J = 7.2 Hz, 3H).

Ethyl 2-Allyl-2,3-dimethyl-4-phenyl-3(E)-butenoate (**E-10a**); Typical Procedure for the Synthesis of Compounds **E-10**

To a dried Schlenk tube were added $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (2.1 mg, 1.9 mol%, 0.008 mmol), **3e** (80.2 mg, 0.4 mmol), and toluene (5 mL) sequentially under a nitrogen atmosphere at room temperature. Then a solution of CH_3MgCl in THF (0.4 mL, 3M, 1.2 mmol, 3 equiv.) was added with a syringe to the reaction mixture within 3 min at -78°C . After the reaction was over as monitored by TLC, the cooling bath was removed and the reaction mixture was stirred at room temperature for 2 h followed by sequential addition of CuI (7.9 mg, 0.04 mmol, 10 mol%) and allyl bromide (0.17 mL, d = 1.390 g mL^{-1} , 236.3 mg, 2 mmol, 5 equiv., dropwise). After the reaction was over as monitored by TLC, it was quenched with a saturated aqueous solution of NH_4Cl (1 mL) at 0°C , followed by warming up to room temperature naturally. After extraction with diethyl ether ($30 \times$

3 mL), the organic layer was washed sequentially with diluted HCl (5%, aqueous), a saturated aqueous solution of NaHCO₃, and brine, and dried over anhydrous Na₂SO₄. Filtration, evaporation, and column chromatography on silica gel (eluent: petroleum ether/CH₂Cl₂ = 8/1) afforded *E*-**10a**^[10] (yield: 42.5 mg) as a mixture of *E*-**10a** and ethyl 2,3-dimethyl-4-phenyl-3(*Z*)-butenoate *Z*-**5e** in a molar ratio by NMR of 94/6, content of *E*-**10a**: 95% by weight and a mixture of *E*-**10a**, *Z*-**5e** and *E*-**5e**, molar ratio by NMR: 86/6/8 (yield: 0.0141 g), content of *E*-**10a**: 88% by weight; combined yield: 52% as a liquid; ¹H NMR (300 MHz, CDCl₃): δ = 7.36–7.28 (m, 2H), 7.27–7.17 (m, 3H), 6.40 (s, 1H), 5.79–5.63 (m, 1H), 5.15–5.04 (m, 2H), 4.16 (q, *J* = 7.1 Hz, 2H), 2.67–2.52 (m, 2H), 1.80 (d, *J* = 1.2 Hz, 3H), 1.37 (s, 3H), 1.25 (t, *J* = 7.1 Hz, 3H).

Acknowledgements

Financial support from National Natural Science Foundation of China (20732005) and the Major State Basic Research Development Program (2006CB806105) is greatly appreciated. Shengming Ma is a Qiu Shi Adjunct Professor at Zhejiang University. We thank Mr. W. Kong in this group for reproducing the results presented in entries 3, 13, and 15 of Table 2, entries 2 and 8 of Table 3, entry 4 of Table 4 and entry 3 of Table 5.

References

- [1] a) S. Cao, C. Foster, J. S. Lazo, D. G. I. Kingston, *Bioorg. Med. Chem.* **2005**, *13*, 5830; b) A. Neudörffer, J.-P. Desvergne, D. Bonnefont-Rousselot, A. Legrand, M.-B. Fleury, M. Langeron, *J. Agric. Food Chem.* **2006**, *54*, 1898; c) M. Chinworrungsee, P. Kittakoop, M. Isaka, A. Rungrat, M. Tanticharoen, Y. Thebtaranonth, *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1965.
- [2] For some recent reports on the synthesis of β,γ-unsaturated alkenoates, see: a) F. Brebion, J.-P. Goddard, L. Fensterbank, M. Malacria, *Org. Lett.* **2008**, *10*, 1917; b) H. Tsuji, T. Fujimoto, K. Endo, M. Nakamura, E. Nakamura, *Org. Lett.* **2008**, *10*, 1219; c) M.-Y. Ngai, A. Barchuk, M. J. Krische, *J. Am. Chem. Soc.* **2007**, *129*, 280; d) E. Rossi, G. Abbiati, O. A. Attanasi, S. Rizzato, S. Santeusano, *Tetrahedron* **2007**, *63*, 11055; e) T. Sanada, T. Kato, M. Mitani, A. Mori, *Adv. Synth. Catal.* **2006**, *348*, 51.
- [3] a) *Modern Allene Chemistry*, (Eds.: N. Krause, A. S. K. Hashmi), Wiley-VCH, Weinheim, **2004**; b) S. Ma, *Chem. Rev.* **2005**, *105*, 2829.
- [4] a) J. Berlan, J.-P. Battioni, K. Koosha, *Bull. Soc. Chim. Fr.* **1979**, 3–4, II-183; b) J. G. Knight, S. W. Ainge, C. A. Baxter, T. P. Eastman, S. J. Harwood, *J. Chem. Soc. Perkin Trans. 1* **2000**, 3188; c) T. A. Bryson, D. C. Smith, S. A. Krueger, *Tetrahedron Lett.* **1977**, 525; d) N. Waizumi, T. Itoh, T. Fukuyama, *Tetrahedron Lett.* **1998**, *39*, 6015; e) R. K. Dieter, K. Lu, *Tetrahedron Lett.* **1999**, *40*, 4011; f) R. K. Dieter, K. Lu, S. E. Velu, *J. Org. Chem.* **2000**, *65*, 8715.
- [5] For a general overview on iron catalysis in organic chemistry, see: a) *Iron Catalysis in Organic Chemistry*, (Ed.: B. Plietker), Wiley-VCH, Weinheim, **2008**; b) C. Bolm, J. Legros, J. L. Pailh, L. Zani, *Chem. Rev.* **2004**, *104*, 6217; c) A. Fürstner, R. Martin, *Chem. Lett.* **2005**, *34*, 624; d) S. Enthaler, K. Junge, M. Beller, *Angew. Chem.* **2008**, *120*, 3363; *Angew. Chem. Int. Ed.* **2008**, *47*, 3317; e) A. Correa, O. G. Mancheño, C. Bolm, *Chem. Soc. Rev.* **2008**, *37*, 1108; f) B. D. Sherry, A. Fürstner, *Acc. Chem. Res.* **2008**, *41*, 1500.
- [6] For some recent examples of FeCl₃-catalyzed reactions, see: a) N. Xia, M. Taillefer, *Chem. Eur. J.* **2008**, *14*, 6037; b) J. Bonnamour, C. Bolm, *Org. Lett.* **2008**, *10*, 2665; c) M. Carril, A. Correa, C. Bolm, *Angew. Chem.* **2008**, *120*, 4940; *Angew. Chem. Int. Ed.* **2008**, *47*, 4862; d) B. Alcaide, P. Almendros, T. Martínez del Campo, *Chem. Eur. J.* **2008**, *14*, 7756; e) S.-M. Yiu, W.-L. Man, T.-C. Lau, *J. Am. Chem. Soc.* **2008**, *130*, 10821; f) D. Stadler, T. Bach, *Angew. Chem.* **2008**, *120*, 7668; *Angew. Chem. Int. Ed.* **2008**, *47*, 7557; g) J. Fan, C. Wan, G. Sun, Z. Wang, *J. Org. Chem.* **2008**, *73*, 8608; h) T. Hatakeyama, Y. Yoshimoto, T. Gabriel, M. Nakamura, *Org. Lett.* **2008**, *10*, 5341; i) A. Correa, M. Carril, C. Bolm, *Chem. Eur. J.* **2008**, *14*, 10919; j) K. Wang, M. Lü, A. Yu, X. Zhu, Q. Wang, *J. Org. Chem.* **2009**, *74*, 935; k) W. M. Czaplik, M. Mayer, A. Jacobi von Wangelin, *Angew. Chem.* **2009**, *121*, 616; *Angew. Chem. Int. Ed.* **2009**, *48*, 607; l) T. Zou, S.-S. Pi, J.-H. Li, *Org. Lett.* **2009**, *11*, 453.
- [7] Z. Lu, G. Chai, S. Ma, *J. Am. Chem. Soc.* **2007**, *129*, 14546.
- [8] For some recent examples of reactions utilizing FeCl₃·6H₂O as catalyst, see: a) K. Komeyama, T. Morimoto, K. Takaki, *Angew. Chem.* **2006**, *118*, 3004; *Angew. Chem. Int. Ed.* **2006**, *45*, 2938; b) G. Anilkumar, B. Bitterlich, F. G. Gelalcha, M. K. Tse, M. Beller, *Chem. Commun.* **2007**, 289; c) J. Kischel, K. Mertins, D. Michalik, A. Zapf, M. Beller, *Adv. Synth. Catal.* **2007**, *349*, 865; d) F. G. Gelalcha, B. Bitterlich, G. Anilkumar, M. K. Tse, M. Beller, *Angew. Chem.* **2007**, *119*, 7431; *Angew. Chem. Int. Ed.* **2007**, *46*, 7293; e) F. Shi, M. K. Tse, Z. Li, M. Beller, *Chem. Eur. J.* **2008**, *14*, 8793; f) F. G. Gelalcha, G. Anilkumar, M. K. Tse, A. Brückner, M. Beller, *Chem. Eur. J.* **2008**, *14*, 7687; g) W. Huang, Q. Shen, J. Wang, X. Zhou, *J. Org. Chem.* **2008**, *73*, 1586; h) W. Huang, P. Zheng, Z. Zhang, R. Liu, Z. Chen, X. Zhou, *J. Org. Chem.* **2008**, *73*, 6845.
- [9] Due to the high electron density at C(2) of a metallo-dienolate, reaction at the α-position affording β,γ-unsaturated carbonyl compounds is favored kinetically over reaction at the γ-position affording α,β-unsaturated carbonyl compounds, see: a) I. Fleming, *Frontier Orbitals and Organic Chemical Reactions*, Wiley-Interscience, New York, **1996**, p 45; b) J. L. Herrmann, G. R. Kieczkowski, R. H. Schlessinger, *Tetrahedron Lett.* **1973**, *14*, 2433. For some recent typical examples of reactions of dienolates at the α-position, see: c) V. Reutrakul, J. Jaratjaroonphong, P. Tuchinda, C. Kuhakarn, P. Kongsaeer, S. Prabpai, M. Pohmakotr, *Tetrahedron Lett.* **2006**, *47*, 4753; d) F. A. Davis, H. Qi, G. Sundarababu, *Tetrahedron* **2000**, *56*, 5303; e) B. Bazán-Tejeda, G. Bluet, G. Broustal, J.-M. Campagne, *Chem. Eur. J.* **2006**, *12*,

- 8358; f) P. A. Alexander, S. P. Marsden, D. M. M. Subtil, J. C. Reader, *Org. Lett.* **2005**, 7, 5433; g) K. Fukuhara, H. Urabe, *Tetrahedron Lett.* **2005**, 46, 603; h) A. Otake, A. Yukimasa, J. Watanabe, Y. Sasaki, S. Oishi, H. Tamamura, N. Fujii, *Chem. Commun.* **2003**, 1834.
- [10] Z. Lu, G. Chai, X. Zhang, S. Ma, *Org. Lett.* **2008**, 10, 3517.
- [11] R. W. Lang, H.-J. Hansen, *Org. Synth.* **1984**, 62, 202.
- [12] H. J. Bestmann, H. Hartung, *Chem. Ber.* **1966**, 99, 1198.
- [13] a) C. G. Frost, J. Howarth, J. M. J. Williams, *Tetrahedron: Asymmetry* **1992**, 3, 1089; for some typical reports, see: b) B. Akermark, S. Hansson, A. J. Vitagliano, *J. Am. Chem. Soc.* **1990**, 112, 4587; c) T. Hayashi, A. Yamamoto, T. Hagihara, *J. Org. Chem.* **1986**, 51, 723.
- [14] The radius of Mg^{2+} is 0.65 Å and the radius of Cu^+ is 0.96 Å.
-