

## INDEX

### A

- Ab initio* quantum chemistry
  - average interaction energies, 314
  - clusters of strongly interacting molecules, 341–342
  - mixtures, 314
  - solvent as polarizable continuum, 314
  - solving Schrödinger equation, 316
  - see also* Quantum chemical calculations
- Acetonitrile
  - correlation plots of pair interaction energies, 319
  - critical properties of vapor-liquid equilibria, 321
  - isotropic and anisotropic polarizability, 323–324
  - phase behavior, 321
  - vapor pressure and vapor-liquid coexistence, 320
- Acetylene, pressure-induced polymerization, 387, 388
- Acid catalysts, solid, *see* Zeolite catalysis theory
- Activated carbon, pore size distribution, 220–221
- Activation energies, hydrocarbon, 420
- Activity coefficients
  - predictive methods, 341
  - UNIQUAC and Wilson models, 342, 344–345
- Adam–Gibbs relation, 59–60
- Adsorbates, structure and energetics, 382
- Adsorption isotherms
  - Lennard–Jones methane in silica xerogel, 215
  - mesocarbon microbead (MCMB), 212–213
  - simulation of alkane, 409–411
- Adsorption process
  - models, 205
  - see also* Disordered structure models; Simple geometric pore structure models
- Alkane adsorption and diffusion
  - methods and theory, 407–408
  - simulation, 407–414
  - simulation of adsorption isotherms, 409–411
  - simulation of diffusion, 412–414
  - see also* Zeolite catalysis theory
- Alkanes, *see* Hydrocarbon activation
- Amino acid substitution
  - hierarchical evolution protocol, 105
  - Monte Carlo simulation results, 107
  - schematic diagram, 106
- Amorphous silicon film
  - chemically reactive minority species, 286–290
  - clusters containing Si atoms, 288–290
  - growth mechanism, 280–281
  - mechanism of SiH radical attachment, 286–287
  - SiH penetration mechanism, 287–288
  - surface evolution and structural characterization, 281–283
  - see also* Silicon film growth
- Anisotropic fluctuations, ordered block copolymers, 441–445
- Argon, adsorption isotherms and pore size distribution, 230
- Atomistic density functional theory, 205

### B

- Barrett–Joyner–Halenda (BJH) method, Kelvin equation, 240
- Base substitutions, DNA copying and amplification, 100
- Basin enumeration function
  - description, 60–61
  - isobaric, 62, 63
  - obtaining from experimental data, 61–62
  - schematic, 65
- Basis set expansion, Car–Parrinello, 362–363
- Basis set superposition error (BSSE), 317
- Bicontinuous networks, *see* Microemulsions

- Bicontinuous phase reactions  
 anomalous mean-field (AMF) kinetics, 129  
 diffusion equations, 127–128  
 dynamics of particle, 130  
 effect of thermal fluctuations of, 135–136  
 flow equations, 133–134  
 Kubo formula, 130  
 Langevin equation for particle in potential field, 130  
 mean-field analysis, 129–132  
 objectives, 128–129  
 physical implications of kinetics, 134  
 presence of potential, 129–132  
 probability distribution, 130–131  
 renormalization group theory (RG), 132–134  
 result counterintuitive to expectations, 134–135  
 self-consistent analysis, 131–132
- Bimolecular reactions, droplet phase, 136–137
- Binary collision theory, ion-surface scattering, 179–180
- Biological systems, Car–Parrinello, 389–392
- Blends, *see* Homopolymer solutions and blends
- Block copolymers  
 anisotropic fluctuations in ordered phases, 441–445  
 approximate theory for anisotropic fluctuations, 442–443  
 composition, 439–440  
 computer simulation by Ginzburg–Landau approach, 446  
 dominant fluctuation modes in HEX phase, 444, 445  
 fluctuation in order parameter, 442  
 hyperparallel tempering Monte Carlo (HPTMC), 17–18  
 kinetic paths of order-order/order-disorder transitions, 445–450  
 long-wavelength deformations/ fluctuations, 452–454  
 long-wavelength fluctuations and instabilities, 452–456  
 long-wavelength instability in ordered phases, 454–456  
 metastable states from preordered states, 459  
 method for calculating matrix at equilibrium, 442  
 microphase ordering, 440  
 microphase separation, 439  
 morphology, 441  
 morphology and metastability in ABC triblock, 456–459  
 nature and stability of nonclassical phases, 450–452  
 nucleation/growth kinetics in order-order/order-disorder, 449–450  
 phase diagram, 18  
 proliferation of HEX cylinder orientations, 448–449  
 scattering function of lamellar phase, 443–444  
 stability of PL morphologies, 451–452  
 thermodynamic stability of G phase, 450–451  
 three pair interactions in ABC triblock, 458–459  
 transition from two- to three-domain structure in triblock, 458  
 two-dimensionally ordered ABC triblock phases, 457
- Bonding, zeolite framework, 405–406
- Born–Oppenheimer molecular dynamics, Car–Parrinello, 368–370
- Born repulsion expression, 406–407
- Bulk materials, processes, Car–Parrinello, 376–378

## C

- Carbon tetrachloride, interaction model, 324
- Car–Parrinello methods  
 advances in methodology, 392–393  
 basis set expansion, 362–363  
 Born–Oppenheimer molecular dynamics, 368  
 chain termination and branching in olefin polymerization, 38  
 choice of model, 363  
 classes of problems best applicable, 356–357  
 classical molecular dynamics, 357–358  
 coupling two methanol at acid site of chabazite, 385–386  
 cubic supercell of water molecules, 365

- density functional theory (DFT), 358–362  
DFT to compute electronic state, 357  
diffusivities of Si, Mg, and Li in conductor, 376–377  
disadvantage of path integral methods, 392–393  
dynamics of proton motion in aspartyl dyad, 390–392  
exposing chemical engineering community, 355–356  
formation of defects in GaAs, 371–372  
gas-phase processes, 372–376  
Grand Challenge Problem, 354  
Hartree–Fock method for single electron equation, 358–359  
Hellman–Feynman theorem, 368  
heterogeneous processes on surfaces, 382–386  
history, 353–354  
infrared spectrum of D<sub>2</sub>O, 380, 381  
interpretation, 354–355  
Kohn–Sham approach, 361–362  
many-body electronic system via electronic density, 359  
many-electron system by Hohenberg and Kohn, 359–360  
melting point of solid materials, 387, 389  
methanol adsorption at activity sites of chabazite, 371  
molecular dynamics, 368–370  
new algorithms and methods, 392  
objectives, 356–357  
periodic boundary conditions of plane waves, 363, 367  
phases of electron transfer process, 372–373, 374  
phase transitions, 386–389  
plane waves form, 363  
polymerization reactions on Ziegler–Natta catalysts, 383–385  
pressure-induced acetylene polymerization, 387, 388  
processes in biological systems, 389–392  
processes in bulk materials, 376–378  
propagating orbitals with nuclei without reoptimizing, 369  
properties of liquids, solvation, and reactions in liquids, 378–382  
pseudopotential method, 367  
purposes of simulations, 371  
reactivity of ketones with OH using, 373–375, 376  
slab model of platinum(111) surface, 366  
sliding of grain boundaries in germanium by *ab initio*, 377  
structural and bonding properties of water via supercells, 378–380  
structure and dynamics of aqueous proton transfer, 382  
superposition of oscillations, 369–370  
unit cell of zeolite chabazite, 364  
Catalysts, *see* Zeolite catalysis theory  
Catalytic reaction, rate, 401–403  
Chabazite  
coupling two methanol molecules, 385–386  
energy of adsorption of protonated propylene, 425  
propene during protonation attack, 416, 417  
methanol adsorption, 371  
stability of isobutene activation, 423  
unit cell, 364  
Charging free energy  
continuum solvation models, 327–328  
contributions, 337–338  
Chemical reactivity, microemulsions, 125  
Chem-Shell embedding procedure, 422–424  
Chlorine ions, *see* Ion-assisted etching  
Clusters  
formation in surfactant solutions, 308  
impact on film defects, 290  
interactions of Si<sub>6</sub>H<sub>13</sub> with a-Si:H film, 289  
neutral and cationic with Si atoms, 288–289  
structure of Si<sub>6</sub>H<sub>13</sub>, 289  
Collision cascade, 157  
Combinatorial chemistry  
goal, 81–82  
steps of first round, 89  
task of searching composition space, 82  
*see also* Materials discovery; Protein molecular evolution  
Complex fluids, molecular simulation, 1–2  
Computational quantum chemistry  
isolated molecule, 313  
*see also* Quantum chemical calculations  
Condensed phases, topographic viewpoint, 34–35  
Controlled pore glasses (CPG), disordered structure models, 206–209  
Copolymers, *see* Block copolymers

- Copolymers, block and random,  
hyperparallel tempering, 17–18  
Corradini site, Ziegler–Natta catalysts,  
383–384  
Critical micelle concentration (cmc)  
determination, 306–307  
reduced osmotic pressure vs. surfactant,  
306  
versus inverse temperature for systems  
forming micelles, 307  
Crystal nucleation, pure melt, 38
- D**
- Debye equation, rotational motion, 31  
Density functional theory (DFT)  
adsorption models, 225–231  
Car–Parrinello simulations, 357, 358–362  
catalytic reaction rate, 402–403  
choice of weighting function, 227–228  
comparing to GEMC model isotherms, 229  
DFT excess isotherm, 228  
energy minimization step, 343  
equilibrium density profile, 228  
model isotherms for various gases, 228–229  
pore filling pressure and pore width, 226  
pore filling pressure correlations for  
MK-BET and DFT, 243  
pore size distribution results, 231  
radical-surface interactions, 257–258  
test of robustness, 229–230  
Deposition  
ion-surface interactions, 180–198  
*see also* Ion-assisted etching  
Desorption isotherms, methane in silica  
xerogel, 215  
Diffusion equations  
bicontinuous phase, 127–128  
droplet phase reactions, 139–141  
simulation, 412–414  
Disease organism, evolved resistance, 117  
Disilane formation, initial stages of film  
growth, 278–279  
Disordered structure models  
carbon-carbon radial distribution  
functions (RDFs), 211  
experimental RDFs, 209  
grand canonical Monte Carlo (GCMC),  
214, 216  
geometric definition of pore size  
distribution, 208  
geometric pore size distribution, 207–208  
illustration of porous matrix via  
templating, 216  
microporous carbons, 209–213  
model porous glasses via quench  
molecular dynamics (MD), 207  
Percus–Yevick closure, 217  
perturbations of GCMC simulation,  
213–214  
physical properties vs. reverse Monte  
Carlo (RMC), 212–213  
porous glasses, 206–209  
replica Ornstein–Zernicke (ROZ) integral  
equations, 217–218  
RMC, 209–211  
templated porous materials, 216–218  
visualizations from GCMC of confined  
fluids in xerogels, 215  
xerogels, 213–216  
*see also* Simple geometric pore structure  
models  
Dissociative adsorption, Si incorporation,  
274–275  
DNA, mutation tendencies, 111, 112  
DNA base mutation, protein expression,  
111, 113  
DNA base substitution, local protein space,  
115  
DNA shuffling  
connection between codon usage and, 113  
discovery, 100–101  
genes and operons evolved by, 101  
hierarchical evolution protocol, 105, 107  
incapable of evolving new protein  
folds, 116  
Monte Carlo simulation results, 107  
schematic diagram, 106  
DNA swapping  
evolution of *E. coli* from *Salmonella*, 115  
penicillin-binding proteins, 115  
Dominant precursor  
deposition, 284–286  
SiH<sub>3</sub> for silicon film growth, 285–286  
Droplet phase reactions  
analysis of leading order corrections,  
145–146  
bimolecular reactions  $A + B \xrightarrow{\zeta} 0$ , 136  
considering temporal regimes, 141–143

- diffusion equation and perturbation  
expansion, 139–141  
dynamical fluctuations enhancing rate,  
143  
effect of Péclet number, 144–145  
effects for future work, 146  
fluctuations, 137–138  
intermediate times, 143  
short time regime, 143–144  
steady-state limit of expression, 142
- Dubinin adsorption models  
Dubinin and Astakhov (DA) equation,  
236–237  
Dubinin–Radushkevitch (DR) equation,  
236  
Dubinin–Stoeckli (DS) method, 237  
pore size distribution (PSD) comparisons,  
238  
semiempirical, 205, 236–239
- E**
- Electrolyte solutions  
binodal curves for asymmetric ionic  
systems, 8–9  
clusters from simulation of  $\lambda = 0.1$  system,  
10  
critical parameters as function of size  
asymmetry, 9  
fraction of ions in clusters, 11  
hyperparallel tempering Monte Carlo  
(HPTMC), 7–11  
integral-equation theoretical predictions,  
9–10  
parallel tempering methods, 8  
primitive model, 7–11
- Eley–Rideal reaction, hydrogen abstraction,  
277
- Embedding techniques  
Chem-Shell, 422–424  
hydrocarbon activation, 422–424
- Empirical interatomic interactions, 258–260
- Energy landscape  
application of landscape-based ideas,  
34–35  
constant volume and pressure, 38–39  
crystal nucleation from pure melt, 38  
deepest basin in each symmetry sector,  
35–36  
identification of inherent structures, 35  
insight into stretched liquids and glassy  
state, 67–68  
N-body potential energy function, 33–34  
partitioning configuration space, 35  
potential energy range, 36  
potential enthalpy, 39  
schematic for many-particle system, 34  
single-sector distribution, 37–38  
steepest descent mapping, 35  
supercooled liquids and glasses, 33–39  
temperature variations, 37  
thermal equilibrium, 37–38  
topographic viewpoint of condensates  
phases, 34  
vibrational partition function for each  
basin, 36–37  
void geometry and connections to, 40–45
- Enthalpy  
potential, 39  
relationship to temperature, 26
- Entropy  
configurational, 59–60  
liquid versus crystal, 29, 30
- Etching, *see* Ion-assisted etching
- Ethanol in heptane, infinite dilution activity  
coefficient, 333
- Ethylene polymerization, Ziegler–Natta,  
382–385
- Exchange energy parameters, surfactants,  
300–301
- F**
- Film growth, *see* Silicon film growth
- Flexible polymers  
liquid-liquid phase diagram, 17  
phase diagram, 16  
*see also* Polymers
- Fluctuations  
droplet phase reactions, 137–138  
effect of, of bicontinuous phase on  
kinetics, 135–136
- Fluorine ions, *see* Ion-assisted etching
- Frederickson–Andersen kinetic Ising model,  
55
- Fredholm integral equations  
adsorption models, 219–220  
circumventing problems by, 220

**G**

- GaAs, formation of defects, 371–372
- Gas-phase processes
- Car–Parrinello, 372–376
  - formaldehyde radical anion with methyl chloride, 372, 373
  - phases of electron-transfer process, 372–373, 374
  - reactivity of ketones with OH, 373–375, 376
- Gaussian landscapes
- 1-propanol and 3-methylpentane, 66
  - relations for systems with, 66–67
  - thermodynamic properties of macroscopic system with, 66–67
- GCSKOW model, *see* Group contribution solvation (GCS) model
- Geometric pore size distribution
- advantage of definition, 208
  - determination for porous glasses, 207–208
  - disordered porous carbon model, 212
  - two-dimensional illustration, 208
  - see also* Pore size distribution (PSD)
- Geometry
- algorithm determining volume and surface area, 40–41
  - analysis of void space in inherent structures, 43
  - average pressure in inherent structures, 42–43
  - complexity of hydrogen bonds in aqueous systems, 45
  - expanding system to lower densities, 43–44
  - morphology of Lennard–Jones inherent structures, 42
  - question from Lennard–Jones fluid results, 44
  - shape of pressure versus density curve, 44
  - shifted-force Lennard–Jones system, 42
  - supercooled liquids and glasses, 39–40
  - void, 40–45
  - Voronoi–Delaunay tessellation, 41
- Germanium, sliding of grain boundaries, 377
- Gibbs ensemble Monte Carlo (GEMC)
- advantage over grand canonical Monte Carlo (GCMC), 225
  - comparing to DFT isotherms, 229
  - critical properties of vapor-liquid equilibria, 321
  - isotherms for nitrogen adsorption, 225, 226
  - molecular simulation, 204
  - phase equilibria of confined fluids, 224
  - pore filling pressure and pore width, 226
  - potentials for predicted phase behavior, 319–321
  - simulation adsorption model, 222, 224–225
- Glass transition temperature,  $T_g$
- definition, 24
  - dependence on cooling rate, 24–25
  - transformation range, 25–26
- Glasses
- chemical engineering science, 22–23
  - commercial processes and applications, 22, 71
  - see also* Supercooled liquids and glasses
- Gram–Schmidt procedure, subspace identification, 90–91
- Grand canonical Monte Carlo (GCMC)
- GCMC adsorption simulation method, 214, 216
  - GCMC with percolation theory, 222
  - isotherms for adsorption of bases, 222
  - Lennard–Jones methane in xerogel, 215
  - molecular simulation, 204
  - perturbations for xerogels, 213
  - simulation of xerogels, 213–216
  - visualizations of confined fluids in xerogels, 215
- Grand Challenge Problem, multiscale modeling, 354
- Group contribution methods
- correcting for structure and proximity effects, 336
  - failure from distortion of electron distribution, 336
  - influence of proximity effects, 339–341
  - physical and thermodynamic properties, 335
  - shortcomings, 335–336
- Group contribution solvation (GCS) model
- derivation for octanol-water partition coefficient (GCSKOW), 332
  - GCSKOW predictions using simple group contribution, 337
  - GCSKOW predictions with multipole corrections, 337, 338–339
  - infinite dilute activity coefficients, 330–332
  - scale factor and temperature dependence, 334–335

Group scale factor, functional groups in different solvents, 329

## H

Halogen ions for etching, *see* Ion-assisted etching

Hartree–Fock (HF) method  
energy calculations, 344  
energy minimization, 343  
Schrödinger to single-electron equation, 358–359

Hellman–Feynman theorem, 368

Hexane, isomerization, 430–431, 432

Hexene, isomerization reaction, 428–430

Hierarchical approach, radical-surface interactions, 255–257

HIV-1 protease, Car–Parrinello, 390–392

Homopolymer chains

mole fraction in dilute phase vs. inverse temperature, 303

phase behavior, 303

surfactant solution simulation, 302–304

Homopolymer solutions and blends

coexistence curves, 13–14

compressibility effects of phase behavior, 14–15

end-to-end vector autocorrelation function, 12

hyperparallel tempering Monte Carlo (HPTMC), 11–15

performance of hyperparallel tempering, 12–13

phase diagram of asymmetric blends, 15

phase diagram of long chains, 13

scaling critical density with chain length, 14

Horvath–Kawazoe (HK) method

adsorbate-adsorbate interaction, 233

causes for differences in model results, 234

dispersion coefficients using

Kirkwood–Muller, 233

illustration of modified HK, 236

limitation, 235

modified HK, 235

pore condensation pressure and slit pore width, 233–234

pore filling correlation, 234

pore filling pressure and pore width, 226

semiempirical adsorption model, 231, 233–234

semiempirical models, 205

Hydride transfer reaction, transition states, 426–427

Hydrocarbon activation

activation energies, 420

activation energies for proton-activated isobutane, 419

energy of adsorption for protonated propylene, 425

Chem-Shell embedding procedure, 422–424

corrections to activation energies for cluster acid strength, 421

elementary reaction steps, 414, 415

embedding techniques, 422

energy differences using Monte Carlo, 427

formation of organic cations, 414, 416

geometries of propene protonation, 416, 417

geometries of zeolitic site clusters, 418

corrections to cluster calculations, 424, 426

properties of zeolitic protons, 416, 418–419

quantum-chemical calculations, 416

stability of *i*-butene activation, 423

transition-state energies, 420–421

transition states of hydride transfer, 426–427

*see also* Zeolite catalysis theory

Hydrogen abstraction

Eley–Rideal reaction, 277

Langmuir–Hinselwood mechanism, 277–278

SiH<sub>3</sub> abstracting from H-terminated Si surface, 276

silicon film growth, 276–278

Hydrogen bonding, configurational freedom, 346

Hydroisomerization reaction, kinetics, 427–428

Hyperparallel tempering Monte Carlo

acceptance rate for trial swap, 19

advantages, 18

block and random copolymers, 17–18

factors affecting performance, 19

grand canonical ensemble, 3

homopolymer solutions and blends, 11–15

Lennard–Jones fluid, 5–7

methodology, 3–5

- Hyperparallel tempering Monte Carlo (*continued*)  
 partition function of composite ensemble, 3  
 preweighting function, 4  
 primitive model electrolyte solutions, 7–11  
 schematic of implementation, 5  
 semiflexible polymers and blends with flexible, 15–17  
 swapping, 4–5  
 three types of trial moves, 4  
 weighting function, 4
- I**
- Ideal glass, description, 64  
 Immune system, evasion of, 117  
 Infinite dilution activity coefficients  
   comparing predictions, 331  
   continuum solvation models, 326–327  
   ethanol in heptane, 333  
   group contribution solvation model, 330–332  
 Infrared spectrum, deuterated water, 380, 381  
 Interatomic interactions, radical-surface, 258–260  
 Intermediate time regime, droplet phase, 143  
 Ion-assisted etching  
   Ar from bare Si surface, 177  
   arguments about mechanisms, 161–162  
   atomic configuration after fluence of  $\text{CF}_3^+$ , 185  
   average product molecular weight vs. ion energy, 190  
   average yield of sputtered species per ion, 188  
   binary collision (BC) theory, 179–180  
   bond energy distribution for F/Si top layer, 166  
   C and F content as function of  $\text{CF}_3^+$  fluence, 183  
   C, F, and Si depth profiles for  $\text{CF}^+$  bombardment, 196, 197  
   cascade energy distribution, 162–163  
    $\text{CF}^+$  and  $\text{CF}_2^+$  deposition on Si, 190–198  
    $\text{CF}_3^+$  etching of silicon, 182–186  
   chlorosilyl layer from Cl bombardment, 173  
   coordinate system and angle definition, 176  
   decomposition yield of F and C vs.  $\text{CF}^+$  and  $\text{CF}_2^+$ , 195  
   deposition and etching, 180–198  
   depth profiles for Si, F, and C after  $\text{CF}_3^+$ , 186  
   distributions of total scattering angle, 181  
   etch product flux versus energy, 163  
   etch yield and fluorocarbonsilyl thickness vs.  $\text{CF}_3^+$ , 187  
   etch yield vs. ion energy from MD simulation, 167  
   fluorosilyl layer from F bombardment, 171  
   experimental studies of mechanisms, 162–164  
   fluorine coverage vs.  $\text{F}^+$  fluence, 170  
   flux probability density, 163  
   interfacial  $\text{Si}_x\text{C}_y\text{F}_z$  thickness vs.  $\text{CF}^+$  and  $\text{CF}_2^+$ , 198  
   ion impact and scattering, 174  
   ion-surface scattering dynamics, 172, 174–180  
   kinetic energy distribution of sputter products, 168  
   mechanisms, 161  
   molecular dynamics studies of, mechanisms, 164–172  
   perfluorinated silicon layers with  $\text{Ar}^+$ , 165, 166  
   polar angle scattering, Cl-covered Si, 177–178  
   product angular distributions, 192  
   product distributions in  $\text{CF}_3^+$  etching, 186–190  
   product kinetic energy distributions, 191  
   reflected energy fraction vs. total scatter angle, 179  
   role of surface roughness, 174–178  
   results of MD, 172  
   second stage of simulation, 172  
   Si surface content as function of ion fluence, 184  
   Si surface requiring amount of halogen, 167–168  
   side and top view of surface layers, 175  
   silicon etching with F or Cl, 169  
   square wave modulated ion beam and flux of  $\text{XeF}_2$ , 163–164



- steady-state etching of Si by energetic F or Cl, 168–169, 171
- sticking coefficients of F and C vs.  $\text{CF}^+$  and  $\text{CF}_2^+$ , 195
- uptake of F and C as function of fluence of  $\text{CF}^+$ , 193
- uptake of F and C as function of fluence of  $\text{CF}_2^+$ , 194
- views of Si–Cl surface layers, 176
- yields of major products versus ion energy, 189
- see also* Molecular dynamics (MD)
- Ion-surface interactions**
- collision cascade, 157
- deposition and etching, 180–198
- heating from energetic ion impact, 159–160
- hierarchical scheme for atomistic simulations, 159
- molecular dynamics, 156–161
- plasma processing, 155–156
- quantum mechanics and
- Born–Oppenheimer approximation, 158–159
- side view of simulated Cu layer with  $\text{Cu}^+$ , 157
- simulating, 157–158
- simulating long times between impacts, 160–161
- size of cell crucial, 160
- sputtering, 157
- see also* Ion-assisted etching
- Isobutane**, activation energies for
- proton-activated, 419
- Isobutene**, effect of embedded environment on stability of activation, 423
- Isomerization**
- hexane, 430–431, 432
- hexene, 428–430
- Barrett–Joyner–Halenda (BJH) method**, 240
- classical adsorption model, 239
- classical thermodynamic models, 205
- improved model of statistical adsorbed film thickness, 242, 244
- mean pore sizes of adsorbents, 240
- model nitrogen adsorption isotherms via
- modified Kelvin–BET, 243
- modified Kelvin (MK) equation, 239–240
- pore filling correlations for nitrogen in MCM-41, 241
- pore filling pressure correlations for MK–BET and DFT, 243
- pore size distribution (PSD) using BJH method, 239
- PSD for model porous silica glasses, 241
- schematic of vapor–liquid equilibrium in wetted pore, 239
- shortcomings spurring efforts to modify, 240, 242
- Ketones**, reactivity with OH, 373–375
- Kinetic Monte Carlo (KMC)**
- modeling growth process over coarser time scales, 257
- radical-surface interactions, 255–257
- Kinetics**
- hexane isomerization, 430–431
- hydroisomerization reaction, 427–428
- isomerization reaction of hexene, 428–430
- Kohlrausch–Williams–Watts (KWW)**
- relaxation kinetics, 54–55
- temporal behavior of response function, 28–29
- Kohn–Sham approach**, 360–361

**L**

- K**
- Kauzmann curve**, 68–69
- Kauzmann paradox**, violation of Third Law of Thermodynamics, 29–30
- Kauzmann temperature**
- configurational entropy, 59–60
- description, 29, 30
- Kelvin equation**
- Landscape dynamics and relaxation**
- Adam–Gibbs relation, 59–60
- basin shapes and transitions, 52–53
- basins and kinetic pathways, 58–59
- configurational entropy, 59–60
- dielectric relaxation time, 60
- driving supercooled system from equilibrium, 54
- fragile glass formers, 56–57

- Landscape dynamics and  
relaxation (*continued*)  
Fredrickson–Andersen kinetic Ising  
model, 55  
isochoric dynamical evolution of  
N-particle system, 50–51  
kinetics of interbasin transitions and  
relaxation, 53–54  
Kohlrausch–Williams–Watt (KWW)  
stretched exponential, 54–55  
long-pathway rearrangement processes for  
fragile materials, 58  
mean relaxation time, 54–55  
mean-squared atomic displacements vs.  
temperature, 52  
mode coupling theory (MCT), 51–52  
models for supercooling and glass  
formation, 55–56  
shear viscosity measures, 56  
Stokes–Einstein relation, 58  
supercooled liquids and glasses, 50–60  
system configuration point, 51  
tiling model, 55  
topographic differences, 57–58  
transitions between neighboring  
basins, 53  
viscosity for molten silica ( $\text{SiO}_2$ ), 56  
Langmuir–Hinselwood mechanism, H  
abstraction, 277–278  
Length scales, plasma processing, 152–153  
Lennard–Jones fluid  
adsorption and desorption isotherms for  
methane in silica xerogel, 215  
adsorption simulation, 222  
density variation of inherent structure  
pressure, 43  
geometric algorithm exploring  
morphology, 42  
hyperparallel tempering Monte Carlo  
(HPTMC), 5–7  
phase behavior, 5–6  
phase diagram, 7  
phase diagram calculated vs. literature,  
6–7  
question from results, 44  
replica number as function of Monte Carlo  
steps, 6  
shifted-force, 42  
Library design and redesign  
closing loop between, 96–97  
materials discovery, 85–87  
Liquids, *see* Supercooled liquids and glasses  
Lubachevsky–Stillinger protocol,  
preparation method, 49
- ## M
- Many-particle system, energy landscape, 34  
Materials discovery  
acceptance probability of exchanging  
samples, 89  
analogy with Monte Carlo (MC), 85  
closing loop between library design and  
redesign, 96–97  
combinatorial approach, 82  
effectiveness of MC strategies, 95–96  
figure-of-merit landscape, 87  
further development, 96–97  
Gram–Schmidt procedure, 90–91  
grid method, 95  
grid search on composition and  
noncomposition variables, 85, 86  
library design and redesign, 85–87  
low-discrepancy sequence (LDS)  
method, 95  
material optimization or development, 83  
maximum figure of merit for different  
protocols, 96  
MC library design and redesign strategy, 88  
MC protocols, 94–95  
method for changing variables, 87  
parallel tempering as MC extension, 89  
phase points as function of dimension and  
spacing, 93  
random phase volume model, 92–94  
searching multidimensional space, 84  
searching variable space by MC, 87–89  
significance of sampling, 91–92  
simplex of allowed compositions, 89–91  
space of variables, 84–85  
steps of combinatorial chemistry, 89  
swapping, 87, 88  
synthetic approach, 82  
ways of changing variables, 87–88  
*see also* Protein molecular evolution  
MCM-41  
pore filling correlations for nitrogen  
in, 241  
pore size distribution, 231, 232

- Mean-field analysis  
  anomalous mean-field (AMF) kinetics, 129  
  bicontinuous phase, 129–132  
  presence of potential, 129–132  
  self-consistent analysis, 131–132
- Mean-spherical approximation (MSA)  
  theory, 9–10
- Melting point, Car–Parrinello, 387, 389
- Mesocarbon microbead (MCMB), *see*  
  Microporous carbons
- Metastability, ABC triblock copolymers, 456–459
- Methane, activated carbon to, adsorption, 220–221
- Methanol  
  pair interaction energies, 319  
  phase behavior, 321  
  vapor-liquid coexistence diagram, 320  
  vapor-liquid equilibria, 321  
  vapor pressure, 320
- Micelles, *see* Surfactant solutions
- Microemulsions  
  applications utilizing, 125  
  importance of fluctuations, 125  
  microstructures of, phase, 126  
  phase behavior, 124  
  practical utility, 125  
  primary interest in reactions  $A + B \rightleftharpoons 0$ , 125–126  
  reaction kinetics at oil–water interface, 126  
  reactions in bicontinuous phase, 127–136  
  reactions in droplet phase, 136–146  
  *see also* Bicontinuous phase reactions;  
    Droplet phase reactions
- Microphase ordering, block copolymers, 440
- Microporous carbons  
  agreement between experimental and simulated, 212–213  
  carbon–carbon radial distribution functions (RDFs), 211  
  carbon plate and microstructure from reverse Monte Carlo (RMC), 210  
  nitrogen adsorption isotherms, 212–213  
  experimental RDFs, 209  
  geometric pore size distribution, 212  
  morphological models by RMC, 209–211  
  RMC procedure, 210–211
- Mixing  
  hierarchical evolution protocol, 108–109  
  Monte Carlo simulation results, 107
- Mode coupling theory (MCT), dynamics and relaxation processes, 51–52
- Modified Horvath–Kawazoe method  
  illustration, 236  
  semiempirical adsorption model, 235
- Modified Kelvin equation (MK)  
  relationship between pore filling pressure and pore width, 226  
  *see also* Kelvin equation
- Molecular disorder  
  bond-orientational order parameter, 46–47  
  crystal-independent measures, 48  
  degree of translational order, 47–48  
  jammed hard-sphere structures, 50  
  Lubachevsky–Stillinger protocol, 49  
  non-equilibrium preparation method, 49  
  ordering phase diagram, 48–50  
  quantifying, 45–50  
  random system, 46  
  specific bond-orientational order parameter, 47  
  structural order parameter, 46
- Molecular dynamics (MD)  
  Car–Parrinello, 368–370  
  Car–Parrinello, classical, 357–358  
  classical mechanics, 158  
  collision cascade, 157  
  heating by energetic ion impact, 159–160  
  hierarchical approach for radical-surface interactions, 255–257  
  hierarchical scheme for atomistic simulations, 159  
  ion-assisted etching mechanisms, 164–172  
  ion-surface interactions, 157–158  
  plasma processing, 149–152  
  quantum mechanics and  
    Born–Oppenheimer approximation, 158–159  
  radical-surface interactions, 254–255  
  side view of copper layer with  $\text{Cu}^+$  bombardment, 157  
  simulating long times between ion impacts, 160–161  
  simulation procedure, 156–161  
  size of cell, crucial factor, 160  
  sputtering, 157

Molecular dynamics (MD) (*continued*)

*see also* Ion-assisted etching; Plasma processing

## Molecular evolution

experimental applications of technology, 116

*see also* Protein molecular evolution

## Molecular statics (MS)

hierarchical approach for radical-surface interactions, 255–257

radical-surface interactions, 254

## Monte Carlo methods

alkane adsorption and diffusion, 408

analogy with combinatorial chemistry, 85

benefit of swapping, 2

complex fluids, 2

conventional, 2

effectiveness in materials discovery, 95–96

first molecular simulations, 315

grid method, 94–95

hierarchical approach for radical-surface interactions, 255–257

library design and redesign, 88

low-discrepancy sequence (LDS) method, 94–95

methods requiring energy of assembly of molecules, 315

parallel tempering as extension of, 89

protocols, 94–95

radical-surface interactions, 254

searching variable space by, 87–89

specifying thermodynamic state of replica, 2–3

*see also* Grand canonical Monte Carlo (GCMC); Hyperparallel tempering Monto Carlo; Kinetic Monte Carlo (KMC); Reverse Monte Carlo (RMC)

## Mordenite

hydrogen bonded to bridging O atom, 404

self-diffusivity diffusion constant, 412

structure, 403, 404

*see also* Zeolite catalysis theory

## Morphology

ABC triblock copolymers, 456–459

block copolymers, 441

## Multipool swapping

hierarchical evolution protocol, 109

Monte Carlo simulation results, 107

schematic diagram, 106

## N

Nanostructure formation, *see* Surfactant solutions

## Nitrogen

adsorption isotherm on mesocarbon

microbead (MCMB), 212–213

adsorption isotherms and pore size distribution, 230

## NK model

energy functions, 103

interaction between secondary structures, 104

molecular evolution protocols, 102–104

random energy model, 102

## O

Olefin polymerization, chain termination and branching, 380

Order-disorder transitions, *see* Block copolymers

Order-order transitions, *see* Block copolymers

## P

## Parallel tempering

basic idea, 2

schematic of swapping, 108

*see also* Hyperparallel tempering Monto Carlo

Parasites, evading chemical control, 116

Péclet number, effect in droplet phase reactions, 144–145

Penicillin-binding proteins, 115

## Percolation theory

adsorption and percolation, 223

grand canonical Monte Carlo (GCMC), 222

Percus–Yevick closure, templated structures, 217

Perturbation expansion, droplet phase reactions, 139–141

Perturbation methods, computing interaction energies, 318

Pest organisms, evading chemical control, 116

- Phase behavior, predictive methods, 341
- Phase diagrams
- amphiphiles not forming micelles, 304
  - asymmetric polymer blends, 15
  - block and random copolymers, 18
  - Car–Parrinello simulations, 386–389
  - Lennard–Jones fluid, 7
  - liquid–liquid, of semiflexible/flexible polymer mix, 17
  - long polymer chains, 13
  - semiflexible and flexible polymer, 16
  - soft-sphere plus mean-field model, 70
  - two-parameter ordering, 49
- Phase separation, *see* Surfactant solutions
- Plane-wave basis sets
- Car–Parrinello, 363, 367
  - Hellman–Feynman theorem, 368
- Plasma-enhanced chemical vapor deposition (PECVD)
- amorphous films on substrates by MD, 262–263
  - growth of thin films, 252
  - hierarchical approach to plasma-surface interactions, 255–257
  - methods of surface preparation, 260–263
  - plasma-surface interactions, 252–253
  - procedure for a-Si:H/c-Si film/substrate systems, 261
  - silane and hydrogen containing discharges, 253
  - surface characterization and reaction analysis, 263–264
  - see also* Radical-surface interactions; Surface chemical reactivity, SiH<sub>x</sub> radicals
- Plasma processing
- crystalline silicon surface exposed to chlorine plasma, 154
  - examples of etch profiles, 153
  - ion-surface interactions, 155–156
  - length scales, 152–153
  - molecular dynamics (MD) simulations, 149–150
  - nature of plasma-surface interactions, 153–155
  - processing wafers, 151–152
  - schematic of typical plasma reactor, 151
  - semiconductors, 150–152
  - simulation procedure, 156–161
  - typical conditions of nonequilibrium plasmas, 151
  - see also* Ion-assisted etching; Molecular dynamics (MD)
- Plasma-surface interactions
- nature, 153–155
  - steady-state depth composition profile, 154
- Platinum, slab model, 366
- Polarizable continuum model (PCM), 328
- Polarizable potentials, multibody interactions, 322–323
- Polymerization of olefins, 380
- Polymers
- block and random copolymers, 17–18
  - configuration snapshot of semiflexible systems, 16
  - hyperparallel tempering Monte Carlo (HPTMC), 11–15
  - liquid–liquid phase diagram for semiflexible/flexible mix, 17
  - phase diagram for semiflexible and flexible polymers, 16
  - semiflexible, and blends with flexible, 15–17
- Pore size distribution (PSD)
- activated carbon to methane adsorption, 220–221
  - density functional theory (DFT) results, 231, 232
  - Dubinin–Stoeckli (DS), Horvath–Kawazoe (HK) and DFT methods, 238
  - grand canonical Monte Carlo (GCMC) for CH<sub>4</sub>, CF<sub>4</sub>, and SF<sub>6</sub> isotherms, 223
  - model silica glasses, 240, 241
  - testing robustness for DFT, 229–230
  - see also* Geometric pore size distribution
- Porous glasses
- generation of model, 207
  - geometric pore size distribution, 207–208
  - interpreting adsorption isotherms, 208–209
  - quench molecular dynamics methods, 206–209
  - reverse Monte Carlo technique, 209
- Porous materials
- classical thermodynamic models, 205
  - disordered model microstructure, 204
  - modeling pore structure, 204

- Porous materials (*continued*)  
   molecular simulation calculations, 204  
   pore structure characterization, 203–204  
   semiempirical models, 205  
   simple geometric model, 204  
   statistical thermodynamic theories, 205  
   templated, 216–218  
   thermodynamic models, 204–205  
   *see also* Disordered structure models;  
     Simple geometric pore structure  
     models  
 Propene, geometries during protonation,  
   416, 417  
 Propylene, changes in energy of adsorption  
   of protonated, 425  
 Protein molecular evolution  
   *ab initio* evolution of protein with specific  
     function, 99  
   amino acid substitution, 105  
   background on experimental, 100–102  
   base substitutions, 100  
   challenge of swapping protocols, 110  
   connection between codon usage and  
     DNA shuffling, 113  
   current state-of-the-art techniques,  
     101–102  
   description, 98–99  
   discovery of DNA shuffling, 100–101  
   DNA base mutation, 111, 112, 113  
   DNA base substitution, 115  
   DNA shuffling, 105, 107, 116  
   energy functions of NK form, 103  
   energy of interaction between secondary  
     structures, 104  
   evasion of immune system, 117  
   evolution of multiprotein complex, 99  
   evolution of new life forms, 99  
   evolved resistance, 117  
   examples of exploiting codon  
     potentials, 113  
   examples of swapping by nature, 114–115  
   experimental conditions and constraints,  
     104–105  
   generalized NK model, 102–104  
   genes and operons evolved by DNA  
     shuffling, 101  
   hierarchical decomposition of protein  
     space, 97–98  
   hierarchical decomposition of sequence  
     space, 99  
   hierarchical evolution protocols,  
     105–109  
   mixing, 108–109  
   Monte Carlo results of evolution  
     protocols, 107  
   multipool swapping, 109  
   mutation tendencies for DNA, 112  
   natural analogs of protocols, 113–115  
   parallel tempering, 108–109  
   pests and parasites evading chemical  
     control, 116  
   possible experimental implementations,  
     109–111  
   random or directed mutagenesis, 100  
   searching space, 97  
   simulated molecular evolution  
     protocols, 106  
   single-pool swapping, 107  
   specific energy function as selection  
     criterion, 103  
 Proximity effect, group contribution  
   models, 336  
 Pseudopotential method, Car–Parrinello,  
   367–368
- Q**
- Quantum chemical calculations  
   *ab initio* molecular simulations, 316  
   activity coefficients, 341  
   application of group contribution methods  
     to large compounds, 341  
   application to chemical engineering,  
     346–347  
   availability of computational  
     packages, 346  
   charging free energy contributions,  
     337–338  
   charging free energy from continuum  
     solvation models, 327–328  
   chosen pair potential and polarization  
     term, 323  
   correcting for structure and proximity  
     effects, 336  
   dilute activity coefficients, 331  
   energies by Hartree–Fock with  
     6-311++G(3d,2p) basis set, 344  
   GCS model for octanol–water partition  
     coefficient (GCSKOW), 332

- GCSKOW model with multipole correction method, 337, 338–339
- GCSKOW model with simple group contribution method, 337
- Gibbs ensemble Monte Carlo (GEMC) for predicting phase behavior, 319–321
- group contribution solvation (GCS) model, 330–332
- group scale factor for functional groups, 329
- hydrogen bonding, 346
- infinite dilute activity coefficient of ethanol in heptane, 333
- infinite dilute activity coefficients, 326–327
- influence of proximity effects, 339–341
- interaction model including polarization, 324
- isotropic and anisotropic polarizability, 323–324
- MC computer simulation methods, 315
- MC simulation of water, 316–317
- molecular simulations, 315
- multibody interactions, 321–322
- pair interaction energies of acetonitrile and methanol, 319
- perturbation methods for computing interaction energies, 318
- phase behavior, 321
- phase equilibria predictions, 341–346
- polarizable continuum model (PCM), 328
- potential energy surface between two molecules, 317
- procedure to determine interaction energies, 342
- scale factor in charging free energy calculations, 330
- scale factors and temperature dependence, 334–335
- sequential use of *ab initio* and molecular simulations, 316
- shortcomings of group contribution, 335–336
- simulations (GEMC) of vapor-liquid equilibria, 321
- small clusters of strongly interacting molecules, 341–342
- solvation models for thermodynamic properties, 325–326
- solvation process in two steps, 326
- supermolecular method, 317–318
- temperature effects by continuum solvation, 332–334
- thermodynamic property predictions, 335–341
- two-body potentials and macroscopic property predictions, 318–319
- UNQUAC and Wilson activity coefficient models, 342, 344–345
- using polarizable potentials, 322–323
- vapor-liquid coexistence diagrams, 320
- vapor pressure, 320
- water models, 324–325
- R**
- Radial distribution functions (RDFs), 209, 211
- Radical-surface interactions
- atomic-scale simulation studying deposition, 255
- atomic-scale theoretical studies, 254
- atomistic simulation methods, 254–255
- characterizing surfaces, reactivity, 264–265
- clusters containing Si atoms in plasmas, 288–289
- density functional theory (DFT), 257–258
- disilane formation, 278–279
- dissociative adsorption mechanism, 274–275
- dominant deposition precursor, 284–286
- Eley–Rideal H abstraction, 277
- empirical description of interatomic, 258–260
- film growth mechanisms by PECVD, 273
- film surface composition, 283–284
- hierarchical approach, 255–257
- hydrogen abstraction by SiH<sub>3</sub> radical, 276–278
- impact of Si<sub>6</sub>H<sub>13</sub> clusters on film defects, 290
- kinetic Monte Carlo (KMC) methods, 255
- Langmuir–Hinselwood H removal, 277–278
- mechanism of amorphous silicon film growth, 280–281
- mechanism of attachment for SiH radical, 286–287
- mechanism of SiH penetration, 287–288

- Radical-surface interactions (*continued*)  
 nanosecond-time scale MD of film growth, 273  
 reactions of  $\text{SiH}_2$  radical with H:Si surface, 268–269  
 role of chemically reactive minority species, 286–290  
 $\text{SiH}_2$  radical, possible adsorption sites, 268  
 $\text{SiH}_3$  insertion reactions and dissociative adsorption events, 275  
 $\text{SiH}_3$  radical attaching to pristine Si surface, 269  
 $\text{SiH}_x$  radicals with amorphous silicon films, 270–272  
 $\text{SiH}_x$  radicals with crystalline silicon surfaces, 266–269  
 $\text{Si}_6\text{H}_{13}$  cluster with a-Si:H surface, 289  
 structure of crystalline and amorphous silicon, 265–266  
 surface characterization and reaction analysis, 263–264  
 surface chemical reactions during film growth, 274–280  
 surface evolution and characterization, 281–283  
 surface preparation, 260–263  
*see also* Surface chemical reactivity,  $\text{SiH}_x$  radicals
- Random copolymers  
 hyperparallel tempering Monte Carlo (HPTMC), 17–18  
 phase diagram, 18
- Random phase volume model  
 functional form, 94  
 materials discovery, 92–94  
 number of phase points, 93
- Reactivity of surfaces, *see* Surface chemical reactivity,  $\text{SiH}_x$  radicals
- Reactor design and kinetics, 123–124
- Relaxation, viscous liquids, 28
- Renormalization group theory  
 bicontinuous phase reactions, 132–134  
 flow equations, 133–134  
 framework for identifying system behavior, 132  
 implementing analysis, 133  
 phase-space flow diagram, 134
- Replica Ornstein–Zernicke (ROZ) integral equations, 217–218
- Reverse Monte Carlo (RMC)  
 agreement with physical properties, 212–213  
 morphological model for microporous carbon, 209–211  
 procedure, 210–211
- Rotational diffusion, 31–32
- S**
- Sampling, materials discovery, 91–92
- Sastry density, description, 68
- Sastry point  
 superheated liquid and, 68  
 theoretical predictions, 69
- Self-assembled system, applications, 124
- Semiconductors, plasma processing, 150–152
- Semiflexible polymers  
 configuration snapshot, 16  
 liquid-liquid phase diagram, 17  
 phase diagram, 16  
*see also* Polymers
- Sequence space, hierarchical decomposition, 99
- Short time regime, droplet phase, 143–144
- $\text{SiH}_x$  radicals, *see* Surface chemical reactivity,  $\text{SiH}_x$  radicals
- Silica ( $\text{SiO}_2$ )  
 strong glass former, 56  
 topographic illustration, 57  
*see also* Supercooled liquids and glasses
- Silicon  
 atomic configuration after  $\text{CF}_3^+$  fluence, 185  
 average yield of major sputtered species, 188  
 C and F content as function of  $\text{CF}_3^+$  fluence, 183  
 C, F, and Si depth profiles for  $\text{CF}^+$ , 196, 197  
 $\text{CF}^+$  and  $\text{CF}_2^+$  deposition, 190–198  
 $\text{CF}_3^+$  etching, 182–186  
 deposition yield of F and C for  $\text{CF}^+$  and  $\text{CF}_2^+$ , 195  
 depth profiles for Si, F, and C after  $\text{CF}_3^+$ , 186  
 etch yield and fluorocarbonsilyl layer thickness, 187  
 interfacial  $\text{Si}_x\text{C}_y\text{F}_z$  layer thickness for  $\text{CF}^+$  and  $\text{CF}_2^+$ , 198



- product angular distributions as function of angle, 192
- product distributions in  $\text{CF}_3^+$  etching, 186–190
- product kinetic energy distributions, 191
- sticking coefficients of F and C for  $\text{CF}^+$  and  $\text{CF}_2^+$ , 195
- surface content, 184
- uptake of F and C as function of  $\text{CF}^+$  fluence, 193
- uptake of F and C as function of  $\text{CF}_2^+$  fluence, 194
- yields of major products, 189
- see also* Ion-assisted etching
- Silicon film growth
- clusters with Si atoms in plasmas, 288–289
  - disilane formation, 278–279
  - dissociative adsorption, 274–275
  - Eley–Rideal H abstraction, 277
  - evolution of surface roughness in MD simulation, 282–283
  - film surface composition, 283–284
  - hydrogen abstraction by  $\text{SiH}_3$  radicals, 276–278
  - impact of  $\text{Si}_6\text{H}_{13}$  clusters, 290
  - Langmuir–Hinselwood H removal, 277–278
  - MD simulations of  $\text{Si}_6\text{H}_{13}$  clusters, 289
  - mechanism of amorphous, 280–281
  - mechanism of SiH radical attachment, 286–287
  - mobility of  $\text{SiH}_3$  radical on amorphous growth surface, 279–280
  - plasma-surface interactions, 273
  - role of chemically reactive minority species, 286–290
  - role of dominant deposition precursor,  $\text{SiH}_3$ , 284–286
  - $\text{SiH}_3$  insertion reactions, 275
  - SiH penetration mechanism, 287–288
  - surface chemical reactions during, 274–280
  - surface evolution and characterization, 281–283
  - see also* Radical-surface interactions; Surface chemical reactivity,  $\text{SiH}_x$  radicals
- Simple geometric pore structure models
- adsorption isotherms and pore size distribution (PSD), 230
  - carbon micropore distributions, 237, 239
  - classical adsorption models, 239–244
  - comparing pore filling correlations for nitrogen in MCM-41, 241
  - adsorption and percolation, 223
  - density functional theory (DFT) adsorption models, 225–231
  - DFT model isotherms for various gases, 228–229
  - Dubinin adsorption models, 236–239
  - Dubinin and Astakhov (DA) equation, 236–237
  - Dubinin–Radushkevitch (DR) equation, 236
  - Dubinin–Stoeckli (DS) method, 237
  - excess adsorption by adsorption integral equation, 218–219
  - Fredholm integral equations, 219–220
  - Gibbs ensemble Monte Carlo (GEMC), 222, 224–225
  - grand canonical Monte Carlo (GCMC), 222
  - Horvath–Kawazoe (HK) method, 231, 233–234
  - interchange steps, 224
  - Kelvin equation, 239
  - Lennard–Jones models, 222
  - mean pore sizes of adsorbents, 240
  - model nitrogen adsorption isotherms, 243
  - modified Horvath–Kawazoe models, 235
  - modified Kelvin (MK) equation, 239–240
  - molecular displacements, 224
  - molecular simulation adsorption models, 222–225
  - percolation theory with GCMC, 222
  - pore filling model, 242
  - pore filling pressure and pore width, 226
  - pore filling pressure correlations, 243
  - pore size distribution (PSD) for methane adsorption, 220–221
  - PSD using GCMC for  $\text{CH}_4$ ,  $\text{CF}_4$ , and  $\text{SF}_6$ , 223
  - PSDs by various methods, 238
  - PSDs for model porous silica glasses, 241
  - schematic of GEMC method, 224
  - semiempirical adsorption models, 231, 233–239
  - shortcomings spurring efforts to modify Kelvin, 240, 242
  - volume exchange steps, 225

- Simple geometric pore structure
  - models (*continued*)
  - weighting function for DFT, 227–228
- Single-pool swapping, 107
- Slab model, platinum(111) surface, 366
- Solid acid catalysts, *see* Zeolite catalysis theory
- Solids, melting points by Car–Parrinello, 387, 389
- Solvation process, 325, 326
- Space of variables, materials discovery, 84–85
- Sputtering
  - description, 157
  - see also* Ion-assisted etching
- Steepest descent mapping, 35
- Stokes–Einstein equation, 31, 58
- Structural order parameters
  - bond-orientational, 46–47
  - crystal-independent measures, 48
  - Lubachevsky–Stillinger protocol, 49
  - measuring degree of translational order, 47–48
  - ordering phase diagram, 48–50
  - specific bond-orientational, 47
  - terminology, 46
  - Voronoi–Delaunay tessellation, 46–47
- Superacid catalysis, 403–405
- Supercells
  - structural and bonding properties of water, 378–380
  - water, 365
- Supercooled liquids and glasses
  - bond-orientational order parameters, 46–47
  - characterization of voids, 40
  - chemical engineering science, 22–23
  - crystal-independent measures, 48
  - density dependence of average pressure, 42–43
  - energy landscape, 33–39
  - entropy for liquids and stable crystals, 29–30
  - expanding system to lower densities, 43–44
  - geometric algorithm, 40–41
  - geometric analysis of void space, 43
  - glass transition temperature, 24
  - glass transition transformation range, 25–26
  - good glass-forming materials, 44–45
  - hydrogen bonds in aqueous systems, 45
  - isobaric relationship between volume and temperature, 25
  - Kauzmann paradox, 30
  - Kauzmann temperature, 29–30
  - kinetics and thermodynamics, 30–31
  - Kohlrausch–Williams–Watts (KWW) function, 28–29
  - landscape dynamics and relaxation phenomena, 50–60
  - morphology of Lennard–Jones inherent structures, 42
  - nonexponential relaxation, 28
  - ordering phase diagram, 48–50
  - parameter for degree of translational order, 47–48
  - quantifying molecular disorder, 45–50
  - questions from Lennard–Jones fluid, 44
  - relationship between enthalpy and temperature, 26
  - relationship between volume and temperature, 24, 25
  - rotational diffusion and viscosity, 31–32
  - scalar measures for translational order, 47
  - shape of pressure versus density curve, 44
  - shifted-force Lennard–Jones system, 42
  - specific bond-orientational order parameter, 47
  - statistical geometry and structure, 39–50
  - strong-fragile classification of liquids, 28
  - structural order parameter, 46
  - substances known to form glasses, 24
  - T<sub>g</sub> depending on cooling rate, 24–25
  - thermodynamics, 60–70
  - tools and protocols, 32
  - translational and rotational motion, 31–32
  - truly random system, 46
  - viscosity by Vogel–Tammann–Fulcher (VTF), 26–27
  - viscosity of glass-forming liquids, 27, 28
  - viscosity of supercooled liquids, 27
  - vitrification, 23–29
  - void geometry and energy landscape, 40–45
  - volume and surface area of packings, 40, 41
  - see also* Energy landscape; Landscape dynamics and relaxation; Thermodynamics of supercooled liquids and glasses
- Superionic conductor, 376–377

- Supermolecular method, 317–318  
 Surface chemical reactivity,  $\text{SiH}_x$  radicals  
   characterizing surface reactivity, 264–265  
   crystalline and amorphous silicon surfaces, 265–266  
   possible adsorption sites of  $\text{SiH}_2$  radical, 268  
   projections of  $\text{SiH}_3$  center-of-mass trajectories, 271  
   radical impingement on a-Si:H surfaces, 272  
   reactions of  $\text{SiH}_2$  radical with H:Si surface, 268–269  
   reactivity maps, 266, 270  
    $\text{SiH}$  radical, 272  
    $\text{SiH}_3$  radical attaching to pristine Si surface, 269  
    $\text{SiH}_x$  radicals and silicon surfaces, 267–269  
    $\text{SiH}_x$  radicals with amorphous silicon films, 270–272  
    $\text{SiH}_x$  radicals with crystalline silicon surfaces, 266–269  
 Surface roughness  
   evolution during film growth, 282–283  
   ion-surface scattering, 174–178  
 Surfactant solutions  
   aggregation to large, stable clusters, 302  
   assembly into nanostructures, 298  
   behavior of (I1) and (I2) interaction sets, 304–305  
   cmc vs. inverse temperature, 307  
   critical micelle concentrations (cmc), 306–307  
   dimensionless ratio, 298  
   exchange energy parameters, 300–302  
   homopolymer chains, 302–304  
   macroscopic phase separation to micellar aggregates, 308–309  
   methodological issues, 301–302  
   micellization processes, 299–300  
   models and methods, 300–301  
   mole fraction in dilute phase vs. inverse temperature, 303  
   normalized cluster probability distributions, 308  
   phase behavior of homopolymers, 303  
   phase diagrams of amphiphiles, 304  
   reduced osmotic pressure vs. volume fraction surfactant, 306  
   role of different interaction sets, 304–308  
   self-assembly of amphiphilic molecules, 299  
   simulation details, 300–302  
   snapshot of  $\text{H}_1\text{T}_4(\text{I2})$ , 305  
   snapshot of  $\text{H}_2\text{T}_4(\text{I1})$ , 305  
   snapshot of  $\text{H}_2\text{T}_4(\text{I2})$ , 306  
   surfactant architectures and interaction energy sets, 300  
   time scales for micellar aggregate formation, 299  
 Swapping  
   acceptance rate, 19  
   benefit, 2  
   changing variables, 87–88  
   DNA, 115  
   examples by nature, 114–115  
   hyperparallel tempering Monte Carlo, 4–5  
   ITCHY and SCRATCHY techniques, 110–111  
   Monte Carlo simulation results, 107  
   multipool, 109  
   schematic diagram, 106  
   single-pool, 107  
   technical challenge by protocols, 110  
 Symmetry-adapted perturbation theory (SAPT), 318
- T**
- Temperature effects, 332–334  
 Templated porous materials  
   illustration of porous matrix by templating, 216  
   Monte Carlo simulations, 217–218  
   Percus–Yevick closure, 217  
   replica Ornstein–Zernicke (ROZ) integral equations, 217–218  
 Templating, porous matrix, 216–218  
 Temporal regimes, droplet phase, 141–143  
 o-Terphenyl (OTP)  
   fragile glass former, 56–57  
   topographic illustration, 57  
 Thermodynamics of supercooled liquids and glasses  
   basin enumeration function, 60–62  
   disordered materials, 69–70  
   macroscopic system with Gaussian landscape, 66–67

## Thermodynamics of supercooled liquids and

glasses (*continued*)

- Gaussian landscape parameters, 66
- Helmholtz energy, 61
- ideal glass, 64
- isobaric basin enumeration functions, 62, 63, 65
- isobaric excitation profiles, 64
- isobaric excitation profiles and isochoric profiles, 65–66
- low-temperature termination of
  - Kauzmann curve, 68–69
- maximum tensile strength of amorphous material at Sastry density, 68
- partition function as one-dimensional integral over basin depth, 61
- phase diagram of soft-sphere plus mean-field model, 70
- relations for systems with Gaussian landscapes, 66–67
- spinodal curve for superheated liquid and Sastry point, 68
- stretched liquids and glassy state, 67–68
- temperature-dependent basin depth, 62–64
- theoretical predictions of Sastry point, 69
- thermodynamic properties and energy landscape, 61
- Third Law of Thermodynamics, Kauzmann's paradox, 29–30
- Tiling model, supercooling and glass formation, 55
- Transition state energies, hydrocarbon activation, 420–421
- Triblock copolymers
  - complex two-dimensionally ordered ABC phases, 457
  - metastable states from preordered states, 459
  - morphology and metastability, 456–459
  - transition from two- to three-domain structure, 458
- see also* Diblock copolymers

## U

- UNIQUAC model, activity coefficients, 342, 344–345

## V

- Vapor-liquid coexistence diagram
  - acetonitrile, 320
  - methanol, 320
- Vapor-liquid equilibrium diagram, acetone + water, 345
- Vapor pressure
  - acetonitrile, 320
  - methanol, 320
- Variables, space of, materials discovery, 84–85
- Viscosity
  - glass-forming liquids, 26–27
  - glass-forming liquids in Arrhenius fashion, 27, 28
  - rotational diffusion and, 31–32
  - Vogel–Tammann–Fulcher (VTF) equation, 26–27
- Vitrification, *see* Supercooled liquids and glasses
- Vogel–Tammann–Fulcher (VTF) equation, viscosity, 26–27
- Void space, geometry of pore structure, 40
- Voronoi–Delaunay tessellation, 41

## W

- Wafers processing, 151–152
- Water
  - cubic supercell, 365
  - models based on effective potential, 324–325
  - Monte Carlo from *ab initio* calculations, 316–317
  - structural and bonding properties, supercells, 378–380
- Water, deuterated, infrared spectrum, 380, 381
- Wilson model, activity coefficients, 342, 344–345

## X

- Xerogels
  - grand canonical Monte Carlo (GCMC), 213–216
  - perturbations, 213

## INDEX

485

simulated adsorption and desorption  
isotherms, 215  
visualizations from GCMC, 215

**Z**

Zeolite catalysis theory  
activation energies for isomerization and  
protonation, 431, 432  
bonding within zeolite framework,  
405–406  
Born repulsion expression, 406–407  
catalytic cycle of zeolite-catalyzed  
reaction, 400  
density functional theory (DFT), 402–403  
dependence of energy of adsorption for  
linear alkanes, 409–410  
energies of adsorption for hydrocarbons  
and zeolites, 410–411  
entropy expression, 403  
flexibility of zeolite framework, 405  
hexane in zeolite micropore, 406  
hydrocarbon activation by zeolitic protons,  
414–427  
hydrogen attached to bridging oxygen  
atom, 403–404  
hydroisomerization reaction, 427–428

interaction potential, 406  
isomerization reaction of hexene, 428–430  
kinetics, 427–431  
kinetics for hexane isomerization,  
430–431  
methods and theory of alkane adsorption  
and diffusion, 407–408  
Monte Carlo computing thermodynamic  
averages, 408  
overall rate of catalytic reaction, 411  
quantum-chemical computations, 402  
rate of catalytic reaction, 401–403  
simulation of alkane adsorption and  
diffusion, 407–414  
simulation of alkane adsorption isotherms,  
409–411  
simulation of diffusion, 412–414  
structure of mordenite, 403, 404  
superacid catalysis, 403–405  
transition rate theory, 402  
zeolites as solid acid catalysts, 403–407  
*see also* Hydrocarbon activation  
Zeolites  
unit cell of chabazite, 364  
*see also* Chabazite  
Ziegler–Natta catalysts  
Corradini site, 383, 384  
polymerization reactions, 382–385