$\mathbf{A}$	BCNU, 210
	Bethanechol, 210
Acrylamides, 118	Bioactive agents, 2
Actin cytoskeleton, 17–18	Bioactive factor delivery, 159-160
Adenosine deaminase deficiency, 131	Biocompatibility, 1–5
Adhesive peptide sequences within ECM	concept of, 3
proteins, 12	host response central to, 5
Adhesive peptides, 37	Biodegradable conduits, comparative
Adsorbed proteins, conformation of, 22	studies, 59
Aliphatic polyanhydrides, 177	Biodegradable materials, 50
copolymers, 182	Biodegradable polymer scaffolds, 61
Allografts, 47	Bioerodible polymers, 169
Alpha, 1-glycoprotein (alpha 1-GP) 65	functional groups found in, 171
Alzheimer's disease, 64, 67, 210	Biohybrid nerve guide, 56
Amine-modified surfaces, 29	Biological response, 5
Amyotrophic lateral sclerosis (ALS), 67	Biologically active materials vs. drugs, 4
Angiogenesis inhibitors, 210	Biomaterials
Angiogenic biomaterials, 3	as agonists of biological response, 3
Anhydride interchange mechanism, 175	inert, 1–2
Anti-adhesive matrix molecules, 17–18	overview, 1–6
Anti-PECAM, 120	synthesis, 15
Apoptosis, 23–24	three-dimensional polymeric networks as
adhesion-dependent control, 23	76–105
Appropriate host response, 3	Blood compatibility, 3
Arg-Gly-Asp peptides (RGD), 11, 37-38,	Bone and GAMs
63, 104	larger animals, 160-161
Asialoglycoprotein, 40	small animals, 160
Astrocytes, 48	summary, 161
Atomic force microscopy (AFM), 195	Bone morphogenetic proteins (BMPs), 159
ATR-FTIR spectroscopy, 111	Bone regeneration, 158–159
Autografts, 47	Bone tissue engineering, 159–161
Axonal growth	Bovine serum albumin (BSA), 65
control of, 55	Bulk erosion, 172
electrically mediated guidance, 60	
longitudinally directed, 61	
Azo-compounds, 121	C
	<sup>13</sup> C-NMR studies, 96
В	Carbohydrates, 13, 15
	Carcinomas, 210
Baby hamster fibroblast cell line (BHK),	Cationic lipids, 136–138
67–68	Cationic polymers, 138–141
Basic fibroblast growth factor (b-FGF), 65	CD36, 17
BCECF-am, 122	CD44, 16–17
	- ,

CD44–ligand interaction, 16	
	Collagen nerve guide, 56
CD44 receptors, 16	Collagen-NGF mixtures, 51
Cell adhesion, 7–8, 11, 21–25, 28–29,	Combinatorial chemistry, 75
32–33, 40	Complementary DNA, 122
Cell-adhesion ligands, 36–41	Complexation in polymers, 92-98
Cell attachment to quaternary amine	effect of PEG chain length, 92-96
surfaces, 29	interpolymer, 92–96
Cell contractility, 25	overview, 92
Cell-derived matrix, 21	Complexation of poly(ethylene glycols), 92
Cell differentiation, 21–23	Compound muscle action potential
Cell encapsulation see Cell encapsulation	(CMAP), 59
Cell integrity, 28	Compression molding, 201
Cell-material interactions, 7-46	Concanavalin-A (Con-A), 118
controlled, 33–41	Conduit materials, comparative studies,
nonspecific, 25-33	57–60
Cell migration, 24–25	Configurational biomimetic imprinted
Cell motility, 24–25	networks, 78
Cell placement and growth, 55-56	Controlled cell-material interactions, 33-41
Cell-recognition processes, 14	Controlled delivery, polymers for, 141
Cell–substrate contacts, 22	Controlled DNA delivery, polymeric
Cell-substrate interactions, 8, 18-19, 24	scaffolds for, 157–162
Cell-surface glycosaminoglycan, 29	Controlled DNA release, polymer based
Cell surface receptors	particles for, 141–157
activation, 13	Controlled release, polyanhydride carriers
and their ligands, 8–18	for, 209–213
recognition, 34	Controlled-release device, 86
Cellular differentiation, 32	Controlled-release polymers, 64-66
Central nervous systems, polymers for	Corticospinal tract (CST), 60
regeneration, 60–64	Covalently immobilized adhesive peptide
Cetyltrimethylammonium bromide (CTAB),	sequences, 36–37
153	
155	*
	Cystic fibrosis, 131
Charge transfer complexes, 92	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19
Charge transfer complexes, 92 Charged surfaces, 28	Cystic fibrosis, 131
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19
Charge transfer complexes, 92 Charged surfaces, 28	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155 Chitosan–DNA nanospheres, 155	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139 Dichloromethane (DCM), 191–192, 200
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155 Chitosan–DNA nanospheres, 155 Chloroquine, 155	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155 Chitosan–DNA nanospheres, 155 Chloroquine, 155 Chondroitin sulfate, 29 Chondroitinase ABC, 29	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30  D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139 Dichloromethane (DCM), 191–192, 200 Differential scanning calorimetry (DSC), 194
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155 Chitosan–DNA nanospheres, 155 Chloroquine, 155 Chondroitin sulfate, 29 Chondroitinase ABC, 29 Ciliary neurotrophic factor (CNTF), 67–68	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30   D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139 Dichloromethane (DCM), 191–192, 200 Differential scanning calorimetry (DSC), 194 Diffusion, fundamentals of, 84–87
Charge transfer complexes, 92 Charged surfaces, 28 cell response to, 29 Chemical cues, 50–51 Chemical potential change due to elastic retractive forces, 79 Chemotherapeutic agents delivery using hydrogels, 121 Chinese hamster ovary (CHO) cells, 28 Chitosan, 112, 142, 153, 155 Chitosan–DNA microspheres, 156 Chitosan–DNA nanoparticles, 155 Chitosan–DNA nanospheres, 155 Chloroquine, 155 Chondroitin sulfate, 29 Chondroitinase ABC, 29	Cystic fibrosis, 131 Cytoplasmic alkalinization, 19 Cytoskeletal organization, 30   D  DDAB, 151–153 Degradation, use of term, 170 Degradation half-lives, 170 Dehydrative coupling, 175–176 Dehydrochlorination, 173, 175 Dendrimers, 121 Dextran, 139 Dichloromethane (DCM), 191–192, 200 Differential scanning calorimetry (DSC), 194 Diffusion, fundamentals of, 84–87 Diffusion coefficients, 85–86

Directed growth methods, 52	Encapsulation, 66–68
DMAEMA, 114	advantages over injection, 143
DMSO, 192	plasmid DNA, 143-150, 152, 154
DNA	Endothelial cells, 21, 33, 104
degradation, 147	Entubulization, 48
delivery, 136–137, 141	degradable polymers for, 52-57
denaturation, 146	nondegradable polymers for, 50–52
release from scaffolds, 161–162	Equilibrium swelling theory, 79–81
release in vaccination applications,	Erosion
153–157	characterization, 170
sequences, 122	use of term, 170
synthesis, 21	Ethyl cellulose, 113
vaccinations, 142, 162	Eucaryotic cells, 13
see also Plasmid DNA	Eudragit 100 gels, 119
DNA/DDAB complexes, 151	Extracellular matrix see ECM
DNA-gelatin microspheres, 154	Extracellular matrix see ECW
DOTAP, 153	F
Double-emulsion solvent-evaporation	Г
technique, 146	Enhancian methods for menous conduits
Drug delivery, 2	Fabrication methods for porous conduits,
surface-erodible biomaterials for,	56–57
169–218	Factor XII, 3
Drug release	FAD, 178
chemically-controlled, 84	FDA, 2
control mechanisms, 170	Fibrillar adhesions, 20
diffusion-controlled, 84	Fibroblast growth factor (FGF), 51
swelling-controlled, 84	Fibroblasts, 20–21, 33, 50
Drugs vs. biologically active materials, 4	Fibronectin, 16, 20, 22, 27, 36, 50, 55, 64
	Fickian diffusion theory, 85
F	Fick's law of diffusion, 84
${f E}$	Flory–Rehner theory, 79–80
	Fluorinated polyanhydrides, 189
E–selectin, 15, 40	Focal adhesions, 19, 25
ECM, 9–10, 17, 104	Foreign body response, 101–102
coatings, 22	FT-IR spectroscopy, 94
components, 16, 20, 24, 48-49, 104	
environment, 20	
geometry, 22	G
layers, 31	
ligands, 19	Galactose, 15
molecules, 15-16, 22	Gelatin, 142, 153–154
orientation, 31–32	Gene activated matrices (GAMs),
production, 33, 39	157–158
properties, 31	and bone
proteins, 11, 34, 36, 41	larger animals, 160–161
adhesive peptide sequences within, 12	
	small animals, 160
synthesis, 39	small animals, 160 summary, 161
synthesis, 39	summary, 161
synthesis, 39 ECM–integrin interactions, 21	summary, 161 Gene delivery

Gene therapy, 131–168 and protein delivery, 132–133	macroporous, 82 microporous, 82
concept of, 162	network structure, 77–83
ex vivo, 133	neutral, 105–110
in vivo, 133–134	nonporous, 82
nonviral, 136	pH-sensitive, 89–90, 110–112, 114–115
biomaterials for, 136–141	poly(glyceryl methacrylate)
overview, 131–132	p(GMA)-collagen, 63
viral, 135	protein-based, 120
Genetic immunization, 142–143	solute permeability properties, 77
Gibbs free energy, 79	solute transport in, 83–87
Glial cell line-derived neurotrophic factor	swollen temperature, 89
(GDNF), 65	synthetic, 63
Glial growth factor (GGF), 53	temperature-sensitive, 90-91, 112-115
Glioblastoma multiforma, polyanhydride	Hydrogen-bonding complexes, 92
discs for treatment of, 210	Hydrogen bonds, 97-98
Glucose response following infusion of	inter-associating, 97
human insulin, 107	self-associating, 97
Glucose-sensitive systems, 117–119	Hydrophilic polymer networks, 76
Glycocalyx, 14, 29	Hydrophilicity/hydrophobicity, 30, 76, 104
Glycopeptides, 39	Hydrophobic monomers, 100
Glycoproteins, 14	Hydrophobicity/hydrophobicity, 30, 76, 104
Glycosaminoglycann (GAG), 15	2-Hydroxyethyl methacrylate (HEMA), 96,
GPC, 191	99
GRGDY, 37	Hydroxypropyl cellulose (HPC), 113
Growth factor administration, 64	
	I
H	-
Н	ICAM-1, 11
H  H-NMR spectroscopy, 96, 198	ICAM-1, 11 ICAMs, 16
H  H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35
H  H-NMR spectroscopy, 96, 198  Hemocompatibility testing, 5  Hemophilia, 131–132	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17
H  H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2
H  H-NMR spectroscopy, 96, 198  Hemocompatibility testing, 5  Hemophilia, 131–132	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17
H  H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98
H  H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76 applications, 76, 105–122	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12 Integrins, 9–13, 17, 19, 36–39
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76 applications, 76, 105–122 chemotherapeutic agents delivery using, 121 classification, 77	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12 Integrins, 9–13, 17, 19, 36–39 Intracellular domain, 9
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76 applications, 76, 105–122 chemotherapeutic agents delivery using, 121 classification, 77 complexing, 91, 115–117	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12 Integrins, 9–13, 17, 19, 36–39
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76 applications, 76, 105–122 chemotherapeutic agents delivery using, 121 classification, 77 complexing, 91, 115–117 drug delivery applications, 121	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12 Integrins, 9–13, 17, 19, 36–39 Intracellular domain, 9
H  1H-NMR spectroscopy, 96, 198 Hemocompatibility testing, 5 Hemophilia, 131–132 Heparin immobilization, 1 Hepatocytes, 39 Host response central to biocompatibility, 5 Human insulin concentration, 107 Human nerve growth factor gene (hNGF), 67 Huntington's disease, 68 Hyaluronic acid (HA), 16–17 Hydrogel tubes, 64 Hydrogels, 60, 76 applications, 76, 105–122 chemotherapeutic agents delivery using, 121 classification, 77 complexing, 91, 115–117	ICAM-1, 11 ICAMs, 16 Ideal surfaces, 34–35 Immunoglobulins, 9, 16–17 Inert biomaterials, 1–2 Inflammatory response, 102 Infrared spectroscopy, 96–97 and polymer complexation, 97–98 Inside-out signaling, 12–13 Insulin delivery systems, 119–120 Insulin-like growth factor (IGF), 159 Integrin–ligand binding, 18–25 Integrin–ligand interactions, 12 Integrins, 9–13, 17, 19, 36–39 Intracellular domain, 9

## L

L-selectin, 15
Lactose, 15
Laminin, 16, 48, 50, 55, 58, 60
Lectin-carbohydrate interactions, 40
Lectins, 15, 40
Leukocyte adhesion, 28
Leukocyte binding, 40
Lower critical solution temperature (LCST), 91, 112–114

 $\mathbf{M}$ 

Macrophages, 103 Matrix-based systems, solute transport in, Medical devices, 2 Melt polycondensation, 174–175 Membrane proteins, 13 Methacrylic acid (MAA), 96 Methyl methacrylate, 100 Mitogen-activated protein kinases (MAPK), 21 Molecular design, 75 Monocyte activation, 3 Monosaccharide asialoglycoprotein ligands, 40 Monosaccharides, 15 Mucosal delivery, aerosols and systems designed for, 212-213 Multigrooved substrates, 32

### N

N-acetyllactosamine, 40
N-methyl pyrrolidinone (NMP), 98
Naked plasmid DNA (pDNA), 142
limitations, 134–135
NCAM, 16
Neomembranes, 56
Nerve growth factor (NGF), 48, 51, 64–65
silicone tubes prefilled with, 59
Nerve growth factor (NGF)-treated
nitrocellulose implants, 60
Nerve guide conduit, 58
Nerve regeneration, polymeric biomaterials
for, 47–74

Neural glial cell adhesion molecule (NgCAM), 16
Neural stem cells, 61
Neurotrophic factors, 64, 66
Neurotrophin-3 (NT-3), 64
Neutral networks, tissue engineering, 98–105
Nitrocellulose implants, 60
Nonviral gene therapy, 136
biomaterials for, 136–141
Nonviral plasmid DNA, 160
Nucleic acids, 131–132

#### O

Office of Combination Products, 2 Oligosaccharides, 14 Osteoblasts, 33 Osteopontin, 16 Outside-in signaling, 13, 20

#### P

p-nitroaniline (PNA), 191, 195 encapulation, 154 P-selectin, 15, 40-41 PAA, 93, 111, 118, 175, 192-193 PAA/PEG complexes, 93-94 Parkinson's disease, 64, 66 PBA, 118 PCDA, 177 PCPA, 179, 191, 200 PCPAM, 187 PCPH, 174, 179, 191, 193-194, 200 P(CPH-CPS), 186 PCPM, 173, 179, 193, 200 PCPO, 179, 191, 200 PCPP, 174-175, 179, 191-194, 199-200 P(CPP-DDA), 174 P(CPP-SA), 174 PCPS, 186, 200 PCPSM, 187 PCPV, 179, 191, 200 PDDA, 174-175, 177, 191-194, 200, 205 PDEAEMA, 110 P(DEAEMA-co-HEMA), 111 PDLA, 56–58 PDLLA, 62 PDMAEMA, 110, 118

P(DMAEMA-co-HEMA), 111	PLGA, 53, 55, 57, 65–66, 142–157, 161
pDNA see Plasmid DNA	PLGA-CTAB microspheres, 153
PDX, 177	PLGA-CTAB nanospheres, 153
PEG, 34–35, 38, 95, 105, 109–110, 115,	PLL, 139, 147–148
119–120, 139, 152, 156	PLLA, 53, 56–57, 59
PEG 400, 66	PMAA, 93, 110, 115–117, 122
PEG chain length, 92–96	PMAA hydrogels, 95
PEG-dimethacrylates, 110	P(MAA-g-EG), 118–119
PEG-monomethacrylates, 110	PMAA/PEG complexes, 93–94
•	PNA, 198
PEG-PLA, 151–152	
PEI, 139	PNIPAAm, 112–113, 115, 118
PELA (2)	PNVP, 105
PELA, 62	Polylactic acid, 57
PEO, 2, 105, 109–110, 112, 121	Polyacrylamide, 34
Peptide-modified surfaces, 36–37	Polyacrylamide (PAAm), 110
Peripheral nerve damage, 47	Poly(acrylic acid) (PAA), 110
Peripheral nervous system, polymers for	Poly(adipic acid) (PAA), 177
regeneration, 48–60	Poly(alpha-hydroxy acid), 61
PEVAc, 105, 161	Poly(amidoamine) dendrimers, 139
PFA, 177, 192, 194	Poly(amidoamine) (PAMAM)
PFAD, 178, 193	dendrimers, 140
PFG-NMR technique, 95	Polyanhydride carriers for controlled
PGA, 59	release, 209-213
pH effect on polymer mesh size, 116	Polyanhydride discs for treatment of
pH-sensitive hydrogels, 89–90, 110–112	glioblastoma multiforma, 210
Phagocytosis, 102	Poly(anhydride-co-alkylene carbonate)s, 188
Pharmacotectonics concept, 66	Poly(anhydride- <i>co</i> -amide)s, 187–188
PHB, 54, 64	Poly(anhydride-co-ester)s, 183–187
PHEMA, 34, 63–64, 77, 97–101, 104–106,	Poly(anhydride-co-ether)s, 183–187
115, 118–119	Poly(anhydride- <i>co</i> -imide)s, 183
PHEMA- <i>co</i> -AA, 111	Poly(anhydride- <i>co</i> -urethane)s, 189
PHEMA-co-MAA, 111	Polyanhydrides
p(HEMA-co-MMA), 64	additional chemistries, 188–189
PHFB, 189	aliphatic, 193
PHMCSA, 188	amorphous phase behavior, 195–199
Photolithographic methods, 22	aromatic, 179–182, 194
PHPMA, 63, 139	as surface erodible biomaterials, 171
PHSRN, 37	biocompatibility, 199–200
PIPA, 179, 191–192, 194	branched, 188
P(IPA–SA), 174	characterization, 189–200
PLA, 121, 211	chemical changes during erosion, 206
PLA/PLGA, 144	chemical characterization, 189–192
Plasmid DNA, 133, 136, 138, 141, 157–158,	chemistry and synthesis, 172–189
161–162	crystalline morphology, 193–195
encapsulation, 143–150, 152, 154	crystallinity, 196–197
entrapment and release, 157	degradation kinetics, 200-208
interaction with cationic lipids, 137	discretized models, 209
nonviral, 160	drug release kinetics, 201, 207–209
PLC, 54, 59-60	drug release rates, 201-205
PLCA, 189	early synthesis, 172–173

erosion, 201, 207–209	Porous materials, tissue response to,
erosion number, 208	102–103
erosion rate constants, 202–203	PPA, 200
experiments, 200–206	PPDP, 174, 179, 191
from unsaturated and fatty acid-derived	Protein adsorption, 26–28, 34, 39
monomers, 177–179	Protein-based hydrogels, 120
FTIR, 189–191	Protein conformation, 26–27
future opportunities, 213–214	Protein delivery
<sup>1</sup> H NMR, 189–191	and gene therapy, 132-133
implantable systems, 210	limitations, 132–133
in vitro degradation, 201	Protein diffusion coefficient, 27
injectable systems, 211–212	Protein diffusivity, 27
laminated devices for pulsatile	Protein–surface interactions, 28
release, 210	Proteoglycans, 13–15, 39–41
microstructure, 195–199	PSA, 175, 177–178, 191–195, 198, 200–201,
model formulations testing in vitro and	204–205
in vivo, 209	FTIR spectrum, 189
modeling, 207–209	P(SA-alt-TA), 179
modulating erosion rates, 201–205	PSA-fatty acid-based copolymers, 179
phenomenological models, 208–209	PTA, 173, 175, 179, 191–192, 194, 199
solubility, 191–192	PTFE, 104 PTMCSA 188
surface changes during erosion, 205–206 synthesis for drug delivery, 173–176	PTMCSA, 188 PTPA, 179
thermal properties, 196–197	PVA, 34, 103–105, 107–109, 111, 118, 145,
thermal transitions, 192–193	148, 152, 161
Poly[ $\alpha$ , $\omega$ ( $p$ -carboxyphenoxy)alkanes], 173	PVP, 145, 152
Poly(azelaic acid) (PAZ), 200	PXDA, 179
Poly(bis(hydroxyethyl)terepthalate-ethyl	
ortho-phosphate/terephthaloyl	
chloride), 58	R
chloride), 58 Polydimethylsiloxane, 48	R
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92	Receptor affinity, 11
Polydimethylsiloxane, 48	<del></del>
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92	Receptor affinity, 11
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate)	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9 REDV (Arginine–Glutamine–Aspartic acid–Valine), 11, 38
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9 REDV (Arginine–Glutamine–Aspartic acid–Valine), 11, 38 Responsive networks, 110–119
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9 REDV (Arginine–Glutamine–Aspartic acid–Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine–Glycine–Aspartic acid), 11,
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9 REDV (Arginine–Glutamine–Aspartic acid–Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine–Glycine–Aspartic acid), 11, 37–38, 63, 104
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor–ligand binding, 9 REDV (Arginine–Glutamine–Aspartic acid–Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine–Glycine–Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared spectroscopy, 97–98	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared spectroscopy, 97–98 Polymeric biomaterials for nerve	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176 Rubber elasticity theory, 81–82
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared spectroscopy, 97–98 Polymeric biomaterials for nerve regeneration, 47–74	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared spectroscopy, 97–98 Polymeric biomaterials for nerve regeneration, 47–74 Poly(phosphoester) conduits, 57	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176 Rubber elasticity theory, 81–82
Polydimethylsiloxane, 48 Polyelectrolyte complexes, 92 Polyesters, 169 Polyethers, 169 Poly(glyceryl methacrylate) p(GMA)-collagen hydrogels, 63 Poly(glycolic acid), 50 Poly(lactic acid) (PLA), 50, 186, 195 Polylactic-co-glycolic acid, 57 Poly(lactide)-co-glycolide, 65–66 Poly(lithocholic acid), 189 Polymer-based particles for controlled DNA release, 141–157 Polymer complexation and infrared spectroscopy, 97–98 Polymeric biomaterials for nerve regeneration, 47–74	Receptor affinity, 11 Receptor binding to ligands, 10 Receptor diffusivity, 10 Receptor-ligand binding, 9 REDV (Arginine-Glutamine-Aspartic acid-Valine), 11, 38 Responsive networks, 110–119 RGD (Arginine-Glycine-Aspartic acid), 11, 37–38, 63, 104 Ricinoleic acid, 179 Ricinoleic half-esters, 179 Ring opening polymerization, 176 Rubber elasticity theory, 81–82

Scaffolds biodegradable polymer, 61 DNA release from, 161–162 for controlled DNA delivery, 157–162 tissue engineering, 162	Thrombogenicity, 3 Thrombospondins (TSPs), 17 Tissue attachment, chemical and physical determinants, 103–105 Tissue sulture reliestures (TCPS) us
Schwann cells, 48, 50, 52, 58, 61	Tissue culture polystyrene (TCPS) vs. laminin-coated surfaces, 8
Secreted protein acidic and rich in cysteine	Tissue engineering
(SPARC), 17	neutral networks, 98–105
Selectins, 9, 13–15, 39–41	scaffolds, 162
Severe combined immunodeficiency	Tissue–implant interactions, 101–105
(SCID), 131	Tissue regeneration, 162
Sialyl-Lewis <sup>x</sup> (sLe <sup>x</sup> ), 41	Tissue response to porous materials,
Signal transduction, 18–25	102–103
Silicone, 60	Transforming growth factor $\beta$ (TGF- $\beta$ ), 159
Silicone chambers, 49–50	Transmembranous domain, 9
Silicone conduits, comparative studies, 59	Trifluoroethanol (TFE), 151
Silicone rubber, 48	Trophic factors, 48, 50
Silicone tubes, 50	Tyrosine-phosphorylated proteins, 20
electrical circuit within, 61	
prefilled with NGF, 59	$\mathbf{U}$
Small-angle X-ray scattering (SAXS), 194	O
Sodium chloride, 66	Ulex europaeus I (UEA I), 40
Solute transport	Unsintered hydroxyapatite/poly-L-lactide
in hydrogels, 83–87	(u-HA/PLLA), 62
in matrix-based systems, 87	(u-11/1/1 LL/1), 02
Specificity, 5	
Spinal cord injuries, 47	$\mathbf{V}$
Spinal cord regeneration, 61–62	
Spinal cord tissue, 60–61	Vaccination applications, DNA release in,
Spray-drying technique, 149–150	153–157
Star polymers, 121	VCAMs, 16
Stereocomplexes, 92	Viral gene therapy, 135
Styrene, 119	VP/AAm matrix, 95
Surface chemistry, 28–30	
Surface-erodible biomaterials for drug	W
delivery, 169–218 Surface erosion, 172	**
Surface closion, 172 Surface plasmon resonance (SPR), 195	Weibel-Palade bodies, 15
Surface receptors, 11, 24	Wide-angle X-ray diffraction (WAXD), 195,
Surface topography, 31–33	198
Syndecans, 17	Wound healing, 158–159
Sylidectalis, 17	
T	X
Teflon, 60	X-ray diffraction (XRD), 194
Tenascins, 17	
Tetrahydrofuran (THF), 191–192	•7
Three-dimensional polymeric networks	$\mathbf{Y}$
as biomaterials, 76–105	VICCD 104

structural characteristics, 82–83 YISGR, 104