

INDEX

A

- Acetylintermediate, 72, 75
Acetyllycopsamine, 72, 75
Acoustic impedance, 117–118
Acoustic signal generator, 122
Actin–DNA cross-links, pyrrolizidine
alkaloid antimitotic activity, 90–91,
92
Adenostyles alliariae (Alpendost), 76
Aegilops cylindrica, 228
Aflatoxins, 198
Agrobacterium tumefaciens transformation
vector, 254
Alfalfa sprouts, ozone treatment, 196
Alginate gels, ultrasonic analysis, 152
Alicyclobacillus acidocaldarius, 197
Alkanes, ultrasonic phase transition
monitoring, 141
Alpendost (*Adenostyles alliariae*), 76
 α -gliadins, 222
 α -trinositol
(inositol(1,2,6)trisphosphate), 28, 40
Aluminium–inositol phosphate
interactions, 32, 36
Animal products, inositol phosphate
content, 26–27
Anticaking agent, gaseous ozone
treatment, 179–180, 198
Anticancer activity, inositol phosphates,
37, 38–39
Antioxidants
inositol phosphates, 28, 29
anticancer activity, 39
ozone interactions, 177
AP-2 (adaptor protein 2), 25, 30, 38
AP-3 (adaptor protein 3), 30
Apple juice, fruit washing water-related
disease outbreaks, 169, 195
Apples, ozone treatment, 192, 195
of washing water, 190
Arabidopsis
Escherichia coli phytase transgenic
expression, 34
phytate (inositol hexakisphosphate)
synthetic pathway, 8
Argon, ozone combined sanitization
procedures, 201
Aspergillus fumigatus, phytase transgenic
expression, 34
Aspergillus niger, 11, 12, 36
phytase, 20
Asteraceae (Compositae), pyrrolizidine
alkaloids, 65
Avocado, inositol phosphate content, 23

B

- Bacillus cereus*, 186
Bacillus spp., 186, 198
Bacillus stearothermophilus, 186, 187
Bacillus subtilis, 187, 199, 202
phytase, 12
Barley
inositol phosphate degradation during
processing, 21
malting, 22
phytases, 9

- reduced phytate gene mutations, 34
 - Beet root, inositol phosphate content, 23
 - β -gliadins, 222
 - Betony (*Pedicularis*), 77
 - BIOTEST, 138
 - Black beans, inositol phosphate degradation
 - during processing, 21, 22
 - Blazing star (*Liatris punctata*), 77
 - Boraginaceae, pyrrolizidine alkaloids, 65
 - Botrytis cinerea*, 181, 191
 - Bread, ultrasonic texture sensing, 146, 147
 - Breadmaking
 - inositol phosphates degradation, 16, 20
 - see also Glutenin high molecular weight subunits
 - Broccoli, ozone treatment, 192
 - Brussel sprouts, chlorination, 188
- C**
- Cabbage
 - chlorination, 188
 - inositol phosphate content, 23
 - Cachana (*Liatris punctata*), 77
 - Cadmium–inositol phosphate interactions, 32
 - Calcium, inositol phosphate effects, 27, 29, 30–31
 - absorption impairment, 32, 35
 - inositol(1,4,5)PU3u signaling, 27–28
 - Campylobacter* spp., 193
 - Canning
 - filling operations, 130
 - phytate degradation, 16
 - Capillary chromatography, inositol phosphates, 6
 - Carbon dioxide, ozone combined sanitization procedures, 201–202
 - Cardiopulmonary toxicity, pyrrolizidine alkaloids, 80–81
 - Carrots, inositol phosphate content, 23
 - Casein, ultrasonic micelle particle sizing, 150
 - Castellija rhexifolia*, 77
 - Castellija* sp. (Indian paint brush), 77
 - Cauliflower, ozone treatment, 192
 - Celery, inositol phosphate content, 23
 - Cereal grains
 - inositol phosphates, 5–6
 - degradation during processing, 16, 21–22
 - ozone treatment, 198
 - pyrrolizidine alkaloids contamination, 68, 69–70
 - Cheese
 - ultrasonic texture sensing, 146, 147
 - ultrasonic velocity changes during melting, 142
 - Chickpeas, inositol phosphate degradation
 - during malting, 22
 - Chinese millet, phytate content, 13
 - Chlorine dioxide sanitization, 188, 189
 - Chlorine sanitization, 169–170, 188–189
 - ozone combined procedures, 200–201
 - Chromium–inositol phosphate interactions, 36
 - Cider
 - fruit washing water-related disease outbreaks, 169, 195
 - pasteurization recommendations, 195
 - Coatmer, 30
 - Cobalt–inositol phosphate interactions, 36
 - Coltsfoot (*Tussilago farfara*), 76, 91
 - Comfrey (*Symphytum*), 74–76, 91
 - Commelina communis*, phytate (inositol hexakisphosphate) synthetic pathway, 8
 - Compositae (Asteraceae), pyrrolizidine alkaloids, 65
 - Compressibility measurements, polymer hydration analysis, 151–152
 - Cookies, ultrasonic texture sensing, 146
 - Copper
 - hepatotoxic synergism with pyrrolizidine alkaloids, 79
 - phytate (inositol hexakisphosphate) interactions, 32, 35–36, 40
 - Corn
 - bran/flour phytate content, 16
 - inositol phosphate degradation during processing, 21
 - Coughwort (*Tussilago farfara*), 76
 - Crotalaria*, 80
 - contamination of staple foods, 69, 70
 - prevention, 91
 - Crotalaria fulva*, 74, 91
 - Cryptosporidium parvum*, 169, 189
 - Cucumbers, ozone/carbon dioxide mixed gas treatment, 202
 - Cyclospora cayatanensis*, 169
 - CYP 2C11, 82
 - CYP 3A, 82

CYP 3A4, 82

Cytochromes P450, pyrrolizidine alkaloids metabolism, 81, 82

D

Dairy products, ultrasonic composition sensors, 138

Dictyostelium

phytase, 12

phytate (inositol hexakisphosphate)

synthetic pathway, 7

Dietary Supplement Health and Education

Act (1994), 74, 91

Digital storage oscilloscope, 122–123

DNA cross-links

formation during pyrrolizidine alkaloid metabolism, 83, 88–90

DNA–protein cross-links, 90, 93

DNA repair, inositol phosphate requirement, 31

Doppler ultrasound

flowmeters, 131–132

viscosity determination, 148, 149

E

Echimidine, 71, 72, 75

Echium plantagineum (Patterson's Curse; Salvation Jane), 72

Echiumine, 72

Eggs

ozone inactivation of microbial contaminants, 186, 193–194
pyrrolizidine alkaloid concentrations, 73, 91

Elastography, carcass grading, 154

Elephant Head (*Pedicularis*), 77

Emulsions, ultrasonic characterization, 112, 114–116, 158

flocculated emulsions, 116, 150

imaging, 154

particle sizing, 115–116, 150

practical applications, 150

Epi-inositol, 37

Escherichia coli, 178, 180, 183, 193

phytase, 9, 10, 11, 21

transgenic expression in *Arabidopsis*, 34

Escherichia coli O157:H7, 169, 178, 181, 188, 190, 195, 203

Europine, 73

F

Fabaceae (Leguminosae), pyrrolizidine alkaloids, 62

Fermentation, phytate degradation, 9, 16, 20, 22

Fish

ozone treatment, 197–198

ultrasonic composition sensors, 138

ultrasonic microscopy for parasite detection, 156

Flavin-containing monooxygenases, 82

Flavobacterium aquatile, 180

Flavobacterium branchiophilum, 181, 190

Fluid flow, ultrasonic sensing, 130–132

Doppler flowmeters, 131–132

transit-time mode, 130–131

vortex shedding, 132

Foams, ultrasonic analysis, 150–151

Foil pouches, ultrasonic leak detection, 154

Food processing equipment, ozone treatment, 192

Foreign bodies, ultrasonic detection, 151

Fourier transform infrared (FT-IR)

spectroscopy, gluten proteins, 239, 278

Fruit

chlorine treatment, 188

inositol phosphates, 22–23

ozone treatment, 190, 191, 192, 194–195

pesticide residues reduction, 199

texture (rheological properties),

ultrasonic sensing, 142

washing water-related disease outbreaks, 169, 194

Fruit juices

Love wave-based component analysis, 153

ozone treatment of ingredients, 196–197

Fulvine, 71, 74

G

Gabo near isogenic line, 252

dough properties, 260, 261

transgenic HMW glutenin subunit studies, 263–264

Galahad near isogenic line, 252

γ -gliadins, 222

Garlic, ozone treatment, 198

Gelation, ultrasonic analysis, 152

- Gliadins, 221
 dough viscoelasticity
 gluten fractionation/reconstitution
 experiments, 242, 243, 245
 protein fraction incorporation studies,
 245–246
 electrophoretic separation, 222, 223
 structural studies, 278
Glu-A1, 224, 226, 256–257, 258, 260, 263
 near isogenic lines, 251, 252
Glu-A3, 258
Glu-B1, 224, 226, 257, 258, 262, 263
 near isogenic lines, 251, 252
Glu-B3, 258
Glu-D1, 224, 226, 256, 257, 258, 260, 262
 near isogenic lines, 251, 252
Glu-D3, 258
 Glutathione metabolism, 79
 Glutathione *S*-transferase, 83
 Gluten
 insolubility, thermodynamic aspects,
 274–275, 276
 macropolymer, 258–259
 dough performance relationship, 259
 viscoelasticity, 279–280
 Glutenin high molecular weight subunits,
 222, 223
 amino acid sequences, 228–233
 atomic force microscopy, 233, 235
 composition manipulation, 249, 251–256
 cross-link formation
 disulphide bonds, 240–242, 278, 284
 low molecular weight subunit
 branches, 240, 242, 278
 mechanical behaviour of dough, 280
 transgenic lines, 265
 tyrosine bonds, 242
 dough viscoelasticity influence, 242–249
 dough mixing process, 284–288
 gluten fractionation/reconstitution
 experiments, 242–245
 in vitro polymerization, 246
 ‘loop and train’ hypothesis, 281–283,
 284, 285, 286, 288–289
 macropolymer interactions, 280–281
 protein fraction incorporation studies,
 245–247
 purified subunits incorporation into
 dough, 247–249
 subunit peptides incorporation into
 dough, 249, 250
 theoretical models, 279–284
 genetics, 224–226
 grain breadmaking quality relationship,
 226–228, 259–265, 266
 biophysical models, 273–289
 gluten macropolymer composition, 259
 molecular basis, 265, 267–273
 quality scoring system, 227
 ranking of subunit loci, 258
 relation to dough rheology, 279
 variation between cultivars, 256–259
 hydration-related molecular
 rearrangements, 278–279, 284
 insolubility, thermodynamic aspects,
 274–278
 mixing/rheological properties influence,
 256–265
 molecular basis of processing properties
 amino acid sequences, 267–269, 271
 subunit interactions, 271–273
 subunit stability, 270, 271
 total amount of protein, 267
 molecular dimensions, 233, 234
 near isogenic lines, 251–253
 dough quality differences, 259–262
 Gabo, 252, 260, 261
 Galahad, 252
 Pegaso, 252–253
 Sicco, 251–252, 259–261
 single subunit, 253
 polymorphisms, 224–226
 SDS-polyacrylamide gel
 electrophoresis, 224, 225
 structure
 repetitive domains, 237–239
 terminal domains, 234, 236–237
 transgenic lines, 253–256
 dough quality differences, 262–265,
 266
 Glutenin low molecular weight subunits,
 222, 223
 B-type, 222
 C-type, 222
 cross-links, 240, 242, 278
 mechanical behaviour of dough, 280,
 281
 D-type, 222
 dough viscoelasticity, incorporation
 studies, 247
 genetic variation, influence on
 technological properties, 257, 258

gluten macropolymer composition, 259
in vitro polymerization, 246
 Glutenins, 221
 dough viscoelasticity
 gluten fractionation/reconstitution
 experiments, 242–245
 protein fraction incorporation studies,
 245–246
 electrophoretic subunit separation, 222,
 223
 subunits *see* Glutenin high molecular
 weight subunits; Glutenin low
 molecular weight subunits
 Grey feather (*Liatris punctata*), 77
 Guided wave sensors, 128

H

Heleurine, 73
 Heliosupine, 66, 67
 Heliotrine, 70, 71, 73, 78
 Heliotrine *N*-oxide, 78
Heliotropium contamination of staple
 foods, 68, 70
 prevention, 91
Heliotropium europaeum, 73
Heliotropium popvoui, 70
 Hepatic veno-occlusive disease,
 pyrrolizidine alkaloid toxicity,
 69–70, 72, 75–76, 77, 79–80, 91
 herbal teas, 74
 synergism with copper, 79
 Hepatitis A, 178
 Herbal medicines, pyrrolizidine alkaloids
 adverse health effects awareness, 91, 93
 sources, 66, 68, 73–78
 High-performance liquid chromatography
 (HPLC), inositol phosphates
 analysis, 2, 5–6, 13, 41
 Honey
 pyrrolizidine alkaloids contamination,
 72, 91
 ultrasonic viscosity measurement, 147
Hordeum chilense, 256
 Hydrogen peroxide sanitization, 189
 ozone combined procedures, 200–201
 Hypochlorite sanitization, 169–170

I

Indian childhood cirrhosis, 79
 Indian paint brush (*Castellija*), 77
 Indian Warrior (*Pedicularis*), 77

Indicine *N*-oxide, 66, 67, 88
 Inositol hexakisphosphate (InsPU6u;
 phytate), 1–2
 analytical methods, 4–5
 anticancer activity, 37, 38–39
 antioxidant activity, 23, 39
 food preservative effects, 39–40
 blood glucose-lowering effects, 40
 cholesterol/triglyceride-lowering effects,
 40
 degradation, 8–12
 digestive processes, 9, 24
 food processing/preparation, 9, 16
 thermal, 12
 extraction during seed processing, 13
 functions in animal tissues, 30–31
 calcium flux regulation, 30–31
 DNA repair, 31
 nuclear mRNA export, 31
 high-performance liquid
 chromatography (HPLC), 5, 6
 mineral absorption impairment, 2, 32–33
 iron, 32, 33–34, 40–41
 optimal dietary levels, 42
 seed content
 processed foods, 16–22
 reduction strategies, 34–35
 whole raw seeds, 12–15
 synthetic pathway, 7–8
 Inositol kinases, 7, 8, 31–32
 Inositol phosphate phosphatases, 11, 12
 Inositol phosphates, 1–43
 analytical methods, 4–6
 animals, 23–32
 absorption, 23–25
 biological functions, 27–32
 tissue content, 26–27
 anticancer activity, 37, 38–39
 antioxidant activity, 23, 28, 29, 30, 39
 calcium mobilisation, 27–28, 29, 30–31
 chemistry, 2–6
 chloride channel regulation, 29
 degradation, 8–12
 breadmaking processes, 16, 20
 digestive processes, 9, 24
 malting, 22
 fruit/vegetable content, 22–23
 hemoglobin oxygen affinity regulation,
 30
 iron binding, 28–29, 30
 metabolism, 7–12

- nomenclature, 2–3
 - nutritional importance, 32–41, 42
 - health disorders prevention, 36–41
 - mineral bioavailability influence, 32–36, 42
 - receptors, 27, 28
 - seed contents, 12–23
 - processed foods, 16–22
 - whole raw seeds, 12–15
 - signaling activities, 7, 27, 28, 29
 - synthesis, 7–8
 - animal cells, 7–8
 - microorganisms, 7
 - plants, 8
 - Inositol pyrophosphates, 31
 - Inositol(1,2,6)trisphosphate (alpha-trinositol), 28, 40
 - Inositol(1,4,5)trisphosphate (Ins(1,4,5)PU3u), 5, 7
 - animal cells, 26
 - receptors (calcium channels), 27–28
 - signaling function, 27–28
 - plant cells, 23
 - Intermedine, 71, 72, 75, 77
 - Iron, inositol phosphates binding, 28–29
 - absorption impairment, 32, 33–34, 40–41
 - low-phytic acid grains, 34–35
 - anticancer activity, 38–39
 - overall dietary absorption impact, 40–41, 42
- J**
- Jack mackerel, ozone treatment, 197
 - Jacobine, 72
 - Jacoline, 70, 72
 - Jaconine, 72
 - Jacozine, 72
 - Japanese flounder, ozone treatment, 197
- K**
- Kimchi preparation, 192
- L**
- Lactobacillus leichmannii*, 203
 - Lactobacillus plantarum*, 22
 - Lasiocarpine, 71, 73
 - Lead–phytate (inositol hexakisphosphate) interactions, 32
 - Legumes, phytate degradation during processing, 16
 - Leguminosae (Fabaceae), pyrrolizidine alkaloids, 62
 - Lentils, inositol phosphates degradation, 20, 22
 - Lettuce
 - chlorination, 188
 - ozone treatment, 192, 195–196, 206
 - Leuconostoc mesenteroides*, 188
 - Level sensors, ultrasonic, 129–130
 - Liatris punctata* (cachana; grey feather; blazing star), 77
 - Lily pollen phytase, 11
 - Listeria monocytogenes*, 169, 188, 193, 203
 - Lousewort (*Pedicularis*), 77
 - Love wave sensors, 128–129, 147
 - arrays, 129
 - smart tongue and nose, 153
 - Love waves, 104
 - Low-phytic acid grains, 34–35
 - Lupin, inositol phosphates degradation, 22
 - Lycopsamine, 71, 72, 75, 77, 78
- M**
- Magnesium absorption, phytate (inositol hexakisphosphate) impairment, 32, 35
 - Maize
 - phytases, 9
 - phytate (inositol hexakisphosphate), 9
 - degradation during processing, 22
 - reduced content gene mutations, 34
 - Malting, inositol phosphates degradation, 22
 - Manganese absorption, inositol phosphate effects, 36
 - Meat
 - elastographic carcass grading, 154
 - foodborne disease outbreaks, 169
 - ozone treatment, 186, 190, 191, 192, 193–194
 - protein ultrasonic analysis, 152–153
 - pyrrolizidine alkaloid residues, 73
 - sanitizing agents, 189
 - ultrasonic composition sensors, 138
 - ultrasonic imaging of carcasses, 154
 - Megalocytosis, pyrrolizidine alkaloid-induced, 88, 89
 - Mercury–inositol phosphate interactions, 32, 36

Micrococcus spp., 198

Milk

contamination with pyrrolizidine
alkaloids, 70, 72, 91

ultrasonic analysis

aggregates, 152

fat globule particle sizing, 150

imaging for spoilage measurement,
154

Mineral bioavailability, inositol phosphate
influences, 32–36

Mixograph, 285

Molybdenum–inositol phosphate
interactions, 36

Monocrotaline, 66, 67, 78, 79, 80, 88

Mucor piriformis, 181

Multiple inositol polyphosphate

phosphatase (MIPP), 11, 12

Multiple scattering theory, 113–116

Mung bean, phytate (inositol

hexakisphosphate) synthetic
pathway, 8

Mussels, ozone treatment, 197

Mycobacterium fortuitum, 173

Myelin proteolipid protein, 30

Myo-inositol, 3, 36

anticancer activity, 37, 38

food sources, 36–37

plasma lipid effects, 40

N

Necic acids, 66

Neosartorya fischeri, 197

Neper (Np), 106

Nickel–phytate (inositol

hexakisphosphate) interactions, 32,
36

Nuclear magnetic resonance (NMR)
spectroscopy

high molecular weight glutenin subunits,
272–273, 279, 282

domain structures, 236, 237, 239

inositol phosphates, 6

whole raw seed studies, 13

O

Oats

inositol phosphate degradation during
processing, 21

zinc absorption influence, 33

Oils, ultrasonic analysis

particle sizing, 150

spoilage correlations, 153

viscosity, 147

Ω -gliadins, 222

structural studies, 278

Oncorhynchus mykiss (rainbow trout)

culture, ozone water treatment, 181,
190

Onions

inositol phosphates content, 23

ozone treatment, 198

Oranges, ozone/carbon dioxide/argon
mixed gas treatment, 201

Ozone, 167–208

analytical methods, 204–205

atmospheric, 170–171

biological oxygen demand (BOD) of
water, 170

chemical oxygen demand (COD) of
water, 170

chemistry/physics, 170–177

cycloaddition reactions, 175

decomposition, 176–177

catalysis by inorganic compounds,
177

free radical species generation, 176

inhibition, 177

initiation, 177

promotion, 177

electrophilic reactions, 175

microbial inactivation, 170

biofilms, 187, 198–199

effect of oxygen demand of medium,
181, 182

food characteristics influencing
efficiency, 190, 191

food sanitizing applications, 189
kinetics, 187–188

mechanisms, 187–188

pulsed electric field combined
treatment, 188, 203–204

relative humidity/microbial cell
hydration, 179

residual ozone values, 180–181

spores, 186, 187

targeted microorganisms, 186–187

nucleophilic reactions, 175

oxidation–reduction potential, 171

pesticide residue reduction, 170, 189, 199

- physical properties, 171
 - processing equipment interactions, 183–185
 - materials compatibility, 184
 - metals, 184–185
 - plastics, 185
 - rubber, 185
 - production methods, 204
 - reactivity, 175–177
 - inorganic compounds oxidation, 177
 - molecular ozone, 175
 - solubility, 172–173
 - stability, 173–175
 - structure, 171
 - toxicity, 206–207
 - transfer from gas to liquid phase (contacting systems), 182–183
 - treatment chambers, 182
 - treatment medium, 178–181
 - oxygen demand, 170, 181, 182
 - relative humidity, 178–180
 - residual ozone, 180–181
 - temperature, 178
 - workplace exposure thresholds, 207–208
 - Ozone sanitization
 - combination treatments, 200–204
 - carbon dioxide, 201–202
 - chlorine, 201
 - heat, 202
 - hydrogen peroxide, 200–201
 - pulsed electric field, 188, 203–204
 - ultraviolet radiation, 202–203
 - dry foods/ingredients, 198
 - equipment treatment, 192
 - filtration of treatment water, 183
 - fish, 197–198
 - food processing applications, 186–199
 - chlorine treatment alternative, 188–189
 - environmental contaminants reduction, 192
 - food characteristics influencing ozone-demand, 190–191
 - food surface structure, 190–191
 - limitations, 206
 - safety issues, 207
 - fruit, 190, 191, 192, 194–195
 - meat, 186, 189, 190, 191, 192, 193–194
 - metal contaminants removal from drinking water, 177
 - packaging materials, 192, 198–199
 - pesticide residue reduction, 170, 189, 199
 - poultry carcasses, 183, 190, 192, 193–194, 201
 - raw products/ingredients, 191–192
 - regulatory status, 206
 - shelf-life extension, 193
 - storage applications (ozone gas), 178
 - surface decontamination, 178
 - food contact surfaces, 198–199
 - vegetables, 190, 191, 194, 195–196
- P**
- Packaging materials, ozone treatment, 192, 198–199
 - Packeria candidissima*, 77–78
 - Paramecium* phytase, 11
 - Parenteral fat emulsions, ultrasonic particle sizing, 150
 - Parsnips, inositol phosphates content, 23
 - Patternson's Curse (*Echium plantagineum*), 72
 - Pea flour, inositol phosphate degradation during processing, 21
 - Peanuts
 - aflatoxins, ozone treatment reduction, 198
 - phytate content, 13
 - Pedicularis* (betony), 77
 - Pegaso near isogenic line, 252–253
 - Penicillium expansum*, 191
 - Peniophora lycii* phytase, 10, 11
 - Pepper, ozone treatment, 198
 - Peracetic acid food sanitization, 189
 - Pesticide residues, reduction by ozone treatment, 170, 189, 199
 - Petasitenine, 66, 67
 - Petasites officinalis*, 76
 - Phaseolus vulgaris* L., iron availability of low-phytic acid varieties, 34–35
 - Phonons, 103
 - Phosphatases, 24
 - Phospholipase C, 27
 - Phytases, 8–9
 - activation in seeds/grains before cooking, 9
 - animal digestive tract, 24
 - cell/tissue activity, 9–10
 - animal cell lines, 24

- classification, 11
 - phytate degradation during food processing, 16, 20, 21
 - purified enzymes activity, 10
 - Phytate *see* Inositol hexakisphosphate
 - Phytophthora parasitica*, 181
 - Pipe fouling, ultrasonic detection, 153
 - Plant pyrrolizidine alkaloid sources, 62–65
 - Plastic pouches, ultrasonic leak detection, 154
 - Polymer hydration, ultrasonic analysis, 151–152
 - Potatoes, inositol phosphate content, 23
 - Poultry carcasses, ozone treatment, 190, 192, 193–194
 - hydrogen peroxide combined treatment, 201
 - prefiltration of treatment water, 183
 - residual ozone values, 180
 - Protein hydration, ultrasonic analysis, 151–152
 - Pseudomonas aeruginosa*, 180
 - Pseudomonas fluorescens*, 180, 188, 199
 - Pseudomonas putida*, 180
 - PTEN, 29
 - Pulsed electric field/ozone combined sanitization procedures, 188, 203–204
 - Pulsed mode ultrasonic measurements, 123–124
 - Punctanecine, 77
 - Pyrrolizidine alkaloids, 61–93
 - carcinogenic potential, 62, 66, 78, 81
 - chemical structure, 65–66, 67, 71
 - consumer protection measures, 69, 73–74, 91–93
 - food contamination, 66, 68–73
 - cereal grains contaminated with weeds, 68, 69–70, 91
 - eggs, 73, 91
 - honey, 72, 91
 - meat, 73
 - milk, 70, 72, 91
 - preventive measures, 91
 - herbal medicine sources, 66, 68, 73–78
 - adverse health effects awareness, 91, 93
 - metabolism, 81–83, 84–85, 86–87
 - dehydrogenation, 82
 - detoxification, 83
 - N-oxidation, 82–83
 - plant sources, 62–65
 - toxicity, 62, 78–81
 - acute, 78
 - cardiopulmonary, 80–81
 - diagnostic problems, 81
 - DNA cross-links formation, 83, 88–90
 - DNA–protein cross-links, 90, 92
 - hepatic veno-occlusive disease, 69–70, 72, 75–76, 77, 79–80, 91
 - hepatotoxic synergism with copper, 79
 - human poisoning incidents, 62, 68, 73, 75–77
 - mechanism, 83, 88–91
 - megalocytosis, 88, 89
 - progressive nature, 79–80
 - species differences in susceptibility, 78–79
 - structural determinants, 65
 - transplacental transmission, 76
- Q**
- Quinoa flour, inositol phosphates
 - degradation during processing, 22
- R**
- Rainbow trout (*Oncorhynchus mykiss*)
 - culture, ozone water treatment, 181, 190
 - Raleigh waves, 104–105
 - Retronecine, 66, 67
 - Retrorsine, 78, 79, 80, 88
 - Rice
 - Aspergillus fumigatus* phytase transgenic expression, 34
 - phytate (inositol hexakisphosphate) content, 12, 13, 16
 - reduced phytate gene mutations, 34
 - Riddelliine, 80, 88
 - Rockfish, ozone treatment, 197
 - Russian comfrey (*Symphytum uplandicum*), 74–75
 - Rye
 - phytase, 9, 21
 - phytate (inositol hexakisphosphate), 9
 - degradation during processing, 21
- S**
- Saccharomyces cerevisiae* phytase, 11
 - Salad dressings, ultrasonic particle sizing, 150
 - Salmon, ozone treatment, 192

- Salmonella enterica* ser. Enteritidis, 193, 194
Salmonella spp., 169, 183, 193, 194
Salmonella typhimurium, 180, 203
 Salvation Jane (*Echium plantagineum*), 72
 Sanitization methods
 ozone applications *see* Ozone
 sanitization
 requirements for food safety, 169–170
 Scanning acoustic microscopy, 156
Schizosaccharomyces pombe, phytate
 (inositol hexakisphosphate)
 synthetic pathway, 7
Scyllo-inositol, 37
 Seeds, inositol phosphates content, 12–15
 Selectins, 30
 Selenium absorption, phytate (inositol
 hexakisphosphate) effects, 32, 35
Senecio, 80
 contamination of staple foods, 68, 69–70
 prevention, 91
Senecio burchelli, 69
Senecio disease, 69
Senecio ilicifolius, 69
Senecio jacobaea (tansy ragwort), 70, 72
Senecio longilobus, 77
Senecio riddellii, 65
Senecio triangularis, 77
 Senecionine, 66, 67, 72, 76, 77, 78, 80, 88
 metabolism, 82, 84–85
 Seneciophylline, 71, 72, 76, 78, 80, 88
 Senkirkine, 66, 67
 Shelf-life extension, ozone treatment, 193
 Shrimp meat, ozone treatment, 197–198
 residual ozone values, 180
 Sicco near isogenic line, 251–252
 dough properties, 259–261
 Sockeye salmon, ozone treatment, 197
 Solid fat content, ultrasonic sensing
 isothermal scanning, 139–140
 temperature scanning, 141
 Sorghum flour, phytate content, 16
 Sour dough breads, phytate content, 20
 Soybean
 inositol phosphates
 calcium absorption influence, 35
 degradation during malting, 22
 phytate (inositol hexakisphosphate), 12
 reduced content gene mutations, 34
 synthetic pathway, 8
Spirodela polyrhiza, 8, 23
Staphylococcus aureus, 180
 Strawberries, ozone treatment, 192
 Sunflower seeds, phytate content, 13
 Supinine, 73
 Symphytine, 75
Symphytum (comfrey), 74–76, 91
Symphytum uplandicum (Russian comfrey),
 74–75
 Symviridine, 75
 Synaptotagmin, 30
- ### T
- Tansy ragwort (*Senecio jacobaea*), 70, 72
 Temperature, ultrasonic sensing, 132–133
 Texture, ultrasonic sensing, 142–148
 viscosity measurements, 147–148, 149
 Thermal degradation, inositol phosphates,
 16
 Thermal sanitization, ozone sequential
 procedures, 202
 Tomato sauce, ultrasonic Doppler
 velocimetry, 148, 149
 Tomatoes
 ozone treatment, 191
 washing water-related salmonellosis
 outbreaks, 169
 Transgenic wheat
 breadmaking quality differences, 262–265
 transformation system, 253–254
 Trichodesmine, 67, 78, 80
 Triglycerides, ultrasonic phase transition
 monitoring, 141
Triticum dicoccoides, 253
Triticum monococcum ssp. *boeoticum*, 226
Triticum monococcum ssp. *monococcum*,
 226
Triticum timopheevi, 228
Triticum tsudchii, 228
Triticum turgidum ssp. *dicoccoides*, 226
Triticum urartu, 226
 Tritordeum
 transformation, 255–256
 transgenic line breadmaking qualities, 263
 Turnip, inositol phosphates content, 23
Tussilago farfara (coltsfoot; coughwort),
 76, 91
- ### U
- Ultrasonic composition sensors, 133–139
 application to animal carcasses, 138–139

- Ultrasonic flow meters, 130–132
 - Doppler flowmeters, 131–132
 - transit-time mode, 130–131
 - vortex shedding, 132
 - Ultrasonic imaging, 153–157
 - carcass grading, 154
 - high-resolution, 156–157
 - limitations, 155
 - noncontact methodology, 155
 - scanning acoustic microscopy, 156
 - Ultrasonic measurement systems, 121–129
 - air–transducer impedance barrier, 124
 - digitizer, 122–123
 - guided wave sensors, 128
 - Love wave sensors, 128–129
 - noncontact measurements, 124–126
 - pulsed modes, 123–124
 - pathlength measurement, 124
 - reflectance methods, 127–128
 - resonance methods, 126–127
 - resonance cell losses (Q -factor), 127
 - signal generator, 122
 - ultrasonic transducers, 122
 - Ultrasonic phase transition monitoring, 139–142
 - isothermal scanning, 139–140
 - temperature scanning, 140–142
 - Ultrasonic sensors, 101–158
 - advantages/disadvantages, 157–158
 - applications, 129–157
 - composition, 133–139
 - dispersed systems, 148, 150–151
 - fluid flow, 130–132
 - imaging, 153–157
 - level sensors, 129–130
 - phase transitions, 139–142
 - polymeric systems, 151–153
 - rheological properties (texture), 142–148
 - temperature, 132–133
 - viscosity measurements, 147–148
 - nondestructive testing, 104
 - theoretical aspects, 103–121
 - Ultrasonic thermometry, 132–133
 - Ultrasonic transducers, 122
 - Ultrasonic wave propagation, 103–104
 - attenuation, 105, 109–111
 - coefficient, 105–106
 - overestimation, 120
 - relaxation, 111
 - diffraction, 120–121
 - homogeneous materials, 105–111
 - adiabatic elastic modulus, 106
 - properties of materials, 106, 107–108
 - inhomogeneous media, 112–117
 - emulsions, 112, 114–116
 - floculated emulsions, 116–117
 - multiple scattering theory, 113–116
 - particle sizing, 115–116
 - scattering interactions, 112–113
 - interfaces, 117–120
 - acoustic impedance, 117–118
 - reflection coefficient, 118, 119
 - refractive index, 118–119
 - transmission coefficient, 118, 119
 - longitudinal (compression) waves, 104, 105, 106, 109
 - critical angle interactions at boundaries, 119–120
 - near-field/far-field regions, 121
 - shear (transverse) waves, 104, 105, 109
 - surface waves, 104–105
 - Love, 104
 - Raleigh, 104–105
 - velocity, 105, 106, 109
 - adiabatic compressibility calculation, 151
 - gases, 106, 109
 - liquids, 109
 - solids, 109
 - wave characteristics, 103–105
 - Ultraviolet radiation, ozone combined sanitization procedures, 202–203
 - Uplandicine, 72
- V**
- Vanadium, inositol phosphate interactions, 36
 - Vegetables
 - chlorine treatment, 188
 - in-plant contamination, 194
 - inositol phosphates, 22–23
 - ozone treatment, 190, 191, 192, 194, 195–196
 - pesticide residues reduction, 199
 - sanitization methods, 170
 - texture, ultrasonic sensing, 142
 - Vibrio cholerae*, 180
 - Vibrio parahaemolyticus*, 180

Viscosity, ultrasonic measurement, 147–148, 149

von Kármán vortex street, 132

Vortex shedding, 132

W

Wheat

flour

calcium bioavailability, 35

iron absorption influence, 33–34

ozone treatment, 192

zinc absorption influence, 33

gluten, 220–221

polymorphisms, 221, 222

proteins, 221–224

glutenin, 219–290

inositol phosphates, 12, 13, 16, 20

degradation during digestion, 24

degradation during processing, 21

phytase, 11

storage proteins, 220

transformation system, 253–254

Whey protein, ultrasonic analysis, 152, 153

Wine, ultrasonic alcohol determination, 138

Winery equipment, ozone treatment, 192

Y

Yogurt, ultrasonic sol–gel transition monitoring, 147

Z

Zinc absorption, inositol phosphates influence, 32, 33, 35, 36, 42

Zygosaccharomyces bailii, 197