

INDEX

A

- Abees* sweet potato, dietary fiber
 - contents of, 13
- Acetyl CoA carboxylase (ACC-1), 90
- Acid/aspartyl proteases, 279
- Actinidia arguta* fruits, 294
- Actinidia deliciosa*, 293
- Actinidia eriantha*, 299, 302
- Actinidia* genus fruits
 - allergenic properties, 317
 - components, 299–316
 - actinidin, 310–316
 - calcium oxalate, 304–305
 - organic acids, 300–301
 - pigments, 305–310
 - sugar and sugar alcohol, 299–300
 - vitamin C, 301–303
 - health benefits and, 317–318
 - lutein and β -carotene, 309
 - perspectives, 318–319
 - species and cultivars, 295–299
 - A. arguta*, 297–299
 - A. chinensis*, 296–298
 - A. deliciosa*, 295–296, 298
 - A. eriantha*, 299
 - A. kolomikta*, 299
 - A. rufa*, 298–299
- Actinidia kolomikta*, 297
- Actinidin in *Actinidia* genus
 - fruits, 310–316
 - actinidin concentration and protease activity, 314
 - cultivars, differences among, 313–316
 - enzyme supplement and, 313
 - food proteins, effects on, 311–313
 - fruit growth and ripening, changes during, 311
 - protease from Shinzan and, 316
 - spatial distribution in fruit, 311
- Acute encephalitis, CDV infection and, 63
- Ad-2, 91
- Ad-5, 65, 68, 90–91
 - effect of infection on, 66
 - human obesity, association with, 66
 - induced obesity, 90–91
 - mechanism of, 91
 - mechanism of action, 67, 91
- Ad-31, 91
- Ad-36, 65, 68, 87–91
 - animal models, 88–89
 - effect of infection on, 65
 - human obesity, association with, 65, 89
 - induced obesity, 87–90
 - mechanism of, 89–90
 - mechanism of action, 67, 89–90
- Ad-37, 65, 67, 91
 - effect of infection on, 65
 - human obesity, association with, 65
 - mechanism of action, 67
- Adenoviruses, 65, 68, 84–90
 - SMAM-1, 65, 68, 85–87
 - type 5, 65, 68, 90–91
 - type 36, 65, 68, 87–90
 - type 37, 65, 90
- Adenovirus type 5, *see* Ad-5
- Adenovirus type 36, *see* Ad-36
- Adenovirus type 37, *see* Ad-37
- Adipogenic pathogens,
 - see also* individual pathogens
 - effects of, 64–66
 - mechanism of action, 67–68
- Aeromonas* sp., 176
- Albumin, 275
- Alexandrium tamarense*, 173–174

- Algae
 cultivation, 249–250
 fucans/fucanoids and other
 polysaccharides from, 271–272
 hydrocolloids, polysaccharides
 from, 269–271
 polysaccharides from, 240–241, 269–274
 transgenics, 250–252
 diatoms, 251
 dinoflagellates, 251–252
 green algae, 252–253
 macroalgae, 253
Alicyclobacillus acidoterrestris, 128
 Allergenic properties, of *Actinidia*
 genus fruits, 317
Alteromonas, 272
Alteromonas tetraodonis, 174, 181
Ambystoma tigrinum, 159
Amphidinium, 252
Amphidinium carteri, 262
Amukeke chips, 37–38
 α -Amylase influence, SPS syrup and, 26–27
 Annelids
 tetrodotoxin in, 172–173
 toxicity of, 173
 Anthocyanins, 2–3
 in *Actinidia* genus fruits, 308–310
 health benefits associated with, 16
 refrigerated fruit juices, contents of, 119–120
 in sweet potato, 14–16
 Antioxidant activity, in refrigerated fruit
 juices, 120–122
Aphanizomenon, 247
Aphanizomenon flos-aquae, 261
 Apple juice, *see also* Refrigerated fruit juices
 consumption of, 104
 Aromatic profile, of refrigerated fruit
 juices, 113–114
Arothron firmamentum, 158
Arothron hispidus, 158
Arothron manilensis, 158
Arothron mappa, 158
Arothron nigropunctatus, 158
Arothron reticularis, 158
Arothron sp., 152
Arothron stellatus, 158
 Arrowworms, tetrodotoxin in, 173
Arthrospira, 247
Aspergillus fischeri, 131
 Astaxanthin, in marine-derived food
 ingredients, 239, 258–259
Astropecten latespinosus, 170, 178
Astropecten polyacanthus, 169, 170, 174, 178, 181
Astropecten scoparius, 170, 178
Astropecten vappa, 170
Atelopus chiriquiensis, 161
Atelopus oxyrhynchus, 161, 176
Atelopus varius ambulatorius, 161
Atelopus varius varius, 160
Atelopus zeteki, 162
Atergatis floridus, 163–166, 174, 181
Atergatisopsis germaini, 163–165, 167
 Autoimmune thyroiditis, RAV-7 and, 76–77
 Avian leukosis viruses, 73, 75
 Avian retroviruses viruses, 73
Ayamurasaki sweet potato, antioxidative and
 radical scavenging activities, 15
- B**
- Baby kiwi fruit, *see Actinidia* genus fruits
Babylonia formosae, 169
Babylonia japonica, 168
Bacillus sp., 174, 281, 283
 Bacteria, tetrodotoxin in, 174–176
Bacteroides thetaiotaomicron, 83–84
 Batata, *see* Sweet potato
 BDV, *see Borna disease virus* (BDV)
 Beauregard sweet potato
 protein contents of, 9
 starch composition and, 23–24
Beauveria bassiana, 256
Beniazuma sweet potato, protein contents of, 9
Betaphycus gelatinum, 270
 Beverage, from sweet potato, 40
Biddulphia sinensis, 262
Birgus latro, 166
 Blue-ringed octopus, tetrodotoxin in, 165, 168
 Body mass index (BMI) in humans
 Ad-36 and, 89
 CP and, 77–78
 SMAM-1 and, 86
 Boniato, *see* Sweet potato
Borna disease virus (BDV), 64, 67, 81–83, 92
 effect of infection on, 64
 human obesity, association with, 64
 induced obesity, 64, 67, 81–83
 mechanism of, 82–83
 mechanism of action, 67, 82–83

Brachycephalus ephippium, 161

Bread supplemented with SPF

development and storage stability of, 33–34

dough enhancers, 33–34

macroscopic and sensory evaluation of, 33

Bryum, 268

Byssoschlamys fulva, 131

Byssoschlamys nivea, 131

C

Calcium oxalate, in *Actinidia* genus

fruits, 304–305

Camote, *see* Sweet potato

Candida lambica, 129

Candida pelliculosa, 127

Candida sake, 129

Canine distemper virus (CDV),

63–64, 67, 69–73, 92

acute encephalitis, 63

alters hypothalamic integrity, 71

cytokine production, 72–73

down regulates

long leptin receptor and increases leptin, 71

melanin-concentrating hormone, 71–72

effect of infection on, 64

F gene mRNA levels, 70

of genus *Morbillivirus*, 63

hit-and-run effect, 62, 70, 72

human obesity, association with, 64

induced obesity, 63, 69

mechanism of, 70–73

mechanism of action, 67, 70–73

neuropeptides and, 72

nucleoprotein transcripts, 70

obesity impact, 69

on brains and reproductive organs, 69

on catecholamine levels, 69, 72

on insulin levels, 69

on leptin levels, 69, 71

on lipid and fat cells, 69

on lipogenesis, 69

on tyrosine hydroxylase, 69

replication in rain, 70

Canthigaster ribulata, 157

Capillary isotachopheresis, detection

method for TTX, 185, 189–190

Carcinoscorpius rotundicauda, 152, 162, 166, 181

β -Carotene, 2–3

in marine-derived food ingredients, 239,

257–258

in sweet potato, 10–13, 16, 31

Carotenoids

in *actinidia* genus fruits, 307–309

in marine-derived food ingredients, 239,
257–261

astaxanthin, 239, 258–259

β -carotene, 239, 257–258

refrigerated fruit juices, contents of, 117–119

Catecholamine levels, CDV obesity

impact on, 69, 72

CDV, *see* *Canine distemper virus* (CDV)

CEBP α gene, 90

CEBP β gene, 90

Cephalothrix linearis, 172

Cephalothrix sp., 172

Cerebratulus lacteus, 172

Charonia sauliae, 148, 168–139, 177

poisoning due to digestive gland of, 147–148

Chelonodon patoca, 158, 179, 219–220

Chick Embryo Lethal Orphan virus

(CELO), 85, 91

China, puffer poisoning cases in, 148–152

Chinese potato, *see* Sweet potato

Chitin and chitosan, polysaccharides

from, 273–274

Chitinolytic enzymes, 281

Chlamydia pneumoniae (CP), 66, 68, 77–78

coronary artery stenosis and, 78

coronary heart disease (CHD) and, 77

effect of infection on, 66

higher BMI and, 77–78

human obesity, association with, 66

mechanism of action, 67, 78

Chlamydia psittaci, 78

Chlamydia trachomatis, 78

Chlamydomonas reinhardtii, 250, 252

Chlorella minutissima, 263, 267

Chlorella protothecoides, 257, 259

Chlorella sp., 245–248, 250, 252

Chlorella vulgaris, 257

Chlorophyll

in *actinidia* genus fruits, 305–307

in marine-derived food ingredients,
239, 260–261

Chymotrypsins, 279

Cilera abana, *see* Sweet potato

Coarse *inginyo*, 37

CODEX, General Standard for

Fruit Juices and Nectars, 104

Coenobitidae, 166

Collagen, 274
 Collagenases, 279–280
 Color, of refrigerated fruit juices, 122–124
Colostethus inguinalis, 161
 CP, *see* *Chlamydia pneumoniae* (CP)
Crypthecodinium, 250
Crypthecodinium cohnii, 245, 262–263
Cryptotethya crypta, 254
Ctenidium, 268
 Cyanobacteria, exopolysaccharides from, 272
Cyanospira capsulata, 272
Cyanothece, 272
Cyclotella, 250
Cynops ensicauda, 159, 180
Cynops pyrrhogaster, 159, 180, 220
 Cytokine production, and CDV, 72–73
 Cytomegalovirus, 77
 Cytotoxicity test, detection method for
 TTX, 185, 195–196

D

Demania alcali, 167
Demania reynaudi, 163–165, 167
Demania toxica, 167
 Diatoms algae transgenics, 251
 Dietary fiber, in sweet potato, 13–14, 31
 Digestive proteases, 278–280
 Dinoflagellates
 algae transgenics, 251–252
 tetrodotoxin in, 173–174
Diodon hystrix, 152
Dunaliella bardawil, 253
Dunaliella salina, 247, 250, 252, 257–258
Dunaliella sp., 245–246, 248–249,
 252–253, 258, 276
Dysidea avara, 255

E

Ectocarpus siliculosus, 253
 Elastase, 279–280
 Electrophoresis, detection method
 for TTX, 185, 189
 Electrospray ionization-time of
 flight/mass (ESI-TOF/MS), 161–162
 TTX, detection method for, 185, 193–194
Endomycopsis fibuligera, 36
Entomophthora obscura, 266
 Enzyme-linked immunoassay (EIA), 197
 Enzyme-linked immunosorbent
 assay (ELISA) system, 196

Enzymes

chitinolytic, 281
 digestive proteases, 278–280
 extremophilic, 282–283
 food industry and, 277–283
 lipases, 280–281
 marine-derived food ingredients,
 242–243, 276–283
 polyphenol oxidases, 281
 from red algae, 283
 transglutaminase, 281–282
Eriphia scabricula, 167
Eriphia sebana, 163, 167
Escherichia coli, in refrigerated fruit
 juices, 130–131
 ESI-TOF/MS, *see* Electrospray ionization-time
 of flight/mass (ESI-TOF/MS)
 Ethanol, SPF fermentation to, 36
Eucheuma denticulatum, 270
Eukrohnia hamata, 173
Eupenicillium, 131
Euphausia superba, 268
 Exopolysaccharides
 cyanobacteria and, 272
 extremophiles and, 272
 Extremophiles
 marine-derived food ingredients
 sources, 253–254
 polysaccharides from, 272
 sources of LC-PUFAs, 267
 Extremophilic enzymes, 243, 282–283

F

FABMS, *see* Fast atom bombardment mass
 spectrometry (FABMS)
 Fast atom bombardment mass spectrometry
 (FABMS), 173–174
 TTX, detection method for, 185, 192–193
 Fat cells, CDV obesity impact on lipids and, 69
 Fatty acids contents, of refrigerated
 fruit juices, 112–113
 Fatty acid synthetase (FAS), 90
 F gene mRNA levels, and CDV, 70
 Fiber one cereal, 34–36
 Fine patterned puffer, poisoning
 due to liver of, 147–148
 Fish
 LC-PUFAs and, 264–266
 marine-derived food ingredients
 and, 256–257

Flaccisagitta enflata, 173
Flaccisagitta scirpassae, 173
Flatworms
 tetrodotoxin in, 170–171
 toxicity of eggs laid by, 171
Flavonoids contents, of refrigerated fruit juices, 119–120
Flavor, of refrigerated fruit juices, 113–114
Food industry, enzymes and, 277–283
Food ingredients production, marine biotechnology for, 237–284
Food Standards Agency, UK, 108
Free amino acids contents, of refrigerated fruit juices, 112–113
French-fries, from sweet potato, 41–42
Frogs, tetrodotoxin in, 160–162
Fruit and vegetables, *see also* Refrigerated fruit juices
 consumption of, 105
 epidemiological studies, 105–106
 protective effect of, 105–106
Fungi, sources of LC-PUFAs, 266
Furan formation, in refrigerated fruit juices, 111–112
Fusarium, 129

G

G. crassissima, 271
Gas chromatography-mass spectrometry (GC-MS), 111
 detection method for TTX, 185, 191
Gastropods
 poisoning due to digestive gland of, 147–148
 tetrodotoxin in, 168–169
 toxicity and toxin composition of, 169
GC-MS, *see* Gas chromatography-mass spectrometry (GC-MS)
Gelatin, 274–275
Gelidium, 270
Geodia cydonium, 255
Geotrichum sp., 129
Gigartina, 270
“Giza 69”, 13
GLUT-4, 90
Gobies
 intoxication in uremic patient in Taiwan, 151–152
 local variation of toxicity in, 161
 tetrodotoxin (TTX) in, 159–161

GPDH gene, 90
Gracilaria, 245, 270–271
Grapefruit juice, *see also* Refrigerated fruit juices
 consumption of, 104
Grape kiwi fruit, *see Actinidia* genus fruits
Grapsus albolineatus, 167
Grateloupia, 270
Green algae transgenics, 252–253
Guava juice, *see also* Refrigerated fruit juices
 consumption of, 104
Gut microbiota, 66–67, 83–84
 effect of infection on, 66
 human obesity, association with, 66
 induced obesity, 83–84
 mechanism, 84
 mechanism of action, 67, 84
Gymnodinium simplex, 262
Gyrodinium cohnii, 262

H

Haematococcus pluvialis, 247, 249, 252, 257, 259
Haematococcus sp., 252, 259
Halobacterium mediterranei, 272
Halosydna brevisetosa, 173
Haplochaena maculosa, 180
Hard red spring wheat (HRSW), 33
Harmothoe imbricata, 173
Hayward fruit, *see Actinidia* genus fruits
Hazard Analysis and Critical Control Points (HACCP) system, 106
Health benefits, of *Actinidia* genus fruits, 317–318
Helicobacter pylori, 77–78
Hemigrapsus sanguineus, 200
Hermenia acanthopeltis, 173
High-performance liquid chromatography (HPLC), 151
 operating conditions for analysis of newt toxins, 188
 TTX, detection method for, 185–188
HIPEF-treated juice, 118–119, 124
Hit-and-run effect of CDV, 70, 72, 92
Hizikia fusiforme, 246
¹H NMR spectrometry,
 see NMR spectroscopy
Hort16A fruit, *see Actinidia* genus fruits
HPLC, *see* High-performance liquid chromatography (HPLC)
Human adenoviruses, 85

Human food systems

- SPF production and utilization
 - for, 28–38, 30, 34
- sweet potato starch utilization in, 18–29 (*see also* SPS syrup)
- sweet potato utilization as value-added product in, 17–18

Human Genome Project, 212

Human obesity association with

- ad-5, 66
- ad-36, 65
- ad-37, 65
- BDV, 64
- CDV, 64
- CP, 66
- gut microbiota, 66
- infections in, 92–93
- inflammation and, 92–93
- macrophage colony-stimulating factor (MCSF), 93
- RAV-7, 64
- scrapie agent, 65
- SMAM-1, 65

Hydrocolloids, polysaccharides from, 269–271

Hydroponic sweet potato starch (HSPS) syrup, 27–28

Hyperglycemia, in ME7-infected mice, 80

Hyperlipidemia, RAV-7 induced, 74, 77

Hypnea, 270

Hypothalami, of CDV infected rats, 70–71

Hypothalamic integrity, and CDV, 71

I

IL- β transcripts, 73

IL-6 transcripts, 73

Immunoassay, detection method for TTX, 185, 196–197

Infatobesity concert, 62

Insulin levels, CDV obesity impact on, 69, 90

IR spectrometry, detection method for TTX, 185, 191–192

Isochrysis galbana, 264

J

Jania sp., 173

Japan, puffer poisoning cases in, 147–148

Japanese potato, *see* Sweet potato

J6/66 sweet potato

- composition of starch processed from, 23–24
- protein contents of, 9

K

Kappaphycus alvarezii, 270*Kara-imo*, *see* Sweet potato

Kenya, sweet potato processing and utilization in, 44–47

Kiwi berry fruit, *see* *Actinidia* genus fruitsKiwifruit, *see* *Actinidia* genus fruits*Kloeckera apis*, 127, 129*Koganesengan* sweet potato, protein contents of, 9*Koinostylochus* sp., 171

Krill, sources of LC-PUFAs, 268

Kumar, *see* Sweet potatoKumara, *see* Sweet potato*Kunuzaki*, from sweet potato, 40

L

Lactobacillus brevis, in refrigerated fruit juices, 129*Lactobacillus plantarum*, 274

in refrigerated fruit juices, 129–130

Lagocephalus gloveri, 157, 213, 216*Lagocephalus inermis*, 157–158, 216*Lagocephalus lunaris*, 157–158, 179, 213, 216*Lagocephalus sceleratus*, 157, 216*Lagocephalus wheeleri*, 157, 216*Laminaria*, 245–246, 253*Laminaria japonica*, 245, 246, 270–271*Laminaria saccharina*, 246LC-MS, *see* Liquid chromatography-mass spectrometry (LC-MS)*Lepidonotus helotypus*, 173

Leptin levels

CDV obesity impact on, 69, 71

and leptin receptor, CDV down regulates, 71

Leptodius sanguineus, 167*Lineus fuscoviridis*, 172

Lipases, 280–281

Lipids

CDV obesity impact on fat cells and, 69

marine-based long-chain PUFAs

(LC-PUFAs), 239, 261–268

extremophiles as source of, 267

fish as source of, 264–266

fungi as source of, 266

krill as source of, 268

macroalgae and mosses as source of, 267–268

microalgae as source of, 262–264

transgenic organisms as source of, 266–267
 marine-derived food ingredients,
 239, 261–269
 sterols, 268–269
 Lipogenesis, CDV obesity and, 69
 Liquid chromatography-mass
 spectrometry (LC-MS), 310
 detection method for TTX, 185, 192–194
Lophozozymus pictor, 163–165, 167
 Lymphotropic RNA virus, 63

M

Macroalgae
 algae transgenics, 253
 and mosses, sources of LC-PUFAs, 267–268
 Macrophage colony-stimulating
 factor (MCSF), in human obesity, 93
Mandazis, from sweet potato, 46–47
 Mango juice, *see also* Refrigerated fruit juices
 consumption of, 104
Marchantia, 268
 Marine-based long-chain PUFAs (LC-PUFAs)
 lipids, 239, 261–268
 production technologies, 262–264
 sources of
 extremophiles as source of, 267
 fish, 264–266
 fungi, 266
 krill, 268
 macroalgae and mosses, 267–268
 microalgae, 262–264
 transgenic organisms, 266–267
 Marine biotechnology, for food ingredients
 production, 237–284
 Marine-derived food ingredients
 categories of, 239–243, 257–283
 enzymes, 242–243, 276–283
 lipids, 239, 261–269
 photosynthetic pigments, 239, 257–261
 polysaccharides, 240–241, 269–274
 proteins, 241, 274–276
 sources of, 244–257
 algae cultivation, 249–250
 algae transgenics, 250–252
 extremophiles, 253–254
 fish and seafood byproducts, 256–257
 macroalgae, 245–246
 macro- and microalgae, 244–253
 marine sponges, 254–256
 microalgae, 246–253

Marine enzymes
 food industry and, 277–283
 chitinolytic enzymes, 281
 digestive proteases, 278–280
 enzymes from red algae, 283
 extremophilic enzymes, 282–283
 lipases, 280–281
 polyphenol oxidases, 281
 transglutaminase, 281–282
 sources of, 276–277
 Marine organisms, 238
 Marine sponges, sources of marine-derived
 food ingredients, 254–256
Matobolwa, from sweet potato, 38
Matteuccia, 268
 Measles virus (MV), 63
 Melanin-concentrating hormone (MCH), 69
 CDV down regulation, 71–72
Michembe from sweet potato, 38
 Microalgae, sources of LC-PUFAs, 262–264
 Mineral contents, in sweet potato, 13
Mnium, 268
Morbillivirus, CDV of genus, 63
Mortierella alpina, 266–267
Mortierella elongata, 266
Mortierella isabellina, 266
Mortierella ramanniana, 266
 Mouse bioassay, detection method for
 TTX, 184–186, 196

N

Nassarius glans, 149
 Nata de coco dessert, 17
Natica alapailionis, 169
Natica lineata, 168–169, 176, 180, 219
Natica pseustes, 170
Natica vitellus, 149, 168–169, 219
Navicula incerta, 262
Navicula pelliculosa, 262
Neosartorya fischeri, in refrigerated
 fruit juices, 131
Neoxanthias impressus, 167
 Neuropeptides expression and CDV, 72
 Neurotropic negative-stranded RNA virus, 63
 Newts
 local and sexual variation of toxicity in, 160
 tetrodotoxin (TTX) in, 159–160
 Nigeria, traditional utilization of
 sweet potatoes in, 6
Niotha clathrata, 149, 168–169, 176, 219

Nitzschia, 250
Nitzschia frustulum, 262
 N-3 LC-PUFAs, *see* Marine-based long-chain PUFAs (LC-PUFAs)
 NMR spectroscopy, detection method for
 TTX, 185, 194–195
 Nonenzymatic browning evaluation, in
 refrigerated fruit juices, 111–112
Nostoc, 247, 272
Nostoc commune, 247
Nostoc flagelliforme, 247
Notocoplana roreana, 171
Notophthalmus viridescens, 159, 180
Notoplana humilis, 171
 Nucleoprotein transcripts, of CDV, 70
 Nutrients, in refrigerated fruit juices, 105

O

O. maculosus, 174
Oliva hirasei, 146, 149, 169, 219
Oliva miniacea, 149, 169, 219
Oliva mustelina, 149, 169, 219
 Omega-3 LC-PUFAs, *see* Marine-based long-chain PUFAs (LC-PUFAs)
 On-the-go food, 17
 Orange-fleshed sweet potatoes, source of
 provitamin A, 10
 Orange juice, *see also* Refrigerated fruit juices
 consumption of, 104, 106
 Organic acids, in *Actinidia* genus
 fruits, 300–301
 Osteopetrosis, 73

P

Prachaeturichtys palynena, 160
 Paralytic shellfish poisons (PSPs), 152
Paramesotriton hongkongensis, 159
 Paramyxoviruses, 63
Parasagitta elegans, 173
 Passion fruit juice, *see also*
 Refrigerated fruit juices
 consumption of, 104
 Patents, on sweet potato products, 40–41
 Pectinesterases activity, of refrigerated
 fruit juices, 124–126
Penicillium, 129
Phaeodactylum tricornutum, 250–251,
 251, 262–264, 266–267
 Photosynthetic pigments
 carotenoids, 239, 257–261

 astaxanthin, 239, 258–259
 β -carotene, 239, 257–258
 chlorophylls, 239, 260–261
 marine-derived food ingredients, 239,
 257–261

Physcomitrella patens, 267
Phytophthora infestans, 266
 Pigments in *Actinidia* genus fruits, 305–310
 anthocyanins, 308–310
 carotenoids, 307–309
 chlorophyll, 305–307
Pilumnus vespertilio, 167
 Pineapple juice, *see also*
 Refrigerated fruit juices
 consumption of, 104
Planocera multitentaculata, 170–171, 170–172
Planocera reticulata, 170–171
Platypodia granulosa, 167
Plsiomonas sp., 176
Pogonatum sp., 268
Pogonatum urnigerum, 268
Polinices didyma, 168, 219
Polinices tumidus, 168
Polypedates sp., 161
 Polyphenol oxidase, 281
 refrigerated fruit juices, inactivation
 in, 126–127
 Polysaccharides from
 algae, 240–241, 269–274
 fucans/fucanoids and other
 polysaccharides from, 271–272
 hydrocolloids, 269–271
 chitin and chitosan, 273–274
 cyanobacteria, 272
 extremophiles, 272
 marine-derived food ingredients,
 240–241, 269–274
Polytrichum, 268
Pomacea canaliculata, 219
Porphyra sp., 245, 246, 253
Porphyridium cruentum, 263–264
 PPAR- γ gene, 90–91
 Pro-opiomelanocortin (POMC) cell bodies,
 loss of, 71
 Protamine, 275–276
 Proteases, 278–280
 Proteins
 marine-derived food ingredients
 from, 241, 274–276
 in sweet potato leaves and roots, 8–10, 31

- Provitamin A, orange-fleshed
sweet potatoes and, 10
- Pseudoalteromonas*, 272
- Pseudomonas* sp., 176
- Pseudopotamilla ocellata*, 172, 174
- Psychrophilic enzymes, 282
- Puffer
edible part of, in Japan, 147
food industry programs for, 208–220
new developed biotechnology
and, 208–209
traditional food puffer liver
kimo, revival of, 209
nontoxic production, 208–209
poisoning cases in
Japan, 147–148
Taiwan and China, 148–152
poisoning deaths, 142
species, 156–158
direct sequence analysis approach, 210
genome techniques, identification
by, 209–213
PCR-RFLP technique, 210–211
PCR-SSCP technique, 211–212
by protein technique, 213–218
RAPD-PCR technique, 212
tetrodotoxin in, 156–158
toxicity of, 147–148, 156
Taiwanese puffer, 157
toxin chemistry, 197–198
traditional food puffer liver kimo, 209
TTX infestation to nontoxic cultured, 200
- R**
- Rain, CDV replication in, 70
- Rapana rapiformis*, 168–169
- Rapana venosa venosa*, 168–169
- RAV-7, 64, 67, 73–77
autoimmune thyroiditis, 76–77
effect of infection on, 64
human obesity, association with, 64
induced hyperlipidemia, 74, 77
induced obesity, 74–75
specificity of, 75–76
mechanism of induced changes, 67, 76–77
stunting and, 74
- Ready-to-eat breakfast cereals (RTEBC), 34–36
- Ready-to-eat sweet potato breakfast cereal
physical properties and sixth graders'
acceptance of extruded, 35–36
- sensory characterization by descriptive
analysis of, 34–35
- Red algae, enzymes from, 283
- Red calcareous alga, tetrodotoxin in, 173
- Redigobius caninus*, 160
- Refrigerated fruit juices
anthocyanins contents, 119–120
antioxidant activity, 120–122
aromatic profile of, 113–114
carotenoids contents, 117–119
color, 122–124
consumer preferences for, 104
consumption of, 104
E. coli, 130–131
fatty acids contents, 112–113
flavonoids contents, 119–120
flavor of, 113–114
free amino acids contents, 112–113
furan formation in, 111–112
inclusion in healthy diet, 104
Lactobacillus brevis, 129
Lactobacillus plantarum, 129–130
market for, 104
Neosartorya fischeri, 131
nonenzymatic browning evaluation, 111–112
nutrients in, 105
pectinesterases activity, 124–126
physicochemicals and quality
characteristics of, 108–111
polyphenol oxidase inactivation, 126–127
processing, 107–108
quality and safety issues, 104–131
quality parameters, 106–107
Salmonella enteritidis, 131
Staphylococcus aureus, 131
vitamin A contents, 117–119
vitamin C concentration in, 114–117
yeast, 127–129
- Renibacterium salmoninarum*, 252
- Rhodotorula rubra*, 129
- Ribbon worms, tetrodotoxin in, 172
- Rous-associated virus-7 (RAV-7), *see* RAV-7
- S**
- Saccharomyces cerevisiae*, 128, 267
- Salmonella enteritidis*, in refrigerated
fruit juices, 131
- Sargassum confusum*, 270
- Satamu sweet potato, 6
- Satsuma bairdi*, 219

- Satsuma-imo, *see* Sweet potato
- Scenedesmus*, 246
- Schizochytrium*, 263–264
- Scrapie agent
- effect of infection on, 65
 - human obesity, association with, 65
 - induced obesity, 65, 67, 78–81
 - mechanism of, 80–81
 - mechanism of action, 67
- Seafood byproducts, 256–257
- Shewanella alga*, 174
- Shewanella putrefaciens*, 174, 181
- Sillago japonica*, 160
- SMAM-1, 65, 68, 85–87, 92
- effect of infection on, 65
 - human obesity, association with, 65
 - induced obesity, 85–87
 - mechanism of, 87
 - mechanism of action, 67, 87
- Snail poisoning outbreaks and patients, in Zhoushan City, 150
- Sodium channels
- mechanism of action on, 201–202
 - molecular structure of, 202–203
- TTX-R, 204–206
- Soursop juice, *see also* Refrigerated fruit juices
- consumption of, 104
- Space missions, sweet potato vegetarian products intended for, 38–40
- Spadella angulata*, 173
- Species and cultivars, of *Actinidia* genus
- fruits, 295–299
 - A. arguta*, 297–299
 - A. chinensis*, 296–298
 - A. deliciosa*, 295–296, 298
 - A. eriantha*, 299
 - A. kolomikta*, 299
 - A. rufa*, 298–299
- Species identification for puffer
- by genome techniques, 209–213
 - direct sequence analysis approach, 210
 - PCR-RFLP technique, 210–211
 - PCR-SSCP technique, 211–212
 - puffer genome technique, 212–213
 - RAPD-PCR technique, 212
 - by protein technique, 213–218
- SPF, *see* Sweet potato flour
- SPF/whole-wheat bran (SPFWWB)
- cereal, 34–36
- Sphagnum*, 268
- Sphoeroides pachygaster*, 216
- Spirulina (Arthrospira) maxima*, 248, 257
- Spirulina (Arthrospira) platensis*, 248, 250, 261
- Spirulina* sp., 245–248, 261
- Sporamins A and B proteins, in sweet potato, 9
- SPS, *see* Sweet potato starch (SPS)
- SPS syrup
- α -amylase influence, 26–27
 - enzymatic hydrolysis into glucose syrup, 25
 - pH effects and concentration times on
 - functional properties of, 26
 - physicochemical and viscometric properties of, 25–26, 29
 - sensory and consumer evaluation of, 27–28
- Staphylococcus aureus*, in refrigerated fruit juices, 131
- Starfish, tetrodotoxin in, 169–170
- Sterols, marine-derived food ingredients in, 268–269
- Stunting, and RAV-7, 74
- Stylochoplana clara*, 171
- Stylochus ijimai*, 171
- Stylochus orientalis*, 171
- Suberites domuncula*, 255
- Sugar and sugar alcohol, in *Actinidia* genus fruits, 299–300
- Suioh* sweet potato greens, nutritional contents of, 8–9
- Sweet potato, *see also* SPF (sweet potato flour)
- anthocyanins, 14–16
 - antioxidative and radical-scavenging activities, 14–15
 - beverage, 40
 - biochemical composition of, 7–16
 - β -carotene in, 10–13, 16
 - dietary fiber, 13–14
 - French-fry-type product from, 41
 - growing season, 5
 - health benefit from antihypertensive and antidiabetic properties of, 15–16
 - in human diets, 6–7
 - kunuzaki, 40
 - leaves and roots, 5
 - mineral contents, 13
 - nutritional composition of, 7–16
 - origin of, 4
 - patents regarding products, 40–41
 - per capita consumption of, 17
 - potential in Ugandan food system, 42–44
 - potential products, 38–42

- processing and utilization in Kenya, 44–47
 - production of, 7–8
 - products intended for space missions, 38–40
 - protein
 - in leaves, 8–9
 - in roots, 9–10
 - scientific classification, 4
 - sporamins A and B proteins in, 9
 - as staple food, 2, 13
 - starch
 - bulk ingredients from cultivars of, 23–24
 - composition and properties of, 23–24
 - effects of processing technology on yield and quality, 23, 25
 - utilization in human food systems, 18–29
 - starch syrup
 - influence of α -amylase on physical properties and consumer acceptability of, 26–27
 - pH effects and concentration times on functional properties of, 26
 - physicochemical and viscometric properties of, 25–26, 29
 - sensory and consumer evaluation of, 27–28
 - starch utilization [*see* Sweet potato starch (SPS)]
 - textural measurements and product quality of restructured French-fries from, 41–42
 - utilization as value-added products, 17–18
 - varieties, 11–12
 - vegetarian products intended for space missions, 38–40
 - vitamin A deficiency (VAD) and, 10–11
 - yam and, 4–5
 - Sweet potato flour (SPF)
 - breadmaking properties of, 32–33
 - bread supplemented with
 - development and storage stability of, 33–34
 - dough enhancers, 33–34
 - macroscopic and sensory evaluation of, 33–34
 - fermentation to ethanol, 36
 - genetic variation in color of, 36–37
 - like products, 37–38
 - preparation of, 36
 - processed in agroecological sites, and small-scale processing technologies, 37
 - production and utilization for human food systems, 28–38
 - protein content of, 30–31
 - quality characteristics of, 30–31
 - quality evaluation of, 37
 - ready-to-eat sweet potato breakfast cereal, 34–36
 - use as coloring materials, 14
 - use in wheat-based composite flour products, 36–37
 - Sweet potato leaves and roots, 5
 - proteins in, 8–10
 - Sweet potato starch (SPS)
 - bulk ingredients from cultivars of, 23–24
 - composition and properties of, 23–24
 - effects of processing technology on yield and quality, 23, 25
 - production procedures for, 21–22
 - scanning electron micrographs of granules of, 19
 - syrup
 - α -amylase influence, 26–27
 - pH effects and concentration times on functional properties of, 26
 - physicochemical and viscometric properties of, 25–26, 29
 - sensory and consumer evaluation of, 27–28
 - utilization in human food systems, 18–29
 - Symbiodinium*, 252
- T**
- Tachypleus gigas*, 162
 - Taiwan
 - goby intoxication in uremic patient in, 151–152
 - poisoning incident due to ingestion of unknown fish in, 150
 - puffer poisoning cases in, 148–152
 - Takifugu chinensis*, 157
 - Takifugu chrysops*, 156
 - Takifugu exascurus*, 157
 - Takifugu flavidus*, 156–157
 - Takifugu niphobles*, 150, 157–158, 174
 - Takifugu oblongus*, 152, 157, 191, 216
 - Takifugu obscurus*, 156–157
 - Takifugu pardalis*, 156, 158, 179
 - Takifugu poecilonotus*, 156, 158
 - poisoning due to liver of, 147–148
 - Takifugu porphyreus*, 156

- Takifugu pseudommus*, 157, 212
Takifugu rubripes, 156, 171, 212–213
Takifugu snyderi, 156, 174, 181
Takifugu stictonotus, 157
Takifugu vermicularis, 157, 219–220
Takifugu xanthopterus, 157, 216
Talaromyces flavus, 131
Taricha granulosa, 159, 180
Taricha oregon, 159
Taricha rivularis, 180
Taricha torosa, 159, 180
Tetraodon albobreticulatus, 157–158
Tetraodon fangi, 158
Tetraodon nigroviridis, 157–158
Tetraodon ocellatus, 157
Tetraodon steindachneri, 158, 219–220
Tetraodontiformes, *see* Tetrodotoxin (TTX)
Tetraselmis, 250
Tetradotoxification symptoms, 152–154
Tetrodotoxin (TTX)
 4,9-anhydro type, 145
 bearing organisms, 156–176
 annelids, 172–173
 arrowworms, 173
 bacteria, 174–176
 blue-ringed octopus, 165, 168
 dinoflagellate, 173–174
 flatworms, 170–171
 frogs, 160–162
 gastropods, 168–169
 gobies, 159–161
 horseshoe crab, 162–164
 newts, 159–160
 puffer, 156–158
 red calcareous algae, 173
 ribbon worms, 172
 starfish, 169–170
 toxin accumulation from TTX-producing
 bacteria, 182–183
 TTX resistibility in, 199
 xanthid crabs, 163–167
 causative agent, 156–208
 chemical structure, 143–145
 detection methods for, 184–197
 bioassay, 185–186
 capillary isotachopheresis, 185, 189–190
 cytotoxicity test, 185, 195–196
 electrophoresis, 185, 189
 ESI-TOF/MS, 185, 193–194
 FABMS, 185, 192–193
 GC-MS, 185, 191
 HPLC, 185–188
 immunoassay, 185, 196–197
 IR spectrometry, 185, 191–192
 LC-MS, 185, 192–194
 mouse bioassay, 184–186, 196
 NMR spectroscopy, 185, 194–195
 TLC, 185, 189
 UV spectroscopy, 185, 190
 distribution in animals, 146
 elaborator, 176–178
 hemilactal type, 144
 hibernation agent, as attractant and, 219–220
 infection to animals, mechanisms of, 178–184
 infestation to nontoxic cultured puffer
 fish, 200
 lacton type, 144
 mechanisms of accumulation in marine
 animals, 179
 pharmacology of, 198–207
 mechanism of action on sodium
 channels, 201–202
 molecular structure of sodium
 channels, 202–203
 site of action and binding of
 TTX, 206–207
 TTX-R sodium channels, 204–206
 poisoning, 145–156
 due to ingestion of puffer, 142
 due to liver of finepatterned
 puffer, 147–148
 incidents in Japan, 143–148
 incidents in Taiwan and China, 148–152
 incidents in world, 143–152
 prevention, 155–156
 symptoms and signs, 152–154
 treatment, 154–155
 site of action and binding of, 206–207
 tautomers, 143
 therapeutic applications, 207–208
Thalamita sp., 167
TH cell bodies, loss of, 69, 71
Thermococcus chitonophagus, 281
Thin-layer chromatography (TLC), detection
 method for TTX, 185, 189
Thraustochytrium, 263–264
Tissue culture bioassay (TCBA), 195
TLC, *see* Thin-layer chromatography (TLC)
3T3-L1 preadipocytes, 90
TNF- α transcripts, 73

Transgenic organisms, sources of
LC-PUFAs, 266–267

Transglutaminase, 281–282

Triturus alpestris, 159, 180

Triturus cristatus, 180

Triturus marmoratus, 180

Triturus vulgaris, 159, 180

Tropical fruit juices, consumption of, 104

Trypsin, 279

TTX, *see* Tetrodotoxin (TTX)

Tubulanus punctatus, 172

TU-82-155 sweet potato, composition of
starch processed from, 23–24

Tutufa lissostoma, 168

Tylototriton andersoni, 159

Tyrosine hydroxylase (TH), *see* TH cell bodies

U

Ubhatata, *see* Sweet potato

Ugandan food system, sweet potato potential
in, 42–44

Ulva pertusa, 272

Umborium suturale, 170

Undaria pinnatifida, 246, 260

V

Vibrio alginolyticus, 174, 176

Vibrio fischeri, 181

Vibrio parahaemolyticus, 176

Vibrio sp., 174, 181, 272

Vitamin A contents, of refrigerated fruit
juices, 117–119

Vitamin A deficiency (VAD),
sweet potato role in fight
against, 10–11, 47–48

Vitamin C

in *actinidia* genus fruits, 301–303
concentration in refrigerated fruit
juices, 114–117

Volvox carteri, 250

W

White-fleshed sweet potato roots, 5

Whole-wheat bran (WWB) cereal, 34–36

X

Xanthias lividus, 164–165

Xanthid crabs

tetrodotoxin (TTX) in, 163–167

toxicity and toxin composition of, 165,
167–168

Xiphosuridae, 166

“Xushu 18” sweet potato, 24

Y

Yam

β-carotene content, 4–5

growing season, 5

sweet potato and, 4–5

Yeast, in refrigerated fruit juices, 127–129

Yongeichthys criniger, 159, 178

Yongeichthys nebulosus, 151, 160

Z

Zanthin, 259

ZESPRI™ GOLD kiwifruit,
see *Actinidia* genus fruits

Zeuxis castus-like, 169

Zeuxis samiplicatus, 149, 169

Zeuxis scalaris, 149, 168–169

Zeuxis sufflatus, 168, 219

Zosimus aeneus, 163–166