

INDEX

A

- Acute toxoplasmosis, 6
- Alaska pink salmon, 139
- Alaska pollock gelatin, 138–139
- Animal feed
 - C. difficile*, 60
 - E. coli* O157:H7, pathogen transmission, 72–73
 - rat feeding experiments, 38
- Asthenia, 7
- Atlantic cod, 138
- Atlantic salmon, 138

B

- Bacteriophage
 - high-level shedders, 81–82
 - preharvest control, 92–93
- Bill and Melinda Gates Foundation (BMGF), 23
- Biofortification
 - anthropometric measurements and blood tests, 43
 - Burkina Faso, 44–45
 - cereal, 42–43
 - CMF, 44
 - definition, 23
 - lysine plus threonine, 42
 - micronutrient deficiency, 46
 - PDCAAS, 45–46
 - QPM, 44
 - transferrin and hemoglobin levels, 43
 - wheat bread, 43
- Biosorghum project, 23–24
- Bradyzoites, 3–5

C

- Carcass contamination
 - C. difficile*, 60–61
 - E. coli* O157:H7
 - fecal shedding and hide prevalence, 70, 89
 - probability, 81
- Carp skin gelatin, 139
- C. difficile* infection (CDI), 54

- Cerebral biopsy, 10
- Cerebral toxoplasmosis, 9–10
- Clostridial spores, 60
- Clostridium difficile*
 - animal feed, 60
 - detection, food
 - bacteriologic culture, 55
 - enrichment broths, 55–56
 - recovery rates, 55, 57
 - environmental strains/organisms, 57, 60
 - food contamination, 60–61
 - genotypes, 61–62
 - human colonization, 62–63
 - isolation, 54
 - meat and meat products, 57–59
- Cold-water fishes, 129–130
- Collagen
 - α -chains, 122–123
 - amino acids, 123
 - β - and γ -chain, 123
 - fibrils, 121–122
 - gelatin conversion
 - alkali and acid process, 125
 - drying and grinding process, 126
 - extraction method, 125–126
 - pretreatment process, 125
 - removal of unwanted materials, 124
 - hydroxyproline, 123
 - isoelectric point, 137
 - molecular structure, 122
 - molecules arrangement, 121–122
 - stabilization, hydrogen bonds, 123–124
 - structural protein, 25
 - triple helix structure, 122
 - tropocollagen, 122
 - types, 124
- Cow's milk formula (CMF), 44

D

- DDG. *See* Dried distillers grain with/without solubles
- Direct/indirect fecal–oral exposure, 75
- Dried distillers grain with/without solubles (DDGS), 84–85

E

Electrolyzed oxidizing water, 74
Enterobacteriaceae, 93
Escherichia coli O157:H7
 animal stress
 catecholamine norepinephrine, 87
 heat stress, 88
 immune response, 87
 livestock response, 88
 microbial food safety risk, 86
 practical implications, 87
 weaning, 87
 antimicrobial carcass interventions, 97
 bacterial diarrhea, 68
 beta-agonists, 88–89
 diet effects, shedding and persistence
 barley grain, 84
 cattle fed barley *vs.* cattle fed cracked,
 84, 86
 distillers grains, 84
 energy-dense grain rations, 84
 gastrointestinal tract and fecal
 incidence, 86
 potential effects, 83–84
 ruminal fluid, 83
 ruminant animals, 82–83
 WDGS and DDGS, 84–85
 fecal shedding, 70
 feedlot cattle, 89
 high-level shedders
 bacteriophage, 81–82
 cattle hide contamination, 81
 chain of events, 81–82
 colonization, 81
 feedlot pens, 80
 mathematical modeling, 80–81
 super-shedders, 80, 82
 ionophores, 89
 modes of transmission, 69
 outbreak, water, 69
 preharvest control
 bacteriophage, 92–93
 brown seaweed product, 95
 chlorate, 93–94
 cottonseed, 94
 esculin and esculin, 95
 manure and cattle pen surface
 treatments, 95–97
 neomycin sulfate, 94
 probiotics/direct-fed microbials, 91–92
 rumen modifiers, 94
 vaccines, 90–91

risk factors, 98
 seasonality of shedding
 cattle feeds, 77
 cooler temperatures, 77–78
 heat stress, cattle, 78
 human foodborne disease, 80
 melatonin, 78
 percentage of samples, 78–79
 physiological responses, animal, 78
 prevalence, 77
 sorbitol fermentation, 70
 sources and transmission, cattle
 animal feed, 72–73
 drinking water, 73–75
 feces, manures, and soils, 75–76
 flies, 71–72
 on-farm ecology, 70
 potential reservoirs/vehicles, 70–71
 prevalence, 71
 transportation and lairage, 98–99

F

Fecal–oral transmission, 75
 Feedlot calves, 61
 Fish gelatin
 chemical characteristics
 amino acid composition, 134
 peptide size, quality, 134–135
 common and potential sources, 129–130
 effects of processing conditions
 alkali and acid treatments, 135–137
 extraction temperature and duration,
 137–138
 gelatin extraction flowchart, 135–136
 storage and transportation, 138
 physical attributes
 gel strength, 130–132
 melting and gelling temperature,
 132–133
 viscosity, 131–132
 quality *vs.* mammalian gelatins, 138–139
 Freeze drying method, 140
 Frequency sweep tests, 133

G

Gelatin
 biopolymers, 120
 characteristics, 127
 chemical composition, 120
 collagen–gelatin conversion
 alkali and acid process, 125

drying and grinding process, 126
 extraction method, 125–126
 pretreatment process, 125
 removal of unwanted materials, 124
 gelatin market, 128
 gelation mechanism, 126
 industrial applications, 128–129
 manufacturing, raw material, 127–128
 parent molecule, collagen
 α -chains, 122–123
 amino acids, 123
 β - and γ -chain, 123
 fibrils, 121–122
 hydroxyproline, 123
 molecular structure, 122
 molecules arrangement, 121–122
 stabilization, hydrogen bonds, 123–124
 triple helix structure, 122
 tropocollagen, 122
 types, 124
 quality determination, 139–140
 Gelatin hydrolysate, 134
 Gelation mechanism, 126
 Gel strength
 bloom jar, 130–131
 bloom strength, 130
 different instruments and probes,
 131–132
 measurement, standard method, 130
 Glucocorticoid dexamethasone, 87
 Grand Challenges in Global Health, 23

H

Headache, 7
 Heat drying method, 140
 High-level shedders
 bacteriophage, 81–82
 cattle hide contamination, 81
 chain of events, 81–82
 colonization, 81
 feedlot pens, 80
 mathematical modeling, 80–81
 super-shedders, 80, 82
 Houseflies, 71–72
 Hydroxyproline, 123–124, 134

I

Isoionic point, 137

K

Kafirin–tannin complexation, 42

L

Leishmania gondii, 3
 Lymph node enlargement, 7

M

Mediterranean Intensive Oxidant Study
 (MINOS), 27
 Multivariate regression analysis approach, 7
 Muscle mass
 amino acids, 26–27
 death, human starvation, 27
 maintenance, 28
 malnutrition, 26
 metabolism, genesis, 26
 MINOS (*see* Mediterranean Intensive
 Oxidant Study)
 obesity relationship, 27
 protein loss, 28
 rapid starvation/dietary protein
 depletion, 26

N

Neomycin sulfate, 94
N-(*n*-butyl) thiophosphoric triamide
 (NBPT), 96

O

Ocular toxoplasmosis, 8, 10
 Oocysts sporulation, 5
 Osteopenia, 23

P

Parasitophorous vacuole, 4
 Pharmaceutical gelatin, 129
 Preharvest control
 bacteriophage, 92–93
 brown seaweed product, 95
 chlorate, 93–94
 cottonseed, 94
 esculetin and esculin, 95
 manure and cattle pen surface treatments,
 95–97
 neomycin sulfate, 94
 probiotics/direct-fed microbials, 91–92
 rumen modifiers, 94
 vaccines, 90–91
 Proline, 123–124, 134, 139
 Protein content and composition
 PDCAAS calculation, sorghum, 33, 36
 sorghum *vs.* other cereals, 32, 34–35
 tryptophan, 36

Protein digestibility
 changes, protein body, 39
 cross-linking, kafirin prolamin proteins,
 36–37
 diets, 38, 41
 disulfide bonding, 36
 grain improvement, 37
 gruels, 38
 high-*vs.* normal-lysine sorghum, 37
 nutritional parameters, 38
 rat feeding experiments, 38
 sorghum lines, 39–40
 transgenic biofortified sorghum, 41–42
 wet cooked sorghum, 36

Protein Digestibility Corrected Amino Acid
 Score (PDCAAS)
 biological value prediction, 31
 calculation, sorghum, 31–33
 protein quality, 32, 34–35
 transgenic biofortified sorghum, 42
 Protein Efficiency Ratio (PER), 31
 Pulsed-field gel electrophoresis (PFGE)
C. difficile genotypes determination,
 61–62
 patterns, 72

Q

Quality Protein Maize (QPM), 44

R

Rheological method, 132–133
 Ribotyping, 61–62

S

Sorghum protein
 biofortification
 anthropometric measurements and
 blood tests, 43
 Burkina Faso, 44–45
 cereal, 42–43
 CMF, 44
 definition, 23
 lysine plus threonine, 42
 micronutrient deficiency, 46
 PDCAAS, 45–46
 QPM, 44
 transferrin and hemoglobin levels, 43
 wheat bread, 43
 human health
 amino acids classification, 24–25
 cereal, 29

1985 FAO/WHO/UNU *vs.* 2002
 WHO/FAO/UNU requirements,
 28–29

fat, 26
 indispensable amino acids, 24
 meta-analysis, 28
 muscle mass, 26–28
 nitrogen and amino acid needs, 28
 nitrogen-containing compounds, 25
 nutrition security planning, 28
 pregnancy and lactation period, 30
 protein types, 25
 safe level protein recommendation, 29
 scoring patterns, amino acid, 30

quality
 chemical mutagenesis, 39
in vitro pepsin method, 41
 lysine-rich protein synthesis, 40
 and measurement, 31–32
 protein body change, 39
 protein content and composition, 32,
 34–36
 protein digestibility, 36–39
 sorghum lines, 39–40
 sorghum *vs.* other cereals, 34–35, 40–41
 stunted and underweight children,
 Africa, 22–23

Stress and strain sweep tests, 133

T

Tachyzoites, 3–4
 Tannins, 42
 Temperature sweep tests, 133
 Thermoreversible gels, 127
 Thymol, 96
 Tilapia skin gelatin, 138
 Time sweep tests, 133
Toxoplasma gondii. *See also* Toxoplasmosis
 control in foods, 13–14
 life cycle
 asexual developmental cycle, 3–4
 felines, 3
 interaction, host cells, 3–4
 sexual phase, 5
 stagespecific markers, 4
 Toxoplasmosis
 congenital toxoplasmosis, 2
 discovery, 3
 laboratory diagnosis and treatment
 cerebral toxoplasmosis, 9–10
 differential diagnosis, 10
 DNA detection, PCR, 9

- IgG and IgM antibodies detection, 9
 - immunocompetent *vs.*
 - immunocompromised patients, 9
 - ophthalmologic examination, 10
 - serologic tests, 9
 - misdiagnosis/underdiagnosis, 2
 - outbreaks, 12–13
 - pathogenesis and human infection
 - spectra
 - asymptomatic infection, 7
 - immunodeficiency, 8
 - infection, pregnancy, 6–7
 - interferon-gamma (IFN- γ) production, 8
 - ocular toxoplasmosis, 8
 - risk factors, 7–8
 - transmission
 - cyst infection, ME-49 strain, 11
 - foodborne transmission, 5
 - food ingestion, 11
 - oocysts ingestion, 5
 - sausage samples, 11
 - tachyzoite, 5–6
 - undercook meat, 11
 - unpasteurized milk, 11
 - Trimester, 6–7
- U**
- Unpasteurized milk, 11, 69
- W**
- Water dessert gels, 128
 - Water troughs, 73–74
 - Wet distillers grain with solubles (WDGS), 84–85