

Modulation by glycine on vascular effect of NMDA: in vivo experimental researches

L. Berrino, S. Vitagliano, S. Maione, and F. Rossi

Institute of Pharmacology and Toxicology, 1st Faculty of Medicine and Surgery,
University of Naples Federico II, Naples, Italy

Accepted July 8, 1992

Summary. The present study has been carried out to determine if glycine, an allosteric modulator of NMDA receptor, is involved in the vascular effect induced by the activation of the CNS NMDA receptors. *Icv* NMDA (from 0.01 to 1 $\mu\text{g}/\text{rat}$ in the 3rd ventricle) determined a significant increase in arterial blood pressure in conscious freely moving rats. Moreover, the hypertension was associated with behavioural modifications (jumping, rearing, teething and running). Glycine pretreatment (1 and 10 $\mu\text{g}/\text{rat}$ *icv*), significantly increased the NMDA hypertension. Alone glycine did not cause any arterial blood pressure modification while it induced a slight sedation. HA-966 (an antagonist of the glycine site on NMDA receptor) administration (1–10 $\mu\text{g}/\text{rat}$ *icv* 5 min before glycine) significantly antagonized the glycine effects on NMDA hypertension.

Alone HA-966 neither modified arterial blood pressure nor antagonized NMDA hypertension. In conclusion, our investigations confirm NMDA receptor involvement in cardiovascular function and they demonstrate that *in vivo* glycine positively modulates NMDA receptor.

Keywords: Amino acids – N-Methyl-D-aspartate (NMDA) – Glycine – HA-966 – Rat – Strychnine

Introduction

Our previous investigations have shown the excitatory amino acid (EAA) effects on the cardiovascular system and both physiological and pathological action have been seen. NMDA and KA receptors play a primary role in the cardiovascular effects exerted by EAA (Lampa et al., 1988b; Maione et al., 1992); in conscious rats intracerebroventricular administration of EAA produced hypertension and tachycardia associated with behavioural modifications (Berrino et al., 1990); in anaesthetized rats glutamate, NMDA and kainate when administered intracerebroventricularly produce hypertension and bradycardia (Lampa

et al., 1988a). It has been shown in neuropharmacological studies that glycine greatly potentiates NMDA effects acting on the glycine modulator site of NMDA receptor (Ascher and Nowak, 1987; Henderson et al., 1990; Kaplita and Ferkany, 1990). Recent evidence showed that glycine also inhibited excitatory responses to kainate, quisqualate and NMDA in the rat thalamus in vivo (Salt, 1989). The aim of the present investigation has been to determine if the allosteric modulation of glycine on NMDA receptor is also involved in the cardiovascular effects induced by the activation of NMDA receptor in vivo.

Material and methods

Subjects

Researches have been carried out on conscious freely moving male Sprague Dawley rats (220–240 g).

Surgical preparation and treatment

Arterial blood pressure was measured continuously by a Statham pressure transducer, introduced by catheter into the femoral artery and connected to a Hellige polygraph (mean basal values: 112 ± 8 mm Hg). The direct application of drugs intracerebroventricularly (*icv*) was done by a guide cannula implanted into the 3rd ventricle under ketamine anaesthesia (100 mg/Kg *ip*) 2 days before experimentation, using a stereotaxic apparatus and applying the coordinates of the atlas of Paxinos and Watson (1986). Microinjections were made with a Hamilton micro-syringe, in a volume of 1 μ l. Control injections were carried out with the same volume of the solvent (0.2 M phosphate buffer pH 6.5 adjusted to 7.2 with 12 mM NaOH) used to dissolve the drugs. *Icv* (3rd ventricle) NMDA from 0.01 to 1 μ g/rat was delivered in 5 sec in freely moving animals. Glycine (1 and 10 μ g/rat) was delivered in 5 sec (in the 3rd ventricle) 7 min before NMDA, in both freely moving. HA-966 (1 and 10 μ g/rat in the 3rd ventricle 5 min before glycine) was delivered in 5 sec in freely moving rats. Afterward, the stereotaxic coordinates of the cannula were checked histologically.

Brain tissue was taken out fresh and immersed in buffered formaline for 2 min; the injection site was verified using 2 consecutive sections, on stained with cresyl violet to identify areas and the other unstained to determine the diffusion and position of the dye.

Drugs

The following drugs were used: NMDA (N-methyl-D-aspartate), glycine and diethyl-ether (Sigma Chemical Co. St. Louis MO, USA), ketamine hydrochloride (Parke-Davis S.p.A., Lainate – Milan, Italy); HA-966 (RBI, Natick, USA); NMDA, glycine and HA-966 [R(+)-3-amino-1-hydroxy-2-pyrrolidinone] were solubilized in 0.2 M phosphate buffer (pH 6.5) and the pH of the solution was adjusted to 7.2 with 12 mM NaOH.

Statistics

All results are expressed as mean \pm standard error (S.E.) with $p < 0.05$ considered as the level of significance; cardiovascular changes were compared by a paired or unpaired Student's *t*-test (Burn et al., 1952; Snedecor and Cochran, 1978).

Results

Icv NMDA (from 0.01 to 1 μ g/rat 3rd ventricle) determined a significant and dose-dependent increase in arterial blood pressure (Figs. 1 and 2) in freely moving




Fig. 1. Percentual variation of arterial blood pressure (ABP) after NMDA microinjections (from 0.01 to 1 µg/rat) in the 3rd ventricle (*icv*) in freely moving rats pretreated with glycine (GLY) (1 µg/rat *icv*) and/or HA-966 (HA) (1 µg/rat *icv*). Experimental groups of 5 animals were used. Significant differences are shown by asterisks (* $p < 0.05$ vs. NMDA; ** $p < 0.05$ vs. NMDA + GLY) determined by Student's *t*-test




Fig. 2. Percentual variation of arterial blood pressure (ABP) after NMDA microinjections (from 0.01 to 1 µg/rat) in the 3rd ventricle (*icv*) in freely moving rats pretreated with glycine (GLY) (10 µg/rat *icv*) and/or HA-966 (HA) (10 µg/rat *icv*). Experimental groups of 5 animals were used. Significant differences are shown by asterisks (* $p < 0.05$ vs. NMDA; ** $p < 0.05$ vs. NMDA + GLY) determined by Student's *t*-test

rats. The pressor effect was associated, only at the highest dose, with behavioural modifications (jumping, rearing, teething and running). The glycine pretreatment (1 and 10 µg/rat 3rd ventricle), significantly and in a dose-dependent manner, increased the NMDA hypertension (Figs. 1 and 2); moreover it slightly reduces

the NMDA behavioural effects in freely moving rats. Glycine caused no arterial blood pressure modifications, and induced a dose-dependent sedation. HA-966 (an antagonist of the glycine site on NMDA receptor) administration (1 and 10 $\mu\text{g}/\text{rat}$ *icv* 5 min before glycine), significantly and in dose-dependent manner, antagonized glycine effects on NMDA hypertension. Alone HA-966 neither modified arterial blood pressure and behavioural, nor antagonized NMDA hypertension (Figs. 1 and 2).

Control injections, carried out with the same volume of the solvent used to dissolved the drugs, produced not significant changes in arterial blood pressure and behaviour.

Discussion

Glycine is the main inhibitory neurotransmitter in the posterior regions of the brain and spinal cord, where its strychnine-sensitive receptors appear to be largely concentrated (Young and Snyder, 1973). However, neurochemical and neurophysiological studies on the glycine in the forebrain have yielded new data on glycine's role (Raiteri et al., 1990; Vamvakides, 1989; White et al., 1989). In rat hippocampus, Pittaluga and Raiteri (1990) and Raiteri et al. (1990) both found strychnine-inhibited and non inhibited glycine-modulated NMDA receptors. Moreover, Schmidt and Taylor (1990) also found non competitively inhibited glycine receptors to strychnine in rat hippocampus. Our results confirm that the NMDA-receptors present in the nuclei near the 3rd ventricle as the anterodorsal medial hypothalamus and the ventromedial hypothalamic area are involved in the cardiovascular function and are positively modulated *in vivo* by glycine. In fact, NMDA induced an increase in arterial blood pressure that was potentiated by glycine pretreatment. The glycine effect is due to its specific and direct activity on its NMDA recognition site because it was antagonized by HA-966, a selective antagonist of the glycine site on NMDA receptor. These findings agree with recent evidence, showing the double role of the amino acid glycine in supraspinal regions of the central nervous system, acting both at the classical strychnine sensitive receptors and at the site coupled to the NMDA receptor complex (Raiteri et al., 1990). By comparison, the behavioural effects induced by NMDA were reduced after the glycine pretreatment. These results are in agreement with those of Salt (1989) who demonstrated that glycine blocks excitatory responses in rat thalamus *in vivo*. Moreover, our data differ from the findings of Koek and Colpaert (1990) since the glycine antagonist HA-966 blocked the NMDA-induced convulsions in *icv* administered mice. They are also in contrast with Toth and Lajtha (1989) who observed that the motor effects of the NMDA injection into the rat striatum were not blocked by glycine. Our results suggest that the well known inhibitory nature of glycine reduced the NMDA elicited jumping, rearing, teething and running and that NMDA receptors, positively modulated by glycine, are present in parts of the hypothalamus which regulate the cardiovascular apparatus. *In vivo* modifications on the effects of NMDA induced by glycine suggest that this may have different action sites which modulate the cardiovascular and behavioural effects of NMDA. In conclusion, glycine could both activate the classical strychnine-sensitive receptors and the

strychnine resistant site coupled to the NMDA receptor complex, the former inhibiting the excitatory behavioural, the latter specifically potentiating glutamergic transmission mediated by NMDA subtype receptors.

Acknowledgement

This research was supported by Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST) and Consiglio Nazionale delle Ricerche (CNR), Rome, Italy.

References

- Ascher P, Nowak L (1987) Ionic channels in mouse astrocytes in culture. *Trends Neurosci* 10: 284–287
- Berrino L, Matera MG, Maione S, Vitagliano S, Loffreda A, Marmo E (1990) Excitatory amino acids and cardiovascular apparatus: experimental studies on conscious rats with L-glutamate, *N*-methyl-D-aspartate, kainate and quisqualate. *Acta Physiol Hungarica* 75 [Suppl]: 31–32
- Burn JH, Finney DJ, Goodwin LG (1952) *Biological standardization*, 2nd edn. Oxford Medical Press
- Henderson G, Johnson JW, Ascher P (1990) Competitive antagonists and partial agonists at the glycine modulatory site of the mouse *N*-methyl-D-aspartate receptor. *J Physiol (Lond)* 430: 189–212
- Lampa E, Berrino L, Susanna V, Angrisani M, Marrazzo R, Maione S, De Santis D, Fici F, Marmo E (1988a) Glutamergic transmission and cardiovascular apparatus in normotensive rats. *Drug Exptl Clin Res* 14: 59–69
- Lampa E, Cazzola M, Angrisani M, Matera C, Lisa M, Susanna V, Marrazzo R, Chieppa S, Marmo E (1988b) Cardiovascular effects of intracerebral injection of glutamate and kainate in rats. *Acta Pharmacol Sin* 9: 252–258
- Maione S, Vitagliano S, Berrino L, Lampa E, Rossi F (1992) Participation of arginine vasopressin-mediated and adrenergic system-mediated mechanisms in the hypertension induced by intracerebroventricular administration of NMDA in freely moving rats. *Neuropharmacology* 31: 403–407
- Kaplitia PV, Ferkany JW (1990) Evidence for direct interactions between the NMDA and glycine recognition sites in brain. *Eur J Pharmacol* 188: 175–179
- Koek W, Colpaert FC (1990) Selective blockade of *N*-methyl-D-aspartate (NMDA)-induced convulsions by NMDA antagonists and putative glycine antagonists: relationships with phencyclidine-like behavioral effects. *J Pharmacol Exp Ther* 252: 349–357
- Paxinos G, Watson C (1986) *The rat brain in stereotaxic coordinates*, 2nd edn. Academic Press, San Diego
- Pittaluga A, Raiteri M (1990) Release-enhancing glycine dependent presynaptic NMDA receptors exist on noradrenergic terminals of hippocampus. *Eur J Pharmacol* 191: 231–234
- Raiteri M, Fontana G, Fedele E (1990) Glycine stimulates [³H]noradrenaline release by activating a strychnine-sensitive receptor present in rat hippocampus. *Eur J Pharmacol* 184: 239–250
- Salt TE (1989) Modulation of NMDA receptor-mediated responses by glycine and D-serine in the rat thalamus in vivo. *Brain Res* 481: 403–406
- Schmidt CJ, Taylor VL (1990) Strychnine-sensitive, glycine-induced release of [³H]norepinephrine from rat hippocampal slices. *J Neurochem* 54: 2077–2081
- Snedecor G, Cochran W (1978) *Statistical methods*, 6th edn. The Iowa State University Press, Ames, Iowa
- Toth E, Lajtha A (1989) Motor effects of intracaudate injection of excitatory amino acids. *Pharmacol Biochem Behav* 33: 175–179

- Vamvakides A (1989) Quelques observations pharmacologiques sur l'action des derives de la glycine en relation avec des particularites du systeme glycinergique central. *Aggressologie* 30: 465–466
- White WF, Brown KL, Frank DM (1989) Glycine binding to rat cortex and spinal cord: binding characteristics and pharmacology reveal distinct population of sites. *J Neurochem* 53: 503–512
- Young AB, Snyder SH (1973) Strychnine binding associated with glycine receptors of the central nervous system. *Proc Natl Acad Sci USA* 70: 2832–2836

Authors' address: Prof. Dr. F. Rossi, Chairman of the Institute of Pharmacology and Toxicology, 1st Faculty of Medicine and Surgery, University of Naples, Via Costantinopoli 16, I-80138 Naples, Italy.

Received June 25, 1992