

Amino acid and vitamin requirements in mammalian cultured cells

Review Article

K. Yamamoto and A. Niwa

Department of Microbiology, Dokkyo University School of Medicine, Mibu, Tochigi,
Japan

Accepted January 11, 1993

Summary. The development of the tissue culture technique has enabled us to cultivate mammalian cells in a way which is similar to that in use with bacterial cells. As such, the nutritional requirements of mammalian cells in culture have been studied with simplicity and exactness. According to Eagle's extensive works it is accepted that cultured cells generally require 13 amino acids, 8 or 9 vitamins, glucose and 6 inorganic salts. However, although some cultured cells have a capacity for the biosynthesis of Eagle's essential nutrients and others require non-essential nutrients.

In this review we will discuss the amino acid and vitamin requirements of cultured cells, and a cell line (R-Y121B·cho) which propagates continuously in a chemically defined medium containing 11 amino acids, 7 vitamins, glucose and 6 ionic salts. Arginine, glutamine, tyrosine and choline are synthesized in the R-Y121B·cho cells.

Keywords: Amino acids – Vitamin – Cultured cell – Nutritional requirement

Harrison employed frog lymph as supporting and nutrient media to study living nerve fibers. In his paper he described that the medullary tube from frog embryos survived and nerve fibers grew out into the lymph clot (Harrison, 1907). This showed that cells could survive and develop in tissue culture and could be used effectively to tackle biological questions. In the early studies of tissue culture, cells were usually embedded in biological materials such as plasma or fibrin clot. Single cells could not be kept alive at all and the culture remained in "tissue culture".

In 1948, Earle and his colleagues reported that they were able to grow a single cell isolated from carcinogen-treated mice and to obtain from them cultures which were designated L-cell (Sanford et al., 1948). To establish cultures from single cells it is necessary to dissociate tissues into viable single cells.

Rous and Jones had achieved the release of cells from the outgrowth of plasma clot cultures by digesting the clot and culture with trypsin (Rous and Jones, 1916). The trypsinization technique prepared viable dispersed cells for cell cultures from whole embryos, organs, tissues, or other cultures.

For many years, fluid-medium cultures had been prepared from minced and chopped tissues, originally for virus studies (Maitland and Maitland, 1928; Parker and Hollender, 1945a; Parker and Hollender, 1945b). These medium usually consisted of a balanced salt solution enriched with naturally occurring materials such as serum, body fluids and exudates, and extracts of tissues and organs. But the complexity and variability of these biological materials made it difficult to examine the metabolic processes of cultured cells under controlled conditions.

A major advance for nutritional requirements of cultured cells was introduced by Eagle and his colleagues (Eagle, 1955a; Eagle, 1955b; Eagle, 1959). They employed cell lines which adhered to the surface of the glass containers or grew in suspension. These cell lines propagated in a medium consisting of an arbitrary mixture of amino acids, vitamins, cofactors, and serum protein. In such a system the specific nutrients could be determined by the omission of a single essential component resulting in the early death of the culture. Eagle and his colleagues used this system to study the essential nutritional requirements for cultured cells. Although there were specific requirements for some animal cell lines, a relatively limited number of metabolites were essential for apparently indefinite propagation of cell cultures, whether human or animal in origin, and whether deriving from normal or malignant tissue. The minimal medium contained 13 amino acids, 8 vitamins, 6 ionic species, glucose and serum protein. This medium is called Eagle's minimum essential medium (Eagle's MEM) and is still in use at many laboratories (Eagle, 1959).

Although Eagle's MEM contained minimal components for growth and multiplication of cultured mammalian cells, some components were not essential in feeding experiments in animals. The fundamental point of our research is to explain the fact that culture cells require some nutrients which are not required for growth *in vivo*. In this review we outline the amino acid and vitamin requirements in cultured cells and also cell lines which required less than 28 nutrient factors.

Amino acid requirements in mammalian cell cultures

It is known that proteins are composed of twenty amino acids. These amino acids are classified into those which are indispensable and those which are dispensable, with respect to their growth effects. Humans need only 8 amino acids, that is, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine. In addition to the above eight amino acids, rats require histidine (Table 1). Although amino acid requirements of cultured cells differ from one cell culture to another, it is widely accepted that the 13 amino acids in Eagle's MEM were essential for the survival and growth of cultured cells. Eagle pointed out three possibilities to explain this fact; 1) loss of biosynthetic mechanisms, 2) lack of appropriate precursors or cofactors, 3) limited biosyn-

Table 1. Essential amino acids in man and rat¹ and amino acids composition in Eagle's MEM²

Subjects	<i>Homo sapiens</i>			<i>Rattus norvegicus</i>		MEM
	< 1 yr old	> 1 yr old	Adult ♂	Immature	Adult	
Requirement for	Growth	Growth	Maintenance	Growth	Maintenance	Growth
Unit of measurement	mg · kg body wt ⁻¹ · da ⁻¹	mg · da ⁻¹		% of diet		mg · l ⁻¹
L-Arginine	—	—	—	0.60	—	105
L-Asparagine	—	—	—	0.40	—	—
L-Cystine	+	+	+	+	+	24
L-Glutamine	—	—	—	4.00	—	292
L-Histidine*	34	—	—	0.30	0.07	31
L-Isoleucine*	126	1000	700	0.55	0.43	52
L-Leucine*	150	1500	1100	0.75	0.25	52
L-Lysine*	103	1600	800	0.90	0.14	58
L-Methionine*	45 ⁴	800 ⁴	1100 ⁶	0.60 ⁶	0.23 ⁶	15
L-Phenylalanine*	90 ⁵	800 ⁵	1100 ⁷	0.80 ⁷	0.19 ⁷	32
L-Proline	—	—	—	0.40	—	—
L-Tyrosine	+	+	+	+	+	36
L-Threonine*	87	1000	500	0.50	0.17	48
L-Tryptophan*	22	250	250	0.15	0.07	10
L-Valine*	105	900	800	0.60	0.31	46
DAAN ³	—	—	—	0.55	0.42	—

¹ Altmann P. L. and Dittmer D. S. ed.: Biology data book 2nd ed. Vol. III, Federation of American Societies for Experimental Biology, Bethesda, MD, USA, pp1473, 1974.

² Eagle H.: Science 130: 432, 1959.

³ dispensable amino acid nitrogen.

⁴ In presence of adequate L-cystine.

⁵ In presence of adequate L-tyrosine.

⁶ 30–50% of total requirement may be furnished by L-cystine.

⁷ 30–50% of total requirement may be furnished by L-tyrosine.

thetic capacity, sufficient for survival but for growth (Eagle, 1959). Cell lines which do not require some amino acids in Eagle's MEM, have been reported. These cells shows why these amino acids are usually essential for most cultured cells. We will describe individual amino acids in the following sections.

Arginine

The pathway of arginine biosynthesis in animal tissues is illustrated in Fig. 1. Arginine is synthesized from ornithine through citrulline and arginosuccinate. The enzymes for the conversion of citrulline to arginine are detectable in most tissues (Ratner, 1973). Most cell lines can utilize citrulline as a precursor for arginine (Levintow and Eagle, 1961; Naylor et al., 1976; Sun et al., 1979). But while citrulline synthesis from ornithine is unique in liver cells *in vivo*, most cultured cells can not utilize ornithine regardless of whether the cells were derived from a normal liver or hepatoma (Naylor et al., 1976). In a few cases, the cell growth in the arginine-free medium containing ornithine was reported




Fig. 1. Pathways of arginine and glutamine biosynthesis in mammals

in rat hepatoma cultures (Niwa et al., 1979; Niwa et al., 1980; Goss, 1984; Delers et al., 1984) and hepatocyte-hepatoma hybrid cells (Widman et al., 1979; Farmer and Goss, 1991). These cell lines and their ornithine carbamoyltransferase activities are shown in Table 2. Ornithine is made by transamination from glutamate- γ -semialdehyde in mammals. However, any cell lines have not been reported which grow in ornithine-free medium.

In primary culture of liver cells ornithine carbamoyltransferase activity usually decayed to an undetectable level within a month after the initiation of cultivation (Yasumura et al., 1978). Leffert and Paul used an arginine-free and ornithine-containing medium to selectively multiply fetal rat hepatocytes, which were able to form the arginine they needed (Leffert and Paul, 1973; Hasegawa et al., 1982). Even though hepatocytes were selected from arginine-required liver-derived cells in arginine-free and ornithine-containing media, continuous growth of adult hepatocytes has not been reported as yet.

Glutamine

Glutamine is found in relatively high concentrations in many mammalian cells where it serves as an ammonia scavenger and a nitrogen donor for the biosynthesis of a number of important compounds such as nucleotides, amino acids and amino sugars. Furthermore, glutamine was used as the major source of energy for cultured human carcinoma cells (HeLa cell, Reitzer et al., 1979) and mouse macrophage (Newsholme and Newsholwe, 1989). Glutamine is synthesized from glutamate by glutamine synthetase, and glutamate is produced from α -ketoglutarate by the catalytic action of glutamate dehydrogenase (Fig. 1).

The necessity of glutamine in the amino acid requirements of cultured cells was first observed by Ehrensward et al. (1949). Since then, studies with several cell lines have demonstrated that glutamine is an essential requirement for optimal proliferation of cultured cells. Eagle and his colleagues reported that both L cells and HeLa cells degenerated and died in the glutamine-free medium. The maximally effective concentration was 0.2 to 0.5 mM for L cells, and 1 to 2 mM for HeLa cells (Eagle et al., 1956a).

Table 2. Ornithine carbamoyltransferase (OCT) activity and cellular growth in an arginine-deprived medium of various cell lines

Cell line (origine)	Serum ¹ in medium	Growth in – Arg + Orn	OCT activity nmol. Cit/min/mg Protein	Reference
7800Ci (rat Morris hepatoma 7800)	5% HS + 5% FBS	–	1	Richardson et al., 1974
R-Y117-21B (rat Reuber hepatoma)	2% DCS	+	30	Niwa et al., 1979
Clone 119 (rat hepatocyte-hepatoma hybrid)	10% FBS	+	43	Widman et al., 1979
R-Y121B (rat Reuber hepatoma)	0	+	122	Niwa et al., 1980
777orn (rat Morris hepatoma 777)	10% FBS	+	120	Goss, 1984
Fa32A50 (rat Reuber hepatoma)	10% FBS	+	65	Delers et al., 1984
BWsf hybrid (mouse hepatoma-hepatocyte hybrid)	5% FBS	+	6	Farmer and Goss, 1991

¹ HS Horse serum, FBS Fetal bovine serum, DCS Dialysed calf serum

In media supporting growth of L cells, glutamine could not be effectively replaced by any concentration of glutamic acid, although, in the case of HeLa cells, glutamic acid at the optimal concentration of 20 mM was an effective substitute for glutamine (10–20 times the glutamine concentration). Several investigators (DeMars, 1958; Paul and Fottrell, 1963; Goetz et al., 1973; Tiemeier and Milman, 1972; Yasumura et al., 1978) have reported that various cell lines also could grow in a glutamine-free and glutamate-supplemented medium due to the ability to produce glutamine synthetase, though the enzyme was usually induced under a condition of high concentration of glutamate. Although glutamine and/or glutamic acid is essential for the great majority of cell lines, some types of cells can synthesize enough glutamine to satisfy their needs (Table 3). Green monkey kidney cells (VERO-300 cells) have a glutamate dehydrogenase and grow in protein-free medium lacking glutamine and glutamate (Yasumura et al., 1978). Rat hepatoma cells (R-Y121B cells) can also grow in glutamine-, glutamate- and protein-free medium (Niwa et al., 1980).

Tyrosine

Tyrosine is formed from an essential amino acid, phenylalanine. It is known that three enzymes (phenylalanine hydroxylase, tyrosine hydroxylase and tryptophan hydroxylase) are responsible for the conversion of phenylalanine to tyrosine (Breakefield and Nirenberg, 1974). The phenylalanine hydroxylase activity has been found in only three mammalian tissues; the liver, kidney and pancreas (Tourian et al., 1969). In the case of cell culture a variant strain of HeLa cells was able to convert phenylalanine to tyrosine, though not at a rate sufficient enough to support growth (Eagle et al., 1957). Haggerty and his colleagues found the presence of phenylalanine hydroxylase as to be a constitutive enzyme in two rat hepatoma cell lines (H4-II-E-C₃, MH₁C₁, Haggerty et al., 1973). These two cell lines grew in medium lacking tyrosine, when phenylalanine was present (Haggerty et al., 1975). A pigmented murine melanoma line (PS1, Breakefield et al., 1978), a human hepatoma cell line (H-H1, Choo et al., 1980) and a mouse hepatoma cell line (BWTG3, Farmer and Goss, 1991) had phenylalanine hydroxylase and grew in tyrosine-free medium.

The tyrosine hydroxylase is found in the adrenal medulla, brain and other sympathetically innervated tissue (Nagatsu et al., 1964). Mouse neuroblastoma cells (N-115 cells) were able to grow in the absence of tyrosine due to the presence of tyrosine hydroxylase in the cell (Breakefield and Nirenberg, 1974).

Tryptophan hydroxylase is found in the brain (Graham-Smith, 1964). Mouse mastocytoma cells (P815-X2.1) possessing this enzyme were able to grow in a tyrosine-free medium but prolonged cultivation was not successful (Choo et al., 1976).

Cystine and methionine

Cystine is synthesized from methionine by the pathway shown in Fig. 2. In the first step of this sequence, methionine loses the methyl group from its sulfur atom to become homocysteine. Homocysteine reacts with serine to yield cy-

Table 3. Serially cultured cells growing in an essential amino acid-free medium

Amino acid	Enzyme involved	Cell line (origine)	Reference
Glutamine	Glutamic dehydrogenase + Glutamine synthetase	VERO-300 (normal monkey kidney) R-Y121B (rat Reuber hepatoma)	Yasumurat et al., 1979 Niwa et al., 1981
Tyrosine	Phenylalanine hydroxylase	H4-II-EC3 (rat Reuber hepatoma) MH1C1 (rat Morris hepatoma) PS1 (mouse pigmented melanoma) H-H1 (human hepatoma) BWTG3 (mouse hepatoma) N-115 (mouse neuroblastoma)	Haggerty et al., 1975 Haggerty et al., 1975 Breakefield et al., 1978 Choo et al., 1980 Famer and Goss, 1991 Breakefield and Nirenberg, 1974
	Tyrosine hydroxylase		




Fig. 2. Pathways of cysteine and methionine biosynthesis in mammals

stathionine. Cystathionine is cleaved hydrolytically to yield cysteine. Although a number of heteroploid human cell lines were able to synthesize cystine from methionine, the cells in a cystine-free medium died unless the population density was maintained in excess of 200,000 to 500,000 cells/ml. Even if the cells were provided with homocysteine or cystathionine, growth was observed only if the initial population density was in excess of 10,000 to 60,000 cells/ml. A similar population-dependent requirement for metabolites such as asparagine, glutamine, serine, inositol and pyruvate was reported (Eagle et al., 1961; Eagle and Piez, 1962). Further studies showed this to be the case for heteroploid cells, but human diploid cells required cystine (Eagle et al., 1966). Some variant human diploid fibroblast clones were able to grow in the medium containing homocysteine in place of cystine (Jacoby and Littlefield, 1971; Hankinson and Jacoby, 1975). In rats, the enzymes of cystine synthesis are predominantly in the liver, pancreas and kidney (Mudd et al., 1965). A most cell lines of mouse and rat cells could grow using cystathionine in place of cysteine (Naylor et al., 1976). However homocysteine could only be used by hepatoma cells (Goss, 1986; Farmer and Goss, 1991).

In mammals methionine is synthesized by at least two enzymes, vitamin B₁₂-dependent 5-methyltetrahydrofolate-homocysteine methyltransferase and betaine-homocysteine methyltransferase (Fig. 2).

The vitamin B₁₂-dependent methyltransferase activity was widely distributed in rat tissue (Finkelstein et al., 1971) and was detected in various cultured mammalian cells. These cultured cells grew in a methionine deficient medium supplemented with homocysteine and vitamin B₁₂ (Mangum and North, 1968; Mangum et al., 1969; Kamely et al., 1973; Halpern et al., 1974).

Betaine-homocysteine methyltransferase was detectable in only a few mammalian tissues (Finkelstein et al., 1971). Although this transferase was detected in cultured mouse cells (Grzelakowska-Sztart and Balińska, 1980), they did not multiply in the methionine-deficient medium.

α-Keto analogues of the essential amino acids and D-amino acids

Essential amino acids were able to be replaced by the corresponding α-keto acids (Eagle, 1959; Naylor et al., 1976). Because the essential parts of the indispensable amino acids are their carbon skeletons, not their amino groups, the α-keto

analogues of the essential amino acids could accept amino groups from excess nonessential amino acid by transaminase action.

D-amino acid can be converted to the L-form by D-amino acid oxidase and aminotransferases. Human oral carcinoma cells (KB cells) were able to utilize D-cystine (Naylor et al., 1976). Green monkey kidney cells (VERO-317 and VERO-300 cells) propagated continuously in the media in which L-amino acid was replaced by D-isomer of leucine, phenylalanine and/or valine (Yasumura et al., 1978).

Non-essential amino acids

The individual requirements for non-essential amino acids will vary from one cell line to another. In some situations there is a specific requirement for an amino acid normally considered non-essential. Asparagine was required by certain malignant cells, including a number of lines of leukemia (McCoy et al., 1956; Haley et al., 1961; Broome, 1963; Summers and Handschumacher, 1971; Ohnuma et al., 1971; Schrek et al., 1973). In addition, there was an unusual requirement of glycine by Rhesus monkey testicular cells (Tytell et al., 1958) and monkey kidney cells (Eagle et al., 1958). Requirement of serine by rabbit fibroblast cells (Haff and Swim, 1957) and leukemic cells (Regan et al., 1969), that of combination of glycine and serine by Novikoff hepatoma cells (McCoy et al., 1959) and that of alanine and serine by mouse fibroblast cells (Nagle and Brown, 1971) were reported.

Vitamin requirements in mammalian cell cultures

Eagle (1955a) reported the HeLa and L cells required 8 vitamins (biotin, choline, folic acid, nicotinamide, pantothenate, pyridoxal, riboflavin, and thiamine). In this case Eagle's medium was supplemented with serum, which would supply biotin, vitamin B₁₂, and fat-soluble vitamins. Sanford et al. (1963) showed that cells required 8 vitamins (Eagle's 8 vitamins), after they examined the effects of the 16 vitamins, in a serum-free chemically defined medium on cell survival and proliferation rates. Eight vitamins such as A, D, K, E, C, p-aminobenzoic acid, B₁₂, and inositol appeared to have no significant effects on the cell growth. The fat-soluble vitamins were necessary for specialized organs, but did not play a necessary role in the metabolism of cultured cells (Smith, 1981). From these results, it was determined that Eagle's 8 vitamins are essential to sustain cell growth. However it was reported that some cell lines required vitamin B₁₂ (Higuchi and Robinson, 1973; Kan and Yamane, 1982) and inositol (Eagle, 1956b; Jackson and Shin, 1982).

Choline is necessary for a substrate of the biosynthesis of phospholipid rather than as cofactor. All of the nitrogen-containing phospholipids can be derived from phosphatidylserine. Decarboxylation of the serine residue gives phosphatidylethanolamine, and successive methylations yield phosphatidylcholine. There is also some direct incorporation of choline into phospholipids (Fig. 3). In cultured cells, the capacity to synthesize choline *de novo* is inadequate to meet the demand for phospholipid synthesis. Therefore, choline is essential in cultured




Fig. 3. Pathways of choline and phosphatidylcholine biosynthesis in mammals

cells. Phosphatidylcholine can be synthesized by the methylation of phosphatidylethanolamine in liver cells. Some cell lines were able to utilize monomethylethanolamine and dimethylethanolamine as a precursor for choline (Glaser et al., 1974, Schroeder et al., 1976). Rat hepatoma R-Y121B·cho cells were able to grow in choline-free medium (Yamamoto et al., 1985).

The vitamins act as the cofactor and coenzyme associated with many enzyme systems. Co-enzymes can generally be substituted for the parent vitamins. The active form of pyridoxal is pyridoxal phosphate. Pyridoxal phosphate serves as the tightly bound prosthetic group of a number of enzymes catalyzing reactions of amino acids. The most common and best known example of these is transamination. When cells were cultured in the absence of pyridoxal, all the amino acids had to be supplied in the culture medium (Swim and Parker, 1958; Sanford et al., 1963).

Auxotrophic mutants

An auxotrophic mutant is considered here to be a genetically altered cells having at least one more nutritional requirements than the parent cells. Auxotrophic mutants are useful in studying the various problems in mammalian cellular genetics and metabolism of cultured cells. A method of selecting for auxotrophic mutants was first described by DeMars and Hooper (1960). It utilized the fact that aminopterin is more toxic for growing than for nongrowing cells. Glutamine-requiring cells were selected from HeLa cells. Puck and Kao developed a technique to obtain auxotrophic mutants. They treated Chinese hamster ovary and Chinese hamster lung cells with mutagenic agents and exposed them to BrdU followed by illumination with near-visible light (Puck and Kao, 1967). Then they isolated the auxotrophs for proline; glycine; inositol (Kao and Puck, 1968; Kao and Puck, 1972). Auxotrophs for serine (Jones and Puck,

1973) and tyrosine (Choo and Cotton, 1977) were also isolated using the same technique.

Establishment of cultured cells that minimize the requirement for organic nutrient

Among Eagle's 13 amino acids and 8 vitamins, 4 of the amino acids (arginine, cystine, glutamine, and tyrosine) and choline are not essential for rats, however, they are essential for cells in culture (Table 1). Because these amino acids and choline are synthesized in liver *in vivo*, liver cell cultures might be capable of synthesizing these nutrients at a rate sufficient enough to sustain growth. Therefore we have tried to establish prototrophs to these nutrients by a multi-step selection procedure using rat hepatoma cells.

The cell line chosen as a parental line was H4-II-E which was established from Reuber hepatoma H-35 (Pitot et al., 1964). H4-II-E cells were maintained in Eagle's MEM supplemented with biotin (EM # 303) and 10% of fetal bovine serum. Since drug-resistant cells are useful in cell hybridization experiments, the parental cells were adapted to grow in EM # 303 including 5-bromodeoxy-




Fig. 4. The growth curves of R-317B, R-117·21B, R-Y121B and R-Y121B·cho cells. **A** R-317B cells were tested in glutamine-deprived and glutamate-supplemented MEM (EM # 317, ●) or arginine-deprived and ornithine-supplemented EM # 317 (○). Media were supplemented with 1% of DCS. **B** R-117·21B cells tested in EM # 317 (●) or arginine-deprived and ornithine-supplemented EM # 317 (○). Media were supplemented with 1% of DCS. **C** R-Y121B cells were tested in arginine-, glutamine-, and tyrosine-deprived and ornithine-supplemented MEM (EM # Y121, ●) or choline-deprived EM # Y121 (○). **D** R-Y121B·cho cells were tested in choline-deprived EM # Y121 (○) or choline- and inositol-deprived EM # Y121 (●)

uridine (BrdU). The BrdU-resistant line was next adapted to the medium containing glutamic acid in place of glutamine (EM # 317). The concentration of serum in the medium was decreased gradually to 1% dialyzed calf serum (DCS) during 2 to 6 months of serial subculturing. The cells cultivated in EM # 317 containing BrdU (50 μ g/ml) and DCS (1%) were designated R-317B (Fig. 4A).

To establish a cell line that had complete set of the urea cycle enzymes, R-317B cells were maintained in an arginine-deprived, ornithine-supplemented medium (EM # 117·21) containing 1% DCS. Thus, we established a cell line, R-117·21B, which grew continuously in EM # 117·21 containing 1% DCS and 50 μ g/ml BrdU (Fig. 4B). R-117·21B cells, like H4-II-E cells, preserved phenylalanine hydroxylase, and propagated in tyrosine-free medium without any lag period.

Although cells in culture usually require some components of serum, serum contained amino acids as well as many other unknown substances. Even if we performed serum dialysis exhaustively, amino acids do not dialyze out completely from serum, and moreover, during the course of culture, serum protein would yield free amino acids by proteolysis (Piez et al., 1960; Niwa et al., 1979). Therefore, studies in amino acid requirements of cultured cells are best conducted in a totally synthetic medium. Thus we tried to adapt R-117·21B cells in serum-free medium. Then the cells were adapted in a glutamic acid-deprived medium. The cells were designated R-Y121B cells, which grew continuously in fully an autoclavable synthetic medium (EM # Y121) which contained 11 amino acids (cystine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, and ornithine) (Fig. 4C). R-Y121B cells have a complete set of the urea cycle enzymes, phenylalanine hydroxylase, and synthesize urea via the urea cycle pathway (Table 4, Niwa et al., 1980; Yamamoto et al., 1981).

Table 4. Enzyme activities of normal rat liver and R-Y121B cells

Enzyme	R-Y121B	Normal rat liver
	nmol product/min per mg cell protein	
Carbamoylphosphate synthetase (ammonia)	7	36
Ornithine carbamoyltransferase	122	821
Argininosuccinate synthetase	5	17
Argininosuccinate lyase	16	23
Arginase	860	10800
Phenylalanine hydroxylase	10	245
Tyrosine aminotransferase	18	25

From the R-Y121B cells, a choline- and inositol-prototroph was also established. R-Y121B cells were cultivated in EM # Y121 with ethanolamine in place of choline. After 15 subcultures in this medium, the cells were subcultured in a choline- and ethanolamine-deprived medium. The cells cultivated in choline-free medium were named R-Y121B·cho cells which proliferated readily in inositol- and choline-free EM # Y121 (Fig. 4D).

The establishment of R-Y121B·cho cells indicate that the cells have not only biosynthetic mechanisms for some of Eagle's essential nutrients but also the products were sufficient for survival and growth. Further effort to establish an ornithine and/or cysteine non-required cells is continued.

References

- Breakefield XO, Nirenberg MW (1974) Selection for neuroblastoma cells that synthesize certain transmitters. *Proc Natl Acad Sci USA* 71: 2530–2533
- Breakefield XO, Castiglione CM, Halaban R, Pawelek J, Shiman R (1978) Phenylalanine hydroxylase in melanoma cells. *J Cell Physiol* 94: 307–314
- Broome JD (1963) Evidence that the L-asparaginase of guinea pig serum is responsible for its antilymphoma effects. II Lymphoma 6C3HED cells cultured in a medium devoid of L-asparagine lose their susceptibility to the effects of guinea pig serum in vivo. *J Exp Med* 118: 121–148
- Choo KH, Cotton RGH, Danks DM (1976) Phenylalanine hydroxylation and tyrosine requirement of cultured cells. Evidence of phenylalanine hydroxylation in mastocytoma cells. *Exp Cell Res* 101: 370–382
- Choo KH, Cotton RGH (1977) Genetics of the mammalian phenylalanine hydroxylase system. I. Isolation of phenylalanine hydroxylase-deficient tyrosine auxotrophs from rat hepatoma cells. *Somat Cell Genet* 3: 457–470
- Choo KH, Cotton RGH, Jennings IG, Fowler K, Danks DM (1980) Genetics of the mammalian phenylalanine hydroxylase system. IV. Evidence of phenylalanine hydroxylase in a cultured human hepatoma cell line. *Biochem Genet* 18: 955–968
- Delers A, Szpirer J, Szpirer C, Saggioro D (1984) Spontaneous and 5-azacytidine-induced reexpression on ornithine carbamoyl transferase in hepatoma cells. *Mol Cell Biol* 4: 809–812
- DeMars R (1958) The inhibition by glutamine of glutamyl transferase formation in cultures of human cells. *Biochim Biophys Acta* 27: 435–436
- DeMars R, Hooper JL (1960) A method of selecting for auxotrophic mutants of HeLa cells. *J Exp Med* 111: 559–572
- Eagle H (1955a) Nutrition needs of mammalian cells in tissue culture. *Science* 122: 501–504
- Eagle H (1955b) The minimum vitamin requirements of the L and HeLa cells in tissue culture, the production of specific vitamin deficiencies, and their cure. *J Exp Med* 102: 595–600
- Eagle H, Oyama VI, Levy M, Horton CL, Fleischman R (1956a) The growth response of mammalian cells in tissue culture to L-glutamine and L-glutamic acid. *J Biol Chem* 218: 607–616
- Eagle H, Oyama VI, Levy M, Freeman A (1956b) Myo-inositol as an essential growth factor for normal and malignant human cells in tissue culture. *Science* 123: 845–847
- Eagle H, Piez KA, Fleischman R (1957) The utilization of phenylalanine and tyrosine for protein synthesis by human cells in tissue culture. *J Biol Chem* 228: 847–861
- Eagle H, Freeman AE, Levy M (1958) The amino acid requirements of monkey kidney cells in first culture passage. *J Exp Med* 107: 643–652
- Eagle H (1959) Amino acid metabolism in mammalian cell cultures. *Science* 130: 432–437
- Eagle H, Piez KA, Oyama VI (1961) The biosynthesis of cystine in human cell cultures. *J Biol Chem* 236: 1425–1428
- Eagle H, Piez KA (1962) The population-dependent requirement by cultured mammalian cells for metabolites which they can synthesize. *J Exp Med* 116: 29–43
- Eagle H, Washington C, Friedman SM (1966) The synthesis of homocystine, cystathionine, and cystine by cultured diploid and heteroploid human cells. *Proc Natl Acad Sci USA* 56: 156–163
- Ehrensward G, Fischer A, Stjernholm R (1949) Protein metabolism of tissue cells in vitro: 7. The chemical nature of some obligate factors of tissue cell nutrition. *Acta Physiol Scand* 18: 218–230

- Farmer AA, Goss SJ (1991) BWTG3 hepatoma cells can acquire phenylalanine hydroxylase, cystathionine synthase and CPS-1 without genetic manipulation, but activation of the silent OCT gene requires cell fusion with hepatocytes. *J Cell Sci* 98: 533–538
- Finkelstein JD, Kyle WE, Harris BJ (1971) Methionine metabolism in mammals. Regulation of homocysteine methyltransferases in rat tissue. *Arch Biochem Biophys* 146: 84–92
- Glaser M, Ferguson KA, Vagelos PR (1974) Manipulation of the phospholipid composition of tissue culture cells. *Proc Natl Acad Sci USA* 71: 4072–4076
- Goetz IE, Weinstein C, Roberts E (1973) Properties of a hamster tumor cell line grown in a glutamine-free medium. *In Vitro* 9: 46–55
- Goss SJ (1984) Arginine synthesis by hepatomas in vitro. I. The requirements for cell growth in medium containing ornithine in place of arginine, and the isolation and characterization of variant hepatomas auxotrophic for arginine. *J Cell Sci* 68: 285–303
- Goss SJ (1986) Characterization on cystathionine synthase as a selectable, liver-specific trait in rat hepatomas. *J Cell Sci* 82: 309–320
- Grahame-Smith DG (1964) Tryptophan hydroxylation in brain. *Biochem Biophys Res Commun* 16: 586–592
- Grzelakowska-Sztabert B, Balińska M (1980) Induction of betaine : homocysteine methyltransferase in some murine cells cultured in vitro. *Biochim Biophys Acta* 632: 164–172
- Haff RF, Swim HE (1957) The amino acid requirements of rabbit fibroblasts, strain RM3-56. *J Gen Physiol* 41: 91–99
- Haggerty DF, Young PL, Popják G, Carnes WH (1973) Phenylalanine hydroxylase in cultured hepatocytes. I. Hormonal control of enzyme levels. *J Biol Chem* 248: 223–232
- Haggerty DF, Young PL, Buese JV (1975) A tyrosine-free medium for the selective growth of cells expressing phenylalanine hydroxylase activity. *Dev Biol* 44: 158–168
- Haley EE, Fischer GA, Welch AD (1961) The requirement for L-asparagine of mouse leukemia cells L5178Y in culture. *Cancer Res* 21: 532–536
- Halpern BC, Clark BR, Hardy DN, Halpern RM, Smith R (1974) The effect of replacement of methionine by homocystine on survival of malignant and normal adult mammalian cells in culture. *Proc Natl Acad Sci USA* 71: 1133–1136
- Hankinson O, Jacoby LB (1975) Variant human diploid fibroblast clones able to grow in homocysteine in place of cysteine. *Exp Cell Res* 96: 138–144
- Harrison RG (1907) Observations on the living developing nerve fibre. *Proc Soc Exp Biol Med* 4: 140–143
- Hasegawa K, Watanabe K, Koga M (1982) Induction of mitosis in primary cultures of adult rat hepatocytes under serum-free conditions. *Biochem Biophys Res Commun* 104: 259–265
- Higuchi K, Robinson RC (1973) Studies on the cultivation of mammalian cell lines in a serum-free, chemically defined medium. *In Vitro* 9: 114–121
- Jackson MJ, Shin S (1982) Inositol as a growth factor for mammalian cells in culture. In: Sato GH, Pardee AB, Sirbasku DA (eds) *Growth of cells in hormonally defined media*. Cold Spring Harbor, New York; Cold Spring Harbor Laboratory, Book A, pp 75–86
- Jacoby LB, Littlefield JW (1971) Mutant human fibroblast clones able to grow on homocysteine instead of cysteine. *Exp Cell Res* 69: 447–449
- Jones C, Puck TT (1973) Genetics of somatic mammalian cells. XVII. Induction and isolation of Chinese hamster cell mutants requiring serine. *J Cell Physiol* 81: 299–304
- Kamely D, Littlefield JW, Erbe RW (1973) Regulation of 5-methyltetrahydrofolate : homocysteine methyltransferase activity by methionine, vitamin B₁₂, and folate in cultured baby hamster kidney cells. *Proc Natl Acad Sci USA* 70: 2585–2589
- Kan M, Yamane I (1982) In vitro proliferation and lifespan of human diploid fibroblasts in serum-free BSA-containing medium. *J Cell Physiol* 111: 155–162
- Kao F-T, Puck TT (1968) Genetics of somatic mammalian cells. VII. Induction and isolation of nutritional mutants in Chinese hamster cells. *Proc Natl Acad Sci USA* 60: 1275–1281
- Kao F-T, Puck TT (1972) Genetics of somatic mammalian cells. XIV. Genetic analysis in vitro of auxotrophic mutants. *J Cell Physiol* 80: 41–50
- Leffert HL, Paul D (1973) Serum dependent growth of primary cultured differentiated fetal rat hepatocytes in arginine-deficient medium. *J Cell Physiol* 81: 113–124

- Levintow L, Eagle H (1961) Biochemistry of cultured mammalian cells. *Ann Rev Biochem* 30: 605–640
- Mangum JH, North JA (1968) Vitamin B₁₂ dependent methionine biosynthesis in HEP-2 cells. *Biochem Biophys Res Commun* 32: 105–110
- Mangum JH, Murray BK, North JA (1969) Vitamin B₁₂ dependent methionine biosynthesis in cultured mammalian cells. *Biochemistry* 8: 3496–3499
- Maitland HB, Maitland MC (1928) Cultivation of vaccinia virus without tissue culture. *Lancet* 215: 596–597
- McCoy TA, Maxwell M, Neuman RE (1956) The amino acid requirements of the Walker carcinosarcoma 256 in vitro. *Cancer Res* 16: 979–984
- McCoy TA, Maxwell M, Kruse PF Jr (1959) Amino acid requirement of Novikoff hepatoma in vitro. *Proc Soc Exp Biol Med* 100: 115–118
- Mudd SH, Finkelstein JD, Irreverre F, Laster L (1965) Trassulfuration in mammals. Microassays and tissue distributions of three enzymes of the pathway. *J Biol Chem* 240: 4382–4392
- Nagatsu T, Levitt M, Udenfriend S (1964) Tyrosine hydroxylase. The initial step in norepinephrine biosynthesis. *J Biol Chem* 239: 2910–2917
- Nagle SC Jr, Brown BL (1971) An improved heat-stable glutamine-free chemically defined medium for growth of mammalian cells. *J Cell Physiol* 77: 259–264
- Naylor SL, Busby LL, Klebe RJ (1976) Biochemical selection systems for mammalian cells: the essential amino acids. *Somat Cell Genet* 2: 93–111
- Newsholme P, Newsholme EA (1989) Rates of utilization of glucose, glutamine and oleate and formation of end-products by mouse peritoneal macrophages in culture. *Biochem J* 261: 211–218
- Niwa A, Yamamoto K, Yasumura Y (1979) Establishment of a rat hepatoma cell line which has ornithine carbamoyltransferase activity and grows continuously in arginine-deprived medium. *J Cell Physiol* 98: 177–184
- Niwa A, Yamamoto K, Sorimachi K, Yasumura Y (1980) Continuous culture of Reuber hepatoma cells in serum-free, arginine-, glutamine- and tyrosine-deprived chemically defined medium. *In Vitro* 16: 987–993
- Ohnuma T, Waligunda J, Holland JF (1971) Amino acid requirements in vitro of human leukemic cells. *Cancer Res* 31: 1640–1644
- Parker RC, Hollender AJ (1945a) Propagation of Theiler's GD-VII mouse virus in tissue culture. *Proc Soc Exp Biol Med* 60: 88–93
- Parker RC, Hollender AJ (1945b) Propagation of rabies virus in tissue culture. *Proc Soc Exp Biol Med* 60: 94–98
- Paul J, Fottrell PF (1963) Mechanism of D-glutamyltransferase repression in mammalian cells. *Biochim Biophys Acta* 67: 334–336
- Piez KA, Oyama I, Levintow L, Eagle H (1960) Proteolysis in stored serum and its possible significance in cell culture. *Nature* 188: 59–60
- Pitot HC, Peraino C, Morse PA Jr, Potter VR (1964) Hepatomas in tissue culture compared with adapting liver in vivo. *Natl Cancer Inst Monograph* 13: 229–245
- Puck TT, Kao F-T (1967) Genetics of somatic mammalian cells. V. Treatment with 5-bromodeoxyuridine and visible light for isolation of nutritionally deficient mutants. *Proc Natl Acad Sci USA* 58: 1227–1234
- Ratner S (1973) Enzymes of arginine and urea synthesis. *Adv Enzymol* 39: 1–90
- Regan JD, Vodopick H, Takeda S, Lee WH, Faulcon FM (1969) Serine requirement in leukemic and normal blood cells. *Science* 163: 1452–1453
- Reitzer LJ, Wice BM, Kennell D (1979) Evidence that glutamine, not sugar, is the major energy source for cultured HeLa cells. *J Biol Chem* 254: 2669–2676
- Richardson UI, Snodgrass PJ, Nuzum CT, Tashjian AH Jr (1974) Establishment of a clonal strain of hepatoma cells which maintain in culture the five enzymes of the urea cycle. *J Cell Phys* 83: 141–149
- Rous P, Jones FS (1916) A method for obtaining suspensions of living cells from the fixed tissues for the plating out of individual cells. *J Exp Med* 23: 549–555

- Sanford KK, Earle WR, Likely GD (1948) The growth in vitro of single isolated tissue cells. *J Natl Cancer Inst* 9: 229–246
- Sanford KK, Dupree LT, Covalesky AB (1963) Biotin, B₁₂, and other vitamin requirements of a strain of mammalian cells grown in chemically defined medium. *Exp Cell Res* 31: 345–375
- Schrek R, Holcenberg JS, Batra KV, Roberts J, Dolowy WC (1973) Effect of asparagine and glutamine deficiency on normal and leukemic cells. *J Natl Cancer Inst* 51: 1103–1107
- Schroeder F, Perlmutter JF, Glaser M, Vagelos PR (1976) Isolation and characterization of subcellular membranes with altered phospholipid composition from cultured fibroblasts. *J Biol Chem* 251: 5015–5026
- Smith JR (1981) The fat-soluble vitamin. In: Waymouth C, Ham RG, Chapple PJ (eds) *The growth requirements of vertebrate cells in vitro*. Cambridge University Press, Cambridge London New York New Rochelle Melbourne Sydney, pp 343–352
- Summers WP, Handschumacher RE (1971) L5178Y asparagine-dependent cells and independent clonal sublines. Toxicity of 5-diazo-4-oxo-L-norvaline. *Biochem Pharmacol* 20: 2213–2220
- Sun NC, Sun CRY, Tennant RW, Hsie AW (1979) Selective growth of some rodent epithelial cells in a medium containing citrulline. *Proc Natl Acad Sci USA* 76: 1819–1823
- Swim HE, Parker RF (1958) Vitamin requirements of uterine fibroblasts, strain U12-79. Their replacement by related compounds. *Arch Biochem Biophys* 78: 46–53
- Tiemeier DC, Milman G (1972) Regulation of glutamine synthetase in cultured Chinese hamster cells. Induction and repression by glutamine. *J Biol Chem* 247: 5722–5727
- Tourian A, Goddard J, Puck TT (1969) Phenylalanine hydroxylase activity in mammalian cells. *J Cell Physiol* 73: 159–170
- Tytell AA, Rader Y, Krumm DL (1958) Amino acid requirements of primary trypsinized Rhesus monkey testicular cells in tissue culture. *Federation Pro* 17: 326
- Widman LE, Golden JJ, Chasin LA (1979) Immortalization of normal liver functions in cell culture: Rat hepatocyte-hepatoma cell hybrids expressing ornithine carbamoyltransferase activity. *J Cell Physiol* 100: 391–400
- Yamamoto K, Niwa A, Yasumura Y (1981) The production of urea from ornithine in rat hepatoma cells continuously cultured in a chemically defined medium. *Cell Struct Funct* 6: 367–374
- Yamamoto K, Niwa A, Yasumura Y (1985) Continuous growth and phosphatidylcholine synthesis of rat hepatoma cells in choline-deprived chemically defined medium. *J Cell Physiol* 125: 91–97
- Yasumura Y, Niwa A, Yamamoto K (1978) Phenotypic requirement for glutamine of kidney cells and for glutamine and arginine of liver cells in culture. In: Katsuta H (ed) *Nutritional requirements of cultured cells*. Scientific Societies Press, Tokyo, pp 223–255

Authors' address: Prof. A. Niwa, MD, Department of Microbiology, Dokkyo University School of Medicine, Mibu, Tochigi, 321-02, Japan.

Received December 28, 1992