

Incorporation of nitrate nitrogen in rice seedlings transferred to anaerobic conditions

Short Communication

R. Reggiani¹, F. Bertini², and M. Mattana¹

¹Istituto Biosintesi Vegetali and ²Istituto Chimica Macromolecole, C.N.R.,
Milano, Italy

Accepted March 13, 1997

Summary. Incorporation of $^{15}\text{NO}_3^-$ into amino acids was studied in 3-day-old aerobic rice seedlings (with coleoptile and root) subjected for 24h to anaerobic conditions. The incorporation of ^{15}N into glutamate, glutamine and alanine accounted for 89% and 84% of total incorporation in coleoptile and root, respectively. These findings indicate that, after the primary incorporation of ^{15}N into glutamate and glutamine, the main fate of nitrate nitrogen in rice seedlings subjected to anoxia is alanine.

Keywords: Amino acids – Anoxia – Coleoptile – Nitrate nitrogen – Rice – Root

Introduction

Nitrate assimilation is one of the main reductive processes occurring in plant tissues during aerobic metabolism. The importance of this pathway under anaerobiosis is controversial. Rice plants grow well and withstand anaerobiosis better with nitrate as the sole nitrogen source when cultured on liquid nutrient media (Malavolta, 1954). Under such condition, nitrate is better than ammonium with respect to plant growth, grain yield, some metabolic activities and cation absorption of the roots (Yamasaki and Seino, 1965).

Contrasting data are provided by the literature on the nitrate assimilation under anaerobic conditions. Investigations carried out on nitrate assimilation under anoxia have revealed that this pathway is partially or completely blocked at the nitrite reductase (NiR) step (Ferrari and Varner, 1971; Lee, 1978; Reggiani et al., 1985). Thus, the anaerobic nitrite production (Jaworski, 1971) or the loss of endogenous nitrate (Gray and Cresswell, 1984) has been used for the nitrate reductase (NR) *in situ* assay in different plant organs (Ferrari et al., 1973).

Assimilation of nitrate under anaerobic conditions was previously reported (Kemp and Small, 1993; Fan, 1994). In rice, it has been established that nitrate ions are assimilated by the rice coleoptile during the anaerobic germination (Reggiani et al., 1993, 1995). The enzymes for the nitrate reduction and further ammonia assimilation (NR, NiR, glutamine synthetase and ferredoxin-dependent glutamate synthase) are anaerobically expressed in the coleoptile (Mattana et al., 1994a,b; Mattana et al., 1996). However, the rice coleoptile of anaerobically germinating seeds is a highly tolerant tissue, being able to elongate in an oxygen-deprived environment (Opik, 1973). Most plant tissues (i.e. roots) are instead anoxia intolerant. The present study seeks to establish the capacity of anaerobic nitrate assimilation in rice seedlings germinated in air (seedlings with coleoptile and root) and then transferred to anoxia. This allows to compare the anaerobic nitrate assimilation in an intolerant tissue, such as rice root, with that described in the tolerant anaerobic rice coleoptile (Reggiani et al., 1995).

Materials and methods

The dehulled seeds of rice (*Oryza sativa* var. Arborio) were sterilized for 2 min in 70% (v/v) ethanol and for 30 min in 5% (w/v) calcium hypochlorite, each treatment followed by rinsing with distilled water. The seeds were germinated for 3 days on a sterile wet paper in Petri dishes at 29°C in the dark. Three-day-old seedlings were then placed in jars (500 seedlings/250 ml) and made anaerobic by flushing nitrogen gas (99.99% N₂) through the medium. After 3 h (in order to reach a complete anaerobic condition), 1 mM K¹⁵NO₃ was added for 24 h and the coleoptiles and roots collected for analysis. Perchloric acid extracts and amino acid fractions were obtained as previously described (Reggiani et al., 1993). The amino acid analysis by HPLC of the o-phthalaldehyde (OPA) derivatives was performed according to the method of Jarrett et al. (1986) modified by Reggiani et al. (1995).

The amino acid fraction was freeze-dried before derivatization for gas chromatography-mass spectrometry (GC-MS) analysis. Amino acids were derivatized with N-methyl-(*tert*.-butyldimethylsilyl)-trifluoroacetamide (MTBSTFA) (Mawhinney et al., 1986; Reggiani et al., 1995). Derivatized samples were injected into a Hewlett-Packard 5985 B GC-MS apparatus operating in electron impact (EI) mode. GC analysis was performed as previously reported (Reggiani et al., 1995). Mass spectral data were obtained under the following conditions: ionizing electron energy, 70 eV; emission current, 0.3 mA; ion source temperature, 200°C; scanning rate, 1.5 scan/s over the mass range *m/z* 50–600. The amino acid derivatives possess characteristic mass spectra with intense diagnostic ions, generally at (M-57)⁺, which make them useful for selected ion monitoring (SIM)-GC-MS experiments (Patterson et al., 1993). The incorporation of ¹⁵N into amino acids was calculated by integration of the areas obtained for selected ions for both labeled and unlabeled amino acids and expressing the data as atom % excess (Robinson et al., 1991). ¹⁵N incorporation was considered relevant for amino acids exhibiting atom % excess >1. Incorporation of ¹⁵N into amino acids in the rice coleoptile of anaerobic germinating seeds (thereafter named “anaerobic coleoptile”) were obtained by elaborating data from Reggiani et al. (1995).

Results and discussion

In Table 1 are shown the amino acid levels in aerobic and anaerobic rice tissues. Anoxia determined a significant increase in the level of total amino

Table 1. Effect of 1mM K¹⁵NO₃ on the amino acid content in coleoptile and root of 3-day-old aerobic rice seedlings subjected for 24h to anaerobic conditions

Amino acid	Coleoptile			Root		
	Aerobiosis ¹	Anaerobiosis		Aerobiosis	Anaerobiosis	
		–K ¹⁵ NO ₃	+K ¹⁵ NO ₃ (μmol/g FW)		–K ¹⁵ NO ₃	+K ¹⁵ NO ₃
Ala	1.77 ± 0.12	6.53 ± 0.69	7.33 ± 0.37	1.94 ± 0.17	4.59 ± 0.18	5.40 ± 0.10
Gaba	0.37 ± 0.03	1.03 ± 0.03	1.17 ± 0.05	0.54 ± 0.04	1.81 ± 0.04	2.03 ± 0.08
Glu	1.74 ± 0.13	0.50 ± 0.02	0.78 ± 0.05	2.15 ± 0.19	0.70 ± 0.02	0.87 ± 0.03
Gln	0.97 ± 0.08	0.48 ± 0.01	0.56 ± 0.04	0.76 ± 0.06	0.37 ± 0.02	0.55 ± 0.03
Asp	0.71 ± 0.05	0.12 ± 0.01	0.20 ± 0.02	0.88 ± 0.07	0.20 ± 0.02	0.30 ± 0.03
Asn	1.36 ± 0.10	0.49 ± 0.02	0.81 ± 0.04	1.15 ± 0.11	0.57 ± 0.01	0.83 ± 0.03
Met	0.17 ± 0.02	0.21 ± 0.01	0.29 ± 0.03	0.06 ± 0.01	0.07 ± 0.01	0.06 ± 0.01
Arg	0.10 ± 0.01	0.09 ± 0.01	0.14 ± 0.01	0.40 ± 0.03	0.45 ± 0.02	0.39 ± 0.03
Ser	0.75 ± 0.08	0.81 ± 0.05	1.00 ± 0.06	0.67 ± 0.06	0.76 ± 0.06	0.81 ± 0.05
His	0.07 ± 0.01	0.06 ± 0.01	0.08 ± 0.01	0.15 ± 0.02	0.17 ± 0.02	0.17 ± 0.01
Gly	0.17 ± 0.02	0.23 ± 0.03	0.27 ± 0.03	0.30 ± 0.02	0.34 ± 0.02	0.34 ± 0.02
Thr	0.31 ± 0.02	0.38 ± 0.03	0.40 ± 0.03	0.38 ± 0.04	0.43 ± 0.05	0.50 ± 0.06
Tyr	0.10 ± 0.01	0.08 ± 0.01	0.35 ± 0.02	0.19 ± 0.02	0.22 ± 0.01	0.30 ± 0.03
Val	0.32 ± 0.03	0.39 ± 0.02	0.65 ± 0.04	0.20 ± 0.01	0.23 ± 0.01	0.39 ± 0.02
Lys	0.31 ± 0.02	0.29 ± 0.02	0.46 ± 0.03	0.61 ± 0.05	0.69 ± 0.04	0.86 ± 0.04
Phe	0.09 ± 0.01	0.10 ± 0.01	0.17 ± 0.01	0.15 ± 0.01	0.17 ± 0.03	0.14 ± 0.02
Ile	0.10 ± 0.01	0.12 ± 0.02	0.14 ± 0.01	0.26 ± 0.03	0.29 ± 0.02	0.31 ± 0.03
Leu	0.19 ± 0.02	0.23 ± 0.03	0.26 ± 0.02	0.50 ± 0.04	0.57 ± 0.06	0.55 ± 0.05
Total	9.60	12.14	15.06	11.29	12.63	14.80

¹ Amino acid levels before the imposition of the stress. These levels did not show significant changes after 24 h in well-aerated conditions. Values are the mean of three replicates ±SE. Proline is not detectable as OPA derivative.

acids and, in particular, of alanine and Gaba, while a decrease in the level of glutamate, aspartate and their amides was observed. In coleoptile and root transferred for 24h in anaerobiosis, nitrate induced an increase in total amino acids by 24% and 17% in coleoptile and root, respectively (Table 1). Independently from the nitrate treatment, about 50% and 36% of total amino acids in coleoptile and root, respectively, was alanine. In anaerobic rice roots, an increase in total amino acids and alanine in the presence of nitrate was previously described (Reggiani et al., 1985). On the contrary, in the coleoptile of anaerobically germinated rice seeds, the supply of nitrate led to a decrease in total amino acids and alanine while increased the level of Gaba, arginine and glutamine (Reggiani et al., 1995). In coleoptile and root of seedlings transferred to anoxia, the level of almost all the other amino acids was increased by the nitrate treatment (Table 1).

The incorporation of ¹⁵NO₃[–] into amino acids in 3-day-old seedlings transferred for 24h in anaerobic conditions was analysed using SIM-GC-MS (Table 2). In aerobic seedlings transferred to anoxia, the incorporation of ¹⁵N into glutamate and glutamine (first products of NH₄⁺ assimilation through the glutamine synthetase/glutamate synthase cycle) was significantly higher than

Table 2. Incorporation of ^{15}N into amino acids after treatment with 1 mM K^{15}NO_3 of 3-day-old aerobic seedlings transferred for 24 h in anaerobic conditions

Amino acid ¹	Anaerobic coleoptile ²		Transferred coleoptile		Transferred root	
	Atom % excess	nmol/g FW	Atom % excess	nmol/g FW	Atom % excess	nmol/g FW
Ala	1.2	47.1	3.3	240.4	1.9	102.6
Gaba	5.1	44.0	1.5	17.1	1.5	31.1
Glu	2.8	3.6	14.9	116.1	13.2	114.8
Gln	1.7	5.4	14.5	81.4	5.7	31.2
Asp	2.3	1.7	3.9	7.7	5.3	15.8
Asn	<1		2.1	17.0	<1	
Arg	5.0	36.8	3.7	5.2	<1	
Met	2.5	10.5	1.7	5.0	1.8	1.1
Total		102.0		489.9		296.6

¹Amino acid showing atom % excess <1 were omitted, ²data for coleoptile of anaerobically germinating rice seeds were elaborated from Reggiani et al. (1995).

in the coleoptile of anaerobically germinated seeds. The final product of the glutamine synthetase/glutamate synthase cycle is glutamic acid. We have estimated, elaborating data on the amino acid content in the anaerobic coleoptile (Reggiani et al., 1995), that the level of glutamate in this last tissue was similar to that found in coleoptile and root of seedlings transferred to anoxia (Table 1). As a consequence of this, the higher label incorporation into this amino acid would suggest that, in transferred seedlings, the assimilation of nitrate nitrogen is faster.

The incorporation of label into alanine resulted higher in coleoptile and root of transferred seedlings than in the anaerobic coleoptile (5.1 and 2.2 times higher in coleoptile and root, respectively). Under anoxia, alanine is continuously synthesized from glutamate through a reaction catalyzed by alanine aminotransferase (Reggiani et al., 1988). The reasons of this are the greater pyruvate availability for transamination and the utilization of α -ketoglutarate in NH_4^+ (re)assimilation (Reggiani et al., 1988). Considering that, in tissues subjected to anaerobiosis, pyruvate is made available by the stimulation of glycolysis, it could be suggested that a faster fermentation in seedlings transferred to anoxia than in anaerobically germinated seeds occurred. Aspartate, which derives by transamination from glutamate, resulted more labeled in transferred tissues than in the anaerobic coleoptile (Table 2). This may be a consequence of the higher ^{15}N incorporation into glutamate. However, in transferred seedlings, the incorporation of label into aspartate was lower in coleoptile than in root and, in the former, incorporation into asparagine was detected. Thus, in this tissue a part of aspartate might be utilized for the synthesis of asparagine. The incorporation into Gaba, arginine and methionine was lower in coleoptile and root of transferred seedlings than in the anaerobic coleoptile (Table 2). The first two amino

acids belong to the glutamate family while the latter belongs to the aspartate family (Singh and Matthews, 1994). The lower label incorporation into these three amino acids led, as a consequence, that almost all the ^{15}N in seedlings transferred to anoxia was recovered into alanine, glutamate and glutamine (89% and 84% in coleoptile and root, respectively). This is different from that found in the coleoptile of anaerobically germinated rice seeds in which the ^{15}N into alanine, glutamate and glutamine accounted for only 55% of total amino acid labelling (Reggiani et al., 1995). The data here presented would indicate that, in aerobic seedlings subjected to an anaerobic stress, the metabolism of nitrate nitrogen is almost all described in terms of alanine synthesis. The faster fermentation above hypothesized for these seedlings in comparison with those anaerobically germinated would produce both ATP and NADH for nitrate assimilation and pyruvate for transamination.

The complete assimilation of nitrate ions by an anoxia-intolerant tissue such as the rice root indicates that the pathway of assimilation is not blocked at the NiR step as suggested by some authors (Ferrari and Varner, 1971; Lee 1978). This leads to the conclusion that the *in situ* assay for NR, based on the nitrite production, is an underestimation of the NR activity of the tissue.

References

- Fan TWM (1994) Nitrate metabolism maintains energy production in anaerobic rice coleoptiles. Proceedings of the Fifth International Symposium on Genetics and Molecular Biology of Plant Nutrition, Davis, CA, pp 71–72
- Ferrari TE, Varner JE (1971) Intact tissue assay for nitrate reductase in barley aleurone layers. *Plant Physiol* 47: 790–794
- Ferrari TE, Yoder OC, Filner P (1973) Anaerobic nitrite production by plant cells and tissues. Evidence for two nitrate pools. *Plant Physiol* 51: 423–431
- Gray VM, Cresswell CF (1984) Nitrite utilization by excised *Zea mays* L. roots under anaerobic conditions. *Plant Sci Lett* 33: 31–38
- Jarrett HW, Cooksy KD, Ellis B, Anderson JM (1986) The separation of o-phthalaldehyde derivatives of amino acids by reverse-phase chromatography on octylsilica columns. *Anal Biochem* 153: 189–198
- Jaworski EG (1971) Nitrate reductase assay in intact plant tissues. *Biochem Biophys Res Commun* 43: 1274–1279
- Kemp K, Small JGC (1993) Nitrate and nitrate reductase in *Erythrina caffra* seeds: enhancement of induction by anoxia and possible role in germination. *Planta* 189: 298–300
- Lee RB (1978) Inorganic nitrogen metabolism in barley roots under poorly aerated conditions. *J Exp Bot* 29: 693–708
- Malavolta E (1954) Study on the nitrogenous nutrition of rice. *Plant Physiol* 29: 98–99
- Mattana M, Bertani A, Reggiani R (1994a) Expression of glutamine synthetase during the anaerobic germination of *Oryza sativa* L. *Planta* 195: 147–149
- Mattana M, Coraggio I, Bertani A, Reggiani R (1994b) Expression of the enzymes of nitrate reduction during the anaerobic germination of rice. *Plant Physiol* 106: 1605–1608
- Mattana M, Coraggio I, Brambilla I, Bertani A, Reggiani R (1996) Nitrate assimilation during the anaerobic germination of rice: expression of ferredoxin-dependent glutamate synthase. *Planta* 199: 74–78

- Mawhinney TP, Robinett RSR, Atalay A, Madson MA (1986) Analysis of amino acids as their tert.-butyldimethylsilyl derivatives by gas-liquid chromatography and mass spectrometry. *J Chromatogr* 358: 231–242
- Opik H (1973) Effect of anaerobiosis on respiration rate, cytochrome oxidase activity and mitochondrial structures in coleoptile of rice (*Oryza sativa* L.). *L Cell Sci* 12: 725–739
- Patterson BW, Carraro F, Wolfe RR (1993) Measurement of ^{15}N enrichment in multiple amino acids and urea in a single analysis by gas chromatography/mass spectrometry. *Biol Mass Spectrom* 22: 518–523
- Reggiani R, Bertini F, Mattana M (1995) Incorporation of nitrate nitrogen into amino acids during the anaerobic germination of rice. *Amino Acids* 9: 385–390
- Reggiani R, Brambilla I, Bertani A (1985) Effect of exogenous nitrate on anaerobic metabolism in excised rice roots. I. Nitrate reduction and pyridine nucleotides pools. *J Exp Bot* 36: 1193–1199
- Reggiani R, Cantù CA, Brambilla I, Bertani A (1988) Accumulation and interconversion of amino acids in rice roots under anoxia. *Plant Cell Physiol* 29: 981–987
- Reggiani R, Mattana M, Aurisano N, Bertani A (1993) Utilization of seed storage nitrate during the anaerobic germination of rice. *Plant Cell Physiol* 34: 379–383
- Robinson SA, Slade AP, Fox GG, Phillips R, Ratcliffe RG, Stewart GR (1991) The role of glutamate dehydrogenase in plant nitrogen metabolism. *Plant Physiol* 95: 509–516
- Singh BK, Matthews BF (1994) Molecular regulation of amino acid biosynthesis in plants. *Amino Acids* 7: 165–174
- Yamasaki T, Seino K (1965) Use of nitrate fertilizers for the cultivation of paddy rice (part 1). About the physiological character of rice seedlings supplied with nitrate as the source of nitrogen. *J Sci Soil Manure Japan* 36: 153–158

Authors' address: Dr. R. Reggiani, Istituto Biosintesi Vegetali, C.N.R., via Bassini 15, I-20133 Milano, Italy.

Received November 4, 1996