

The proline biosynthesis in living organisms

Review Article

B. Aral and P. Kamoun

Laboratoire de Biochimie Médicale B and Centre National de la Recherche Scientifique (CNRS) URA1335, Hôpital Necker-Enfant Malades, Paris, France

Accepted May 28, 1997

Summary. In this article we review recent work on the physiology of proline and Δ^1 -pyrroline-5-carboxylate (P5C) in living organisms and consider recent progress in our understanding of the role of P5C synthetase in collagen metabolism and the regulation of urea cycle in vertebrates. Much of this recent progress has been made possible by advances in our knowledge of the enzymes and genes involved in proline biosynthesis in man. The availability of well characterized P5C synthetase deficiency in man has been an impetus for the cloning of the cDNA encoding for this enzyme from man and facilitated the establishment of the phenotype-genotype relationships in P5C synthetase deficiency in higher vertebrates.

Keywords: Proline – Δ^1 -pyrroline-5-carboxylate synthetase – cDNA – Collagen – Urea cycle

Abbreviations: GK: γ -glutamyl kinase; GPR: γ -glutamyl phosphate reductase; P5CR: Δ^1 -pyrroline-5-carboxylate reductase; GSA: glutamic- γ -semialdehyde; P5C: Δ^1 -pyrroline-5-carboxylate; P_i : Inorganic phosphate; AMP, ADP, ATP: Adenosine 5'-mono-, di-, triphosphate; NAD^+ , NADH: nicotinamide adenine dinucleotide, and its reduced form; $NADP^+$, NADPH: nicotinamide adenine dinucleotide phosphate, and its reduced form; DEAE: diethylaminoethyle; OAT: ornithine amino transferase; CHO: Chinese hamster ovary; IGF-1: insulin-like growth factor-1; P5CDH: pyrroline 5-carboxylate dehydrogenase; IMP: inosine 5'-monophosphate.

Introduction

Proline represents a unique class of molecules among amino acids; besides being classified as a non-essential amino acid it belongs to the class of “glutamate family” of amino acids whose biosynthesis depends on the ability of the carbon skeleton of glutamic acid to give rise to glutamine, proline,

arginine, ornithine (Conn et al., 1987). In fact, a specific system of enzymes with special properties has evolved to mediate the metabolism of proline. These metabolic interconversions among proline, ornithine, and glutamate have pyrroline-5-carboxylic acid (P5C) as the sole common intermediate. As pointed out by Adams, the inclusion of the α -amino group within a ring structure bestows a number of special properties (Adams, 1970). With its peptide bond within the pyrrolidone ring, proline confers rigidity and three-dimensional stability to proteins. The high content of proline and hydroxyproline in collagen contributes to the unique physical properties of this structural protein. Moreover, in several unicellular species, the intracellular concentration of proline serves as a defense against osmotic challenge (Csonka, 1989; Csonka and Hanson, 1991; Strom et al., 1983; Yancey, 1994). Also as it has been reviewed extensively elsewhere, the accumulation of proline is a common metabolic response of higher plants to water deficits, salinity stress and cold stress (Csonka and Baich, 1983; Delauney and Verma, 1993; Dorffling et al., 1993; Hanson and Hitz, 1982; Stewart, 1981; Stewart and Larher, 1980; Thomson, 1980). In this article we review recent work on the metabolic functions of proline and P5C unraveled by the results of molecular studies from their deficiencies in humans.

I. Pathways of synthesis of P5C and proline

Nutritional and other studies with isotopic carbon have provided strong evidence for the interconversion in vivo of glutamic acid and proline in both animal tissues and microorganisms (Stetten, 1951). The metabolic pathways with P5C as the central intermediate (Fig. 1) were initially defined as a system for proline biosynthesis (Adams, 1970; Adams and Frank, 1980; Baich, 1969) and this physiologic endpoint has been emphasized by a number of investigators (Mezl and Knox, 1977; Smith and Phang, 1978; Yip and Knox, 1972). In both prokaryotes (Baich, 1969) and eukaryotes (Baich, 1977; Smith and Phang, 1979; Valle et al., 1973) the inability to synthesize P5C results in proline auxotrophy. On the other hand if one of the pathways for P5C synthesis is present, i.e. from either ornithine or glutamate, normal growth occurs in proline-free medium (Smith and Phang, 1979).

a. In bacteria

Enzyme systems catalyzing the reduction of this intermediate to proline have been described in microorganisms (Meister et al., 1957). In bacteria such as *E. coli*, *Pseudomonas aeruginosa* and *Salmonella typhimurium*, proline synthesis from glutamate is catalyzed by three enzymes: γ -glutamyl kinase (GK), γ -glutamyl phosphate reductase (GPR) (also called glutamic semialdehyde dehydrogenase) and Δ^1 -pyrroline-5-carboxylate reductase (P5CR) (Fig. 2), encoded by genes *proB*, *proA* and *proC*, respectively (Adams and Frank, 1980; Baich, 1969, 1971; Csonka and Baich, 1983; Deutch et al., 1982, 1984; Gamper and Moses, 1974; Hayzer and Leisinger, 1980, 1981, 1982; Hayzer and




Fig. 1. Metabolic pathways illustrating the relationships between proline, ornithine and citrulline biosynthesis and the urea cycle substrates. (1) Δ^1 -pyrroline-5-carboxylate synthetase (P5CS); (2) Δ^1 -pyrroline-5-carboxylate reductase; EC 1.5.1.2; (3) Δ ornithine amino-transferase (OAT); EC 2.6.1.13; (4) ornithine trans-carbamoylase (OTC); EC 2.1.3.3; P5C Δ^1 -pyrroline-5-carboxylate

Moses, 1978a,b; Krishna and Leisinger, 1979; Krishna et al., 1979; Smith et al., 1984). The step from glutamic- γ -semialdehyde (GSA) to Δ^1 -pyrroline-5-carboxylate (P5C) is reversible and spontaneous. The reaction catalysed by GK is irreversible. The product of the reaction, γ -glutamyl phosphate, is extremely labile. It can be chemically trapped by reaction with hydroxylamine to yield the more stable γ -glutamyl hydroxamate which can be quantified with a ferric chloride reagent (Baich, 1969; Hayzer and Leisinger, 1980). In the absence of hydroxylamine, the γ -glutamyl phosphate product spontaneously degrades to pyroglutamate (5-oxoproline) and inorganic phosphate (P_i) (Seddon et al., 1989). By using two proline-requiring mutants of *E. coli*, Strecker has shown that the addition of pyruvate, lactate, glucose, or formate increased the formation of P5C in washed cells in the presence of either L-glutamate or L-glutamine; further stimulation was obtained on the addition of either AMP or ADP but not ATP. The production of P5C was inhibited by the endogenous proline (Strecker, 1960b) Δ^1 -pyrroline-5-carboxylic acid hydrochloride has been synthesized and isolated and assayed for its conversion to proline in 80% yield by chemical reduction. Solutions of the compound are unstable both at room temperature and in the frozen state but are relatively stable at 3°C; depending on the pH of the reaction, reaction with ninhydrin results in the formation of two products; the 2,4-dinitrophenylhydrazone derivative and the addition product with o-aminobenzaldehyde have been prepared (Strecker, 1960a). The GK of *E. coli* is inhibited by proline (Baich, 1969; Rushlow et al., 1984; Smith, 1985) and has a relatively low affinity for glutamate ($K_m = 7\text{--}10\text{ mM}$) (Baich, 1969). Marked substrate activation of the enzyme is observed when the concentration of glutamate exceeds 50 mM (Baich, 1969). Proline increases the apparent K_m for glutamate (Baich, 1969). Proline overproducing mutants have a GK with markedly altered feedback inhibition (Csonka, 1981; Rushlow et al., 1984; Smith, 1985). The reaction




Fig. 2. Pathways of proline biosynthesis in different species. The enzymes are: *A* γ-glutamyl kinase (γ-GK) (EC 2.7.2.11); *B* γ-glutamyl phosphate reductase (γ-GPR) (EC 1.2.1.41); *AB* P5C synthetase (P5CS); *C* non-enzymatic step; *D* P5C reductase (EC 1.5.1.2)

catalysed by the second enzyme is reversible and this reversibility forms the basis of a selective assay for GPR in which P5C-GSA-dependent, and P_i -dependent reduction of $NADP^+$ is measured. The γ-glutamyl phosphate generated in this reverse reaction spontaneously degrades to pyroglutamate and P_i (Hayzer and Leisinger, 1981). Pyroglutamate has the following distinctive attributes: (i) unlike other amino acids it does not bind to Dowex-50- H^+ (Hayzer and Leisinger, 1981); (ii) it is ninhydrin negative (Hayzer and Leisinger, 1981); and (iii) it can be converted to glutamate by boiling in aqueous acid (Hayzer and Leisinger, 1981; Mazelis, 1979). The assay for GPR ($P5C \rightleftharpoons GSA$ - and P_i -dependent reduction of $NADP^+$) is subject to interference by an active P5CR in crude extracts (Hayzer and Leisinger, 1980). Thus, as $NADPH$ is formed, this can be oxidized back to $NADP^+$ by the catalytic activity of P5CR. This interference by P5CR can lead to markedly non-linear $NADP^+$ reduction kinetics in bacterial crude extracts. However, GPR and

P5CR can be separated easily on a diethylaminoethyl (DEAE)-cellulose column (Hayzer and Leisinger, 1980).

b. In yeast

The yeast *Saccharomyces cerevisiae* synthesizes proline from glutamate via the intermediates γ -glutamyl phosphate, γ -glutamyl semialdehyde, and P5C (Brandriss, 1979; Brandriss and Falvey, 1992; Orser et al., 1988; Tomenchok and Brandriss, 1987). The isolation and preliminary characterization of mutations leading to proline auxotrophy in the yeast *S. cerevisiae* have been reported (Brandriss, 1979). Three complementation groups (pro1, pro2, and pro3) were identified. The requirement for proline in the pro1 and pro2 strains could be satisfied by arginine or ornithine, since these compounds can be converted to glutamate semialdehyde by enzymes in the arginine degradation pathway, bypassing the early biosynthetic steps. pro3 mutants can grow only when proline is added to the medium and are defective in P5C reductase enzyme activity. The three enzymes involved in this pathway are γ -glutamyl kinase (the product of the PRO1 gene), γ -glutamyl phosphate reductase (the product of the PRO2 gene), and P5C reductase (the product of the PRO3 gene). The gene-enzyme relationships were demonstrated by interspecies complementation of proB, proA, and proC mutants of *E. coli* and *Salmonella typhimurium* by the yeast PRO1, PRO2, and PRO3 genes, respectively (Tomenchok and Brandriss, 1987), and of a yeast pro1 mutant by the *E. coli* proB gene (Orser et al., 1988). The first two enzyme activities were never measured in the studies of *S. cerevisiae*. In *S. cerevisiae*, the enzyme P5C reductase also functions in the pathway for arginine degradation (Brandriss and Magasanik, 1980). The ornithine that is formed from arginine is converted to glutamate semialdehyde and P5C in the cytoplasm. P5C is then converted to proline by P5C reductase, and the proline is transported into the mitochondria, where it is converted to glutamate (Brandriss and Magasanik, 1981). Therefore, P5C reductase functions in both the proline biosynthetic and the arginine degradative pathways.

c. In higher plants

Enzyme systems catalyzing the reduction of the glutamate to proline have been described in plants. (Bogges et al., 1976; Morris et al., 1969; Heyser et al., 1989a; Heyser et al., 1989b; Rhodes and Handa, 1989; Rhodes et al., 1986). A number of in vivo labelling studies with ^{14}C -labelled precursors (Bogges et al., 1976; Morris et al., 1969) or $[^{13}\text{C}]$ glutamate (Heyser et al., 1989a,b) or $^{15}\text{NH}_4^+/\text{NO}_3^-$ (Rhodes and Handa, 1989; Rhodes et al., 1986) strongly suggest that glutamate is a major precursor of osmotic stress-induced proline accumulation in plants and that osmotic stress results in an increase of proline biosynthesis rate (Bogges et al., 1976; Rhodes and Handa, 1989; Rhodes et al., 1986; Stewart, 1981). The prevailing evidence [reviewed extensively by others (Delauney and Verma, 1993; Hanson and Hitz, 1982; Stewart, 1981;

Thompson, 1980)] is that the increase in proline synthesis rate may, in part, involve induction or activation of enzymes of proline biosynthesis, possibly coupled with a relaxation of proline feedback inhibition control of the pathway (Bogges et al., 1976; Delauney and Verma, 1993; Stewart, 1981). Interestingly, this feedback control of the proline pathway may also be modulated during development (Oaks, 1992; Oaks et al., 1970). Decreased proline oxidation to glutamate (Elthon and Stewart, 1982; Huang and Cavalieri, 1979; Stewart et al., 1977), decreased utilization of proline in protein synthesis (Bogges and Stewart, 1980; Stewart, 1981) and enhanced protein turnover (Fukutoku and Yamada, 1984) may all contribute to net proline accumulation. There are indications that *de novo* transcription and translation are required for stress-induced proline biosynthesis in higher plants. For example, stress-induced proline accumulation is inhibited by both cycloheximide and cordycepin in *Arabidopsis* (Verbruggen et al., 1993). In contrast, because cycloheximide is inhibitory to 200mM NaCl-induced proline accumulation in *D. spicata* cells, whereas actinomycin D is not inhibitory, Ketchum et al. (1991) suggest that translation, but not transcription, is necessary for production of proline in stressed cells.

In addition to the pathway of synthesis of proline from glutamate, there is also the possibility of synthesis of proline from ornithine (derived from the N-acetyl-glutamate pathway) in higher plants. This can occur by two routes, depending upon whether the α - or δ -NH₂ moiety of ornithine is transaminated (Adams and Frank, 1980; Csonka and Baich, 1983; Delauney and Verma, 1993; Mestichelli et al., 1979; Thompson, 1980). In the case of δ -NH₂ transamination, the product is GSA which, in spontaneous equilibrium with P5C, can be converted readily to proline by P5CR. Ornithine δ -transaminase has been partially purified from several plant sources, including pumpkin cotyledons (Splittstoesser and Fowden, 1973). The alternative pathway involves α -NH₂ transamination of ornithine to α -keto- δ -aminovalerate $\rightleftharpoons$ Δ^1 -pyrroline-2-carboxylate (P2C). This product can be reduced to proline by a P2C reductase. In vivo tracer studies with labelled ornithine strongly suggest that the P2C pathway of ornithine metabolism is the main route of conversion of ornithine to proline in several plant species (Mestichelli et al., 1979). There is a paucity of in vitro enzymological data to support the occurrence of a glutamate pathway of proline synthesis in higher plants similar to that established for bacteria. Although P5CR has been identified and characterized in several plant species (e.g. Krueger et al., 1986; LaRosa et al., 1991; Treichel, 1986), there have been no reports of GK or GPR activities in higher plants since the first report of the occurrence of in vitro synthesis of ¹⁴C GSA from ¹⁴C glutamate, Mg²⁺, ATP and NADPH in cell-free extracts of sugar beet (Morris et al., 1969). The enzyme activity described by Morris et al. (1969) is similar to that reported for the P5C synthetase activity of mammalian cells (Lodato et al., 1981; Wakabayashi and Jones, 1983). Efforts devoted to identifying a GK activity in plant extracts by the γ -glutamyl hydroxamate assay procedure have been hindered by a large background of γ -glutamyl hydroxamate formation activity catalysed by glutamine synthetase in plant tissues (Mifflin and Lea, 1977). There does not appear to have been a sys-

tematic search for GPR in higher plants. In addition to the problem of interference from P5CR in detecting GPR activity in crude extracts there is also an interference from an enzyme capable oxidizing P5C to glutamate; P5C:NAD(P)⁺ oxidoreductase. This enzyme is the second enzyme of the two-step pathway of proline oxidation in higher plants (Elthon and Stewart, 1982; Huang and Cavalieri, 1979; Rayapati and Stewart, 1991; Stewart, 1981; Stewart and Boggess, 1978; Thompson, 1980). These problems associated with measurement of GPR are alleviated by monitoring P_i-dependent pyroglutamate production from P5C $\rightleftharpoons$ GSA and NADP⁺ instead of monitoring NADP⁺ reduction to NADPH.

d. In animals

Enzyme systems catalyzing the reduction of the P5C to proline have been described in rat and other animal tissues (Meister et al., 1957). Workers showed that an enzyme activity converting glutamate to P5C is present in homogenates of intestinal mucosa incubated with [U-¹⁴C] glutamate (Ross et al., 1978). The conversion of glutamate to P5C is most likely catalyzed by two enzymes or a multifunctional enzyme complex. The γ carbon of glutamate is activated by ATP followed by the NADPH-dependent reduction of γ -glutamyl phosphate. As a trivial designation, "P5C synthetase" refers to the cell-free activity in which glutamate is converted to P5C in an ATP and NADPH-dependent reaction. The activity was measured in cultured cells and its presence or absence correlated with proline prototrophy or auxotrophy, respectively (Smith et al., 1980). Using solubilized mitochondrial preparation from rat intestinal mucosa, Wakabayashi et al. characterized this activity in greater detail (Wakabayashi et al., 1983). The activity is sensitive to inhibition by ornithine (Lodato et al., 1981). And unlike the bacterial enzyme, the conversion of glutamate to P5C in mammalian tissues is not inhibited by proline (Kramer et al., 1985). Interestingly, in rat tissues the enzyme has been detected only in intestine and thymus (Wakabayashi et al., 1983). However, the presence of the pathway in a variety of cultured cells is supported by metabolic studies (Lorans et al., 1978; Smith and Phang, 1979; Valle et al., 1973) as well as by direct enzyme assay in cell-free extracts (Smith et al., 1980). Derepression or activation of the enzyme accompanying mitogenesis is an intriguing possibility. Mutants of Chinese hamster fibroblasts which are proline hyperproducers and thereby resistant to the toxic effects of azetidine carboxylic acid have either an extremely high level of P5C synthetase or an enzyme which is insensitive to inhibition by ornithine (Smith et al., 1981; Lodato et al., 1981). The extent and the rate of inactivation depend upon the time and the temperature of prior incubation of the enzyme with ornithine in the absence of substrates, which is suggestive evidence for an allosteric regulation of the enzyme; the inactivation was prevented by 0.2 mM ATP or ADP (Wakabayashi et al., 1985).

Although demonstrable *in vitro* (McGivan et al., 1977), the physiological significance of the carbon transfers between the urea cycle and glutamatic acid-P5C-proline metabolism was in doubt. Widely distributed in mam-

malian tissues and localized to the mitochondrial matrix, ornithine aminotransferase catalyzes the ω -transamination of ornithine to α -ketoacid acceptors thereby forming $P5C \rightleftharpoons GSA$. The K_{eq} of hepatic OAT greatly favors $P5C$ formation. The properties of the purified enzyme have been recently reviewed (Valle and Simell, 1995). Animals fed a high protein diet have markedly elevated levels of OAT activity in liver (Volpe et al., 1969). Moreover, recent studies in patients with gyrate atrophy of the choroid and retina, an inherited disorder with deficiency of OAT (Valle et al., 1977; Valle and Simell, 1995) strongly supports this idea (Valle et al., 1980). The deficiency of OAT results in markedly elevated plasma ornithine, low ammonia, and low glutamate and glutamine, findings which are consistent with the interpretation that OAT controls the level of ornithine available to the urea cycle. In order to prove the importance of the ornithine pathway, Smith and Phang have selected mutant Chinese hamster ovary (CHO) cells with specific defects in the proline metabolic pathways and have found that ornithine can be converted to proline at a rate adequate to support maximal cell growth in the absence of any other sources of proline; although the importance and net direction of the OAT reaction in cultured CHO fibroblasts was in the direction $ornithine \Rightarrow P5C \Rightarrow glutamate$, the OAT-catalyzed reaction is reversible and net flow from glutamate or proline to ornithine may occur in other cell types (Smith and Phang, 1979). Glucagon which induces the hepatic enzyme may be responsible, in part, for the elevated levels of OAT in liver in response to high protein diet. Liver OAT would be inducible not by glucagon but estrogens (Lyons and Pitot, 1976). In fact, transcript quantification experiments showed a largely transcriptional-independent mechanism for the increase in enzyme synthesis with increased dietary protein (Mueckler et al., 1983). The combination of these two activities may supply urea cycle intermediates for the liver. The back reaction by ornithine aminotransferase, i.e. the synthesis of ornithine from $P5C$, clearly can occur in intestine. Ross et al. obtained ornithine synthesis from glutamate in rat intestinal mucosa homogenates: in their studies, ^{14}C -labeled ornithine was separated from other radioactive chemicals by ion-exchange chromatography, and was specifically identified by its capacity to serve as a substrate for OTC. The synthesis of ornithine required ATP, Mg^{2+} and NADPH in addition to glutamate (Ross et al., 1978). Windmueller and Spaeth showed citrulline production from glutamine in isolated perfused rat intestine (Windmueller and Spaeth, 1975). The formation of $[^{14}C]$ ornithine from $[^{14}C]$ glutamate was found in a subcellular fraction enriched in mitochondria of rat small intestinal mucosa; synthesis of ornithine from glutamate was not observed with liver or kidney homogenates; pyridoxal 5'-phosphate stimulated both ornithine synthesis and endogenous OAT activity; partially purified rat liver OAT stimulated ornithine synthesis (Henslee and Jones, 1982). In order to establish the interorgan dependency of arginine synthesis, Wakabayashi et al. quantified $P5C$ synthetase activity in many rat tissues and found that the activity was concentrated only in the upper small intestine. Minor activity was found in pancreas, thymus, lymph node, and some other tissues: this was confirmed by the dependency on the specific substrates, the loss of activity in the pres-

ence of an inhibitor, and identifying the reduced product as proline (Wakabayashi et al., 1991). Riby et al. established the developmental profile of the two intestinal mitochondrial enzymes, P5C synthetase and OAT, responsible for the conversion of glutamate to ornithine. Both enzymatic activities were found to be significantly elevated throughout the suckling period with a peak of activity during the second week of life; in contrast, glutamate dehydrogenase activity in the intestine did not appear to be developmentally regulated during the suckling and weaning periods and ornithine decarboxylase activity was undetectable in the intestine of the mouse during the suckling period and was detected briefly at weaning, indicating that ornithine synthesized in the intestinal mitochondria is not diverted actively into the polyamine pathway and is available for synthesis of arginine by the enzymes of the urea cycle (Riby et al., 1990). Wu et al. showed no evidence of P5C synthetase activity in chick enterocytes therefore resulting in the lack of synthesis of ornithine and citrulline from glutamine, in contrast to cells from post-weaning pigs (Wu et al., 1995ab; Wu, et al., 1994). In the lens, glutamate is necessary for the synthesis of products including glutamine, glutathione, and proline, as well as proteins such as the glutamate-rich crystallins. The transamination reactions in human lenses appeared to preferentially involve amino groups derived from glutamate; but the glutamate used for human lens proline biosynthesis is derived from glutamine and glutaminase, which converts glutamine to glutamate, is located in mitochondria in all tissues examined as well as in rat lens, suggesting that the glutamate at the site where proline synthesis begins (i.e. in mitochondria) may arise primarily from deamidation of glutamine by glutaminase (Jernigan, 1990).

L-Pyrroline-5-Carboxylic Acid (P5C) has been obtainable by chemical synthesis only as a mixture of the D- and L-stereoisomers. Smith et al. reported a method for the enzymatic synthesis of P5C on a preparative scale. The P5C is formed from L-ornithine by purified OAT and then isolated by Dowex-50W cation-exchange resin chromatography. With [^{14}C]ornithine as precursor, high specific activity uniformly labeled [^{14}C]P5C can be obtained (Smith et al., 1977). Using an assay based on a preparation of P5C reductase purified from genetically engineered *E. coli*, Fleming et al. measured P5C in normal human plasma, urine, and saliva. In addition, they have found pyrroline-5-carboxylate levels $10\times$ to $20\times$ normal in two patients with type II hyperprolinemia. P5C concentrations in normal human plasma were 0.38 ± 0.11 nmol/mL; those of saliva were approximately 50% of plasma levels. Urinary excretion of P5C was up to $1.2\mu\text{mol}/24\text{h}$ in normal subjects (Fleming et al., 1984).

Mixson et al. developed a very sensitive and specific new method of assay based on the interaction of P5C with cysteine. [^{35}S] Cysteine at a defined specific activity is reacted with P5C and the P5C-Cys adduct can be separated from the reactants by cation exchange column chromatography (Mixson, 1991).

P5C reductase, a cytosolic enzyme which catalyzes the conversion of P5C to proline with reduced pyridine nucleotide as cofactor, is ubiquitous in ani-

mal cells. The enzyme has been purified from *E. coli* (Deutch et al., 1982; Rossi et al., 1977) and yeast (Matsuzawa and Ishiguro, 1980) but not from mammalian cells. Nevertheless, characterization of the activity from animal tissues reveals a number of interesting features. Heretofore, investigators have considered that NADH is the preferred cofactor for P5C reductase. But studies in intact cells argue that it is NADPH which is oxidized by the enzyme in situ (Yeh and Phang, 1980; Yeh et al., 1981). The explanation may lie in the kinetic characteristics of the enzyme. Although $V_{\max}$ activities are higher with NADH, preference for NADPH may be due not only to a markedly higher affinity for NADPH but also a higher affinity for P5C with NADPH as cofactor (Yeh et al., 1981). Thus, at least in certain cells, NADPH is oxidized concomitantly with the conversion of P5C to proline. Also of interest P5C reductase from various tissue or cellular sources are differentially sensitive to inhibition by proline (Valle et al., 1973; Yeh et al., 1981), NADP^+ (Smith and Greenberg, 1957; Yeh and Phang, 1981), and adenine nucleotides (Smith and Greenberg, 1957). The enzyme from cultured fibroblasts is sensitive to inhibition by proline and its inhibition of the NADH-mediated reaction ($K_i = 2 \times 10^{-4} \text{M}$) is much greater than of the NADPH-mediated reaction ($K_i = 2 \times 10^{-3} \text{M}$). By contrast, the hepatic and erythrocyte enzymes are insensitive to proline but very sensitive to inhibition by NADP^+ (Yeh and Phang, 1981). Adenine nucleotides also inhibit the hepatic enzyme (Smith and Greenberg, 1957) but not the erythrocyte enzyme. These differential patterns of inhibition suggest that there are isozymes of P5C reductase found in various tissues, an interpretation further supported by the relative deficiency of the NADH-mediated activity in a leukemia cell line (Lorans and Phang, 1981). Whether these putative isozymes are products of different genes or are interconvertible forms is an intriguing question which awaits definitive studies on enzyme prepared from different sources. In addition to the kinetic characteristics and allosteric regulation, the amount of P5C reductase in cells appears to be regulated. In contrast to enzyme systems for amino acid synthesis in prokaryotes P5C reductase in cultured cells is not repressed by high concentrations of proline in the growth medium. However, marked increases in enzyme activity are observed in human peripheral lymphocytes activated with mitogenic lectins (Valle et al., 1975). In mammary gland tissues stimulated to undergo lactogenesis, coordinated increases in arginase, ornithine aminotransferase, and P5C reductase are observed (Yip and Knox, 1972). An enzyme has been purified approximately 160-fold from a calf liver extract catalyzing the reduction of P5C to proline and gave preliminary kinetic data. Most of the P5C reductase activity were associated with the microsomal and supernatant fractions whereas both proline oxidase and P5C dehydrogenase activities are associated with the mitochondrial fraction (Johnson and Strecker, 1962). The reductive reaction cannot be reversed, lending support to the hypothesis that the reduction of glutamate to proline proceeds by a pathway which differs from that for the oxidation of proline to glutamate. Phang et al. developed a new radioisotopic assay in which they measured the radioactivity incorporated into the product, L-proline, from precursor

Δ^1 -pyrroline-5-carboxylate [$U\text{-}^{14}\text{C}$] (Phang et al., 1973). A high P5C reductase activity was also shown in epidermal and neural ectoderm such as cornea, lens, retina and retinal pigment epithelium (Matsuzawa et al., 1982b). Bovine retinal P5C reductase (L-proline: NAD(P)⁺ 5-oxidoreductase, EC 1.5.1.2) was purified 180-fold from crude extract, using affinity chromatography with 2',5'-ADP Sepharose 4B. The purified enzyme utilized either NADH or NADPH as the coenzyme. The enzyme activity was inhibited with ATP, and activated with K⁺ and phosphate ions, suggesting that it may be regulated by the energy level and the redox state in the retina. The distribution of the enzyme activity in bovine cornea and lens tissues have been provided (Shiono et al., 1986; Matsuzawa, 1982b).

In brain for example, the correlation between OAT and P5C reductase was significant ($\gamma = 0.75$). This is because the brain uptake index of proline is as low as those of other putative neurotransmitters (glutamate, aspartate and glycine), the existence of an endogenous proline synthetic pathway favouring the ornithine $\Rightarrow$ P5C $\Rightarrow$ proline pathway in the CNS is suggested (Matsuzawa and Obara, 1987). Yeh et al. (1984) reported a P5C reductase activity in human erythrocytes. Moreover, P5C reductase activity was the only proline metabolic enzyme activity present in these cells; The kinetic properties and regulation of the erythrocyte enzyme were distinctly different from those for human fibroblast P5C reductase (First, the enzyme has a preferential affinity for NADPH; second, it lacked of sensitivity to feedback inhibition by proline and instead this enzyme is sensitive to feedback inhibition by product NADP⁺), suggesting that the function of the enzyme in human erythrocytes may be to generate oxidizing potential in the form of NADP⁺ (Yeh and Phang, 1981). Generally, P5C reductase capacity correlates with that of OAT in the same tissue; this indicates that there is a widespread potential among tissues for proline synthesis from ornithine. In contrast, the significant correlations of the activities of the proline oxidase, P5C dehydrogenase with one another indicate a real but less widespread potential for proline oxidation in rat tissues. In fact, a flow to form ornithine would appear to be distinctly atypical, because proline oxidase activities are limited and are not significantly associated with OAT activities in the same tissues, and because there is no detectable conversion of P5C into glutamate by the reversal of the P5C dehydrogenase reaction in the same tissues (Herzfeld et al., 1977).

Proline oxidase catalyzes the conversion of proline to P5C and transfers electrons into the mitochondrial electron transport with an intervening flavoprotein (Meyer, 1977). Thus, the pair of electrons donated by proline can effect the phosphorylation of two ADP molecules. The enzyme is tightly bound to mitochondrial inner membranes and is limited in its tissue location to liver, kidney, brain and heart (Herzfeld et al., 1977). Treatment of mitochondria with Triton X-100 yields a solubilized preparation but enzyme activity then requires the presence of an electron acceptor such as iodonitrotetrazolium (Kowaloff et al., 1977). The enzyme can be induced by glucocorticoids both in animals (Kowaloff et al., 1976) and in cultured cells

(Kowaloff et al., 1978). The activity is sensitive to inhibition by lactate (Kowaloff et al., 1977): the K_m for proline increases as a function of increasing lactate concentration.

P5C dehydrogenase catalyzes the conversion of P5C to glutamic acid (Adams and Goldstone, 1960). The enzyme is ubiquitous but shows marked differences in specific activity among tissues. Although the activity of P5C dehydrogenase is generally high in tissues with proline oxidase, e.g., liver and kidney, suggesting that for these tissues proline serves as a source of glutamate, the activities of these two enzymes are not always parallel. For example, rat brain has P5C dehydrogenase/proline oxidase ratios which are one-tenth of those for liver or kidney (Yeh and Phang, unpublished observations). Considered to be localized primarily in mitochondria, P5C dehydrogenase is also found in the cytosol. Differential utilization of NAD^+ versus $NADP^+$ may help distinguish enzymes from the two subcellular sources. P5C dehydrogenase isolated from both yeast and animal tissues appears to be sensitive to inhibition by amino acids. In yeast cells the sensitivity of the enzyme to specific amino acids in vitro correlates with the sensitivity to growth inhibition of a mutant strain auxotrophic for glutamate (Lundgren and Ogur, 1973). Similarly, the enzyme isolated from kidney is sensitive to inhibition by a variety of amino acids. At saturating concentrations, these amino acids inhibit in an additive fashion suggesting distinct regulatory sites for each amino acid (Lundgren and Ogur, 1972). Small et al. reported an 800-fold purification P5C dehydrogenase from rat liver mitochondria to yield an essentially homogenous protein with a $M_r = 59,000$. The substrate P5C and NAD^+ have apparent dissociation constant of 0.16 and 1.0 mM, respectively. Studies established that the reverse reaction, the conversion of glutamate and NADH to P5C and NAD^+ , is not catalyzed by this enzyme. Survey of various rat tissues for P5C dehydrogenase activity used $[^3H]P5C$ as substrate, revealed higher activities in liver, kidney and heart, intestinal mucosa, retina and xiphoid cartilage had lowest activities (Small and Jones, 1990). The deficiency of P5C dehydrogenase has been identified as the basis for the inborn error of metabolism, Type II hyperprolinemia (Valle et al., 1976). In these patients, plasma proline concentrations are elevated markedly and P5C is excreted in the urine. Due to the insensitivity of previously available assays for P5C, this molecule was not measurable in the plasma of these patients. But recent studies using a sensitive enzyme-coupled radioisotopic assay has quantitated P5C levels in the plasma of patients with hyperprolinemia II as well as in normal subjects. Plasma P5C in this disease is 15 to 20 times elevated over that of normals (Fleming et al., 1984). An assay which measures radiolabeled glutamate production by P5C dehydrogenase in the presence of NAD^+ from radiolabelled P5C has been developed. Separation of substrate from product is achieved by column chromatography using Dowex-50 cation-exchange resin. The product isolated by this procedure was identified as glutamate (Small and Jones, 1987). P5C dehydrogenase has been purified to homogeneity from the human liver by the use of ion exchange chromatography on CM-Sephadex and affinity chromatography on Blue Sepharose CL-6B and 5'-AMP Sepharose 4B. The reaction catalyzed appears to be irreversible.

Although NADP can be used, NAD is the preferred coenzyme. The enzyme also exhibits an unusual property of being subject to substrate inhibition by NAD (Forte-McRobbie and Pietruszko, 1986).

II. Regulation of proline biosynthesis in development, ageing and in other physiological conditions

The physiologic relevance of P5C as an extracellular effector molecule besides being an intracellular intermediate in amino acid metabolism, was supported by the identification of P5C in normal human plasma (Fleming et al., 1989; Fleming et al., 1984). Strikingly, plasma P5C levels showed marked diurnal fluctuations, which correlated with feeding and which were markedly attenuated by fasting. Although the rise in plasma P5C seemed to be nutritionally dependent, it did not relate simply to ingestion of specific precursor amino acids, e.g., large oral bolus of proline did not increase plasma P5C. Presumably, the release of P5C into plasma is dependent on a number of metabolic and perhaps hormonal factors. Semon et al. reported that cultured human fibroblasts not only release P5C into the medium but also respond with increased release when stimulated by either insulin or insulin-like growth factor 1 (IGF-1) (Semon and Phang, 1991).

Isotopic evidence for the differential regulation of arginine and proline synthesis shown in man: In one study the subjects ingested a single dose of a mixture of uniformly (U)- ^{13}C -labeled amino acids and carbohydrate; the appearance of U- ^{13}C -amino acid in the plasma reflects entry of the dietary amino acids, and the appearance of ^{13}C in lower mass isotopomers demonstrates synthesis of the respective amino acids by the subject. The [M+5] isotopomer of plasma arginine was labeled as rapidly as [M+6] (i.e., tracer) arginine, which reflects the synthesis of citrulline from dietary [M+5]-glutamate in the first pass. The [M+1] to [M+3] isotopomers of arginine were also labeled for a prolonged period of time, suggesting that systemic glutamate was also a precursor for arginine synthesis. In fed subjects, only the [M+5] isotopomer of proline was significantly labeled. Fasting was associated with increased labeling of the [M+1] to [M+3] isotopomers of both arginine and proline, suggesting an increase in the contribution of de novo synthesis to their plasma flux (Berthold et al., 1995).

P5C concentration in the small intestine of fetuses was 1/5 that of adults and reached an adult level as early as postnatal Day 1. The highest peak was observed at Day 14, and then activity decreased to the adult level. However, P5C in the brain was highest at birth and quickly inactivated in a few days. Pregnancy and lactation reduced intestinal P5C considerably up to day 14 after parturition. P5C concentration in the small intestine of senescent rats was almost halved compared to young controls on a whole tissue basis. These results indicate that P5C synthetase in the small intestine is highly activated in suckling and weaning and amino acids derived from dietary ornithine are not fully dispensable in pregnancy, lactation, and senescence (Yamada and Wakabayashi, 1991).

III. Pathological states affecting the metabolism of P5C and glutamate family of amino acids

Stewart et al. (1981) showed that cats that develop hyperammonemia after an arginine-free meal have low hepatic ornithine levels. This group concluded that the sensitivity of the cat to an arginine free meal was the result of low levels of hepatic ornithine. With arginine deprivation, the net synthesis of ornithine becomes a critical source of urea cycle intermediates for the removal of excess ammonia. As a possible explanation for the unusual response of the cat to arginine deprivation Rogers and Phang (1985) have focused on the activity of P5C synthetase in intestinal mucosa of the cat. They found that P5C synthetase activity of the intestinal mucosa of cats, when compared to that of rats, was only 18% as high per gram of mucosa and only 5% as high per kilogram body weight. This severe limitation in the first step in the de novo synthesis of ornithine may be the metabolic basis for the severe hyperammonemia found in cats fed an arginine-deficient diet. This lack of ornithine synthesis makes the cat completely dependent on dietary arginine for the ornithine required for the removal of ammonia via urea synthesis in the liver.

Normally, in bovine ocular tissues, cornea and retinal outer layers provide the intracellular proline synthetic pathway from ornithine, but not from glutamate. In C3H retinal degeneration mice, P5C reductase activity in the retinal outer layers was decreased to about one third lesser than that of CRJ control mice in the specific activity, throughout suckling and weaning periods. The activity of an incorporation of ^{14}C -ornithine into proline in the retinal outer layers (including the choroid) which presumably participates in the biosynthesis of proline-rich proteins such as collagen and glycoproteins, was also decreased to the same extent at the weaning period, in the retinal degeneration mice (Matsuzawa et al., 1982a; Matsuzawa, 1982a).

In man, disorders of proline and hydroxyproline metabolism result in hyperprolinemia. Two distinct conditions are known, each due to a mutation in a different genetic locus; both are apparently autosomal recessive. Type I hyperprolinemia is due to a deficiency of proline oxidase, and type II hyperprolinemia is due to a deficiency of P5C dehydrogenase. Proline levels are elevated only three- to fivefold in the former condition but ten- to fifteenfold in the latter. The distinguishing biochemical findings in type II hyperprolinemia are high plasma P5C levels and urinary excretion of P5C. Although the metabolic disorders in the hyperprolinemias are consistent with normal adult life, recent findings strongly suggest that there is a causal relationship between type II hyperprolinemia and neurologic manifestations in childhood (Phang et al., 1995).

Rabier et al. (1992) described a clinical case where two children born to consanguineous parents presented with a joint hyperlaxicity, skin hyperelasticity, bilateral cataract and mental retardation with hyperammonemia and low citrulline, ornithine and proline in plasma. They suggested that these patients affected with Δ^1 -pyrroline-5-carboxylate synthetase (P5CS) deficiency. Molecular studies confirmed this hypothesis (Aral et al., 1997).

Gyrate atrophy of the choroid and retina is a progressive chorioretinal degeneration which is inherited as an autosomal recessive trait. It is caused by a deficiency of OAT. Plasma ornithine values range from 400 to 1400 μ M, and 0.5 to 10 mmol of ornithine is excreted daily. Plasma glutamate, glutamine, lysine, creatine and creatinine concentrations are modestly reduced. OAT activity in the cells and tissues of patients is from 0 to 6% that in control patients, and obligate heterozygotes have intermediate values (Phang et al., 1995).

IV. Gene-enzyme relationships in the proline biosynthetic pathway

In contrast to the poor progress in measuring enzymes of proline synthesis in higher plant extracts, there has been rapid progress in recent years in the cloning of cDNAs encoding these enzymes from higher plants by complementation of proline-requiring mutants of *E. coli*. A cDNA encoding P5CR was cloned by direct complementation of an *E. coli* proC proline auxotroph with a soybean nodule cDNA expression library (Delauney and Verma, 1990). This cDNA then facilitated the isolation of an homologous P5CR gene from *Pisum sativum* (Williamson and Slocum, 1992). The sequence of the cDNA encoding P5CR of soybean nodules also enabled the selection of suitable primers for the isolation of a P5CR gene from *Arabidopsis thaliana* (Verbruggen et al., 1993). In soybean, pea and *Arabidopsis*, P5CR transcripts increase in abundance in response to osmotic stress, indicating that P5CR gene transcription is under osmotic stress control (Delauney and Verma, 1990; Verbruggen et al., 1993; Williamson and Slocum, 1992). It is still not certain, however, whether this induction of P5CR mRNA contributes to enhanced proline accumulation under stress. The P5CR cDNA from soybean has been over-expressed in tobacco (Szoke et al., 1992). The transgenic plants exhibit 100-fold greater P5CR activity than wild-type. However, proline concentrations were not increased significantly in these transgenic plants (Szoke et al., 1992). Thus, P5CR may not be the rate-limiting step in proline accumulation (Delauney and Verma, 1993; LaRosa et al., 1991).

A cDNA encoding an ornithine- δ -aminotransferase was isolated from a mothbean (*Vigna aconitifolia*) cDNA expression library employing a proBA mutant of *E. coli* (CSH26) grown on nitrogen-free medium supplemented with 10 mM ornithine. This mutant cannot utilize ornithine as a proline source. Introduction of the ornithine- δ -aminotransferase cDNA from *Vigna* allowed conversion of ornithine to GSA $\rightleftharpoons$ P5C which was then readily converted to proline by P5CR encoded by the endogenous *E. coli* proC gene, facilitating growth on ornithine in the absence of exogenous proline (Delauney and Verma, 1993; Delauney et al., 1993; Verma et al., 1992). In striking contrast to P5CR, ornithine- δ -aminotransferase appears to be markedly down-regulated under salinity stress (Delauney et al., 1993).

A cDNA encoding both GK and GPR similarly was isolated from a mothbean cDNA expression library employing *E. coli* proA, proB and proBA

proline auxotrophs, and screening for cDNAs which permit growth in the absence of proline. This cDNA was found to encode a bifunctional enzyme Δ^1 -pyrroline-5-carboxylate synthetase (P5CS) (Hu et al., 1992). The single major open reading frame of this cDNA encodes a polypeptide of 73.2 kDa which has two distinct domains exhibiting 55.3% overall similarity to *E. coli* GK and 57.9% similarity to 168 *E. coli* GPR, respectively (Hu et al., 1992). The GK activity (as measured by the γ -glutamyl hydroxamate assay (Hayzer and Leisinger, 1980) of this bifunctional enzyme is inhibited by proline (50% inhibition with 6 mM proline) (Hu et al., 1992). This enzyme appears to be much less sensitive to feedback inhibition than the wild-type GK of *E. coli* (50% inhibition by 0.2 mM (Hu et al., 1992) to 0.1 mM proline (Smith, 1985). Northern analyses indicate that the P5CS gene is induced (particularly in leaves) by treatment of *Vigna* plants with 200 mM NaCl (Hu et al., 1992). A single substitution of an alanine for a phenylalanine at amino acid residue 129 of the P5CS resulted in a significant reduction of proline feedback inhibition. The 50% inhibition values of γ -GK activity of the wild type and the mutant P5CS were observed at 5 mM and 960 mM proline, respectively (Zhange et al., 1995). Garcia-Rios et al. (1991, 1994) have also recently cloned a cDNA encoding both GK and GPR, from a tomato fruit cDNA expression library by complementation of prob *E. coli* mutants. It was subsequently found that this cDNA (designated PRO1) complemented both proA and proBA mutants of *E. coli* (Garcia-Rios et al., 1991, 1994). The nucleotide sequence of the PRO1 cDNA revealed two protein domains; a GK domain consisting of 283 amino acids at the 5' end, and a GPR domain consisting of 429 amino acids at the 3' end (Garcia-Rios et al., 1994). The GK domain has a 66–69% amino acid sequence similarity of the GKs of *E. coli* and yeast, and a 69% similarity to the GK domain of mothbean P5CS. The GPR domain has a 69 and 66% similarity to the corresponding enzyme from *E. coli* and the corresponding domain of mothbean P5CS, respectively (Garcia-Rios et al., 1994). In marked contrast to mothbean P5CS, the tomato PRO1 cDNA has an inframe TAA (translation termination) codon between the two segments that specify GK and GPR (Garcia-Rios et al., 1994). This codon is followed after 5 bp (i.e. at a -1 or $+2$ frameshift with respect to GK) by a potential translation start ATG codon (Garcia-Rios et al., 1994). Consistent with this, when expressed in *E. coli* maxi cells, PRO1 directs the synthesis of two separate polypeptides (30 and 45 kDa) of the approximate mass predicted from the sequence analysis (Garcia-Rios et al., 1994).

The conversion of L-glutamate to proline in *E. coli* is achieved in a four-step reaction catalysed by GK, GPR and P5CR which are encoded by proB, proA, and proC, respectively. The *E. coli* proB, proA (Deutch et al., 1984) and proC (Deutch et al., 1984) loci have been cloned and sequenced. In *E. coli* and *Serratia marcescens*, the proB gene is located directly upstream of the proA gene and they appear to share a common promoter (Deutch et al., 1984). Similarly, the proB and proA genes of *Salmonella typhimurium* and of *C. jejuni* (Mahan and Csonka 1993; Lee et al., 1985) have been reported to form a single operon with proB located proximally to the promoter. The gene-enzyme relationships were demonstrated by interspecies complemen-




Fig. 3. Gene-enzyme relationship of the P5CS in different organisms (adapted from Hu et al., 1992)

tation of proB, proA, and proC mutants of *E. coli* and *S. typhimurium* by the yeast PRO1, PRO2, and PRO3 genes, respectively (Tomenchok and Brandriss, 1987) and of a yeast pro1 mutant by the *E. coli* proB gene (Orser et al., 1988). There is substantial deduced amino acid sequence similarity between the *E. coli* (Deutch et al., 1984) and *S. cerevisiae* (Li and Brandriss, 1992) GK, between the *E. coli* (Deutch et al., 1984) and *S. cerevisiae* (Brandriss, unpublished data) GPR, and among the P5C reductases of bacteria (Deutch et al., 1982; Hamilton and Reeve, 1985; Savioz et al., 1990), *S. cerevisiae* (Brandriss and Falvey, 1992), plants (Delauney and Verma, 1990) and mammals (Dougherty et al., 1992).

A human P5CS cDNA has been cloned by "database cloning" strategy and 2,907bp sequenced from this cDNA which has a closed open reading frame (ORF) of 2,385bp coding for a polypeptide of 795 amino acid residues (Aral et al., 1996). This cDNA, as its plant counterpart, encodes a bifunctional enzyme (Fig. 3), with both γ -glutamyl kinase GK and γ -glutamyl phosphate reductase GPR activities that catalyzes the first two steps in proline biosynthesis and it hybridizes to a 4.5kb mRNA from various tissues, including spleen, thymus, prostate, testis, ovary, small intestine, heart and skeletal muscle, pancreas, among others. In humans, the expression levels found in heart, skeletal muscle and pancreas were relatively high compared to other tissues (Aral et al., 1996). In rats, assay for P5CS activities showed values 10–1,000 times higher in small intestine in comparison to other tissues (Wakabayashi et al., 1991). However, our results showed that in humans the mRNA levels detected in other tissues equals, or even higher, those in small intestine, contrasting with the tissue distribution of P5CS activity in rats given by Wakabayashi (Wakabayashi et al., 1991). Northern blot analyses showed that the P5CS gene is induced (particularly in chondrocytes) by treatment of rabbit cartilage cells in culture with IGF-1 (Aral, unpublished data; Smith et al., 1979), indicating that P5CS gene transcription is under hormonal control. The human gene encoding P5CS has been assigned to 10q24.3 by in situ hybridization (Liu et al., 1996). Dougherty et al. (1992) cloned a human P5C reductase

cDNA by complementation of proline auxotrophy in a *S. cerevisiae* mutant strain (DT1100). The 1,810-bp P5C reductase cDNA hybridizes to a 1.85-kb mRNA in samples from human cell lines and predicts a 319-amino acid, 33.4-kDa protein. The derived amino acid sequence is 32% identical with that of *S. cerevisiae*. By genomic DNA hybridization analysis, the human P5C reductase appears to be encoded by a single copy gene which maps to chromosome 17. The tissue expression profile of P5C reductase mRNA is ubiquitary. The P5C dehydrogenase, the gene defective in type II hyperprolinemia has also been cloned by using published peptide sequence data and degenerate primer PCR. Hu et al. (1996) cloned two full length human P5CDH cDNAs, differing in length by 1 kb. Both have a 1,689bp ORF encoding a protein of 563 residues with a predicted molecular mass of 60 kDa. The predicted amino acid sequence has 100% identity to human P5CDH peptide sequence and 43% identity to *S. cerevisiae* P5CDH. The human P5CDH mRNA transcript has 3.2kb length. The human OAT cDNA and structural gene have been cloned and characterized and the OAT locus mapped to 10q26 (Mitchell et al., 1988; Inana et al., 1986; Ramesh et al., 1986, 1987; Barrett et al., 1987).

V. The physiological roles of P5C and proline

A specific system of enzymes with special properties has evolved to mediate the metabolism of proline. These metabolic interconversions among proline, ornithine, and glutamate have P5C as the sole common intermediate. There are three poles of production, namely the intestinal mucosa which produces ornithine and citrulline, the liver which produces urea and the kidney which produces arginine. The intestinal mucosa, produces and exports both proline and ornithine. Only in the light of human genetic pathology and animal models related to the deficiencies of enzymes surrounding P5C-proline metabolism, we can truly speak about tissue specificity for each metabolic function.

The P5CS deficiency has shed some light on the importance of the ability to synthesize P5C and convert it to ornithine in enterocytes. The mutation these patients show is in the structural gene for P5C synthetase (Aral et al., 1997). As a result, these patients have a paradoxical hyperammonemia associated to hypocitrullinemia, hypoornithinemia and hypoprolinemia. This situation, coupled with the impaired production of urea, indicates that the conversion of P5C to proline and to ornithine is important for the maintenance of normal blood (and tissue) glutamate family amino acids and for the normal turnover of urea cycle and blood ammonemia levels. Whereas hyperammonemia is worsened after a protein load in most inborn errors of metabolism (such as urea cycle disorders and organic acidemias), in these patients hyperammonemia was more severe preprandially. It has been suggested that the cause of the paradoxical nature of hyperammonemia is that food intake provides the necessary ornithine to urea cycle whereas in fasting the ornithine is provided from the conversion of the P5C to ornithine in enterocytes via OAT reaction. Besides, in OAT $-/-$ mice, during the neonatal periods, there is a marked reduction in plasma ornithine,

arginine and citrulline (Wang et al., 1995). These animals also exhibit hyperammonemia and profound orotic aciduria which suggests impaired ammonium detoxification due to deficiency of urea cycle intermediates and the availability of intramitochondrial ornithine is inadequate for citrulline synthesis (Hauser et al., 1990). These data indicate that the flux in the OAT reaction in the neonatal period is in the direction of ornithine synthesis rather than degradation and that the survival of the rapidly growing neonatal mouse depends on an intact P5CS and OAT reaction to synthesize ornithine and arginine. The other animal model which support the usefulness of glutamate $\Rightarrow$ P5C $\Rightarrow$ ornithine pathway is the deficiency of P5CS in the intestinal mucosa of the cat, which makes the cat completely dependent on dietary arginine for the ornithine required for the removal of ammonia via urea synthesis in the liver (Rogers and Phang, 1985).

On the other hand the lack of P5CS in the liver, and the very high level of P5CDH which forms glutamate from P5C, indicates that glutamate synthesis may be important in liver. The high level of OAT in liver may indicate that OAT play a central role in regulating ornithine levels available to urea cycle; There is a human genetic disease, gyrate atrophy (GA), where patients suffering from it present themselves with limitations in peripheral vision due to atrophy of the retina, which starts at the periphery and then gyrates toward the center of the retina. The mutation these patients show is in the structural gene for OAT. As a result, these patients have high levels of blood ornithine. This situation, coupled with the normal production of urea, indicates that the conversion of ornithine to P5C (ornithine $\Rightarrow$ P5C) is important for the maintenance of normal blood (and tissue) ornithine levels. It has been suggested that the cause of the retinal atrophy is that cellular conversion of ornithine to P5C and conversion of P5C to proline (ornithine $\Rightarrow$ P5C $\Rightarrow$ proline pathway) is essential. Knockout adult mice (OAT $-/-$) for OAT on a standard diet exhibit profound hyperornithinemia to an extent similar to that of humans with GA (Wang et al., 1995).

In some initial studies, it was shown that P5C markedly activated the metabolism of glucose through the pentose phosphate pathway (Phang et al., 1979, 1982). The effect was a function of P5C concentrations and was observed at all concentrations of glucose (Phang, 1985). The unaltered response to P5C in cells deficient in either OAT or P5CDH, suggested that P5C reductase is the enzyme mechanism mediating the effect. In human erythrocytes, P5C reductase is the only enzyme of the proline metabolic system (Yeh and Phang, 1981), and in incubations of human red cells P5C stimulated the metabolism of glucose through the pentose phosphate pathway and there was a good correlation between two molecules of P5C converted to proline for every glucose 6-phosphate metabolized to ribose 5-phosphate. These initial studies support the general hypothesis that the interconversions of proline, ornithine, and glutamate can stimulate a metabolic pathway dependent on the NADP⁺/NADPH redox ratio. There was also an effect of P5C on PP-ribose and purine nucleotides. Indeed, as PP-ribose-P levels in incubated human erythrocytes markedly increased when cells were treated with P5C (Phang, 1985; Yeh and Phang, 1981; Yeh and Phang, 1983). Although circulating

human erythrocytes lack the pathways for de novo purine synthesis they readily incorporate preformed purines into nucleotides by the salvage pathway in a PP-ribose-P-dependent process (Pritchard et al., 1975). P5C added to red cells markedly increased the incorporations of labeled purines into their respective nucleotides (Phang, 1985; Yeh and Phang, 1981), P5C also stimulated the IMP synthesis from hypoxanthine. A possible mechanism for the P5C effect on nucleotides via PP-ribose-P synthetase has been evoked (Phang, 1985).

Proline belongs to a class of compounds known as “compatible” or “counteracting solutes” which includes several amino acids (e.g. proline, alanine, β -alanine and taurine). These compounds tend to be uncharged at neutral pH, and are of high solubility in water (Ballantyne and Chamberline, 1994). In fact, exogenously supplied proline is osmoprotective for bacteria, facilitating growth in highly saline environments (Csonka, 1989; Csonka and Hanson, 1991; Strom et al., 1983; Yancey, 1994). Overproduction of proline in bacteria caused by altered feedback inhibition of the proline biosynthesis pathway can result in increased osmotolerance (Csonka, 1981; Smith, 1985). In higher plants, the accumulation of proline is a common metabolic response of higher plants to water deficits, salinity stress and cold stress (Csonka and Baich, 1983; Delauney and Verma, 1993; Dorffling et al., 1993; Hanson and Hitz, 1982; Steward, 1981; Steward and Larher, 1980; Thompson, 1980). *Fasciola hepatica* manipulates the metabolism of the host animal using a mechanism involving proline. While still in the host liver, the immature fluke has high proline concentrations and has a large proline biosynthetic capacity (Isseroff, 1979). The fluke releases proline into the host's bile increasing the concentration of biliary proline more than 1,000-fold, and inducing ipso facto, proliferation and enlargement of the host's bile duct so that it becomes a suitable habitat.

Conclusions

Plants, microorganisms and animal species have all three necessary enzymes which produce (the P5C-producing enzymes) and utilize (the P5C-utilizing enzymes) the P5C. But they use the P5C for different purposes. Certainly, the glutamate $\Rightarrow$ P5C $\Rightarrow$ proline pathway is the most studied in three organisms. Proline biosynthesis as a substrate to be incorporated into proteins must exist in plants, microorganisms and animal species. The tissues with high collagen content use this pathway in order to supply proline during tissue development. In microorganisms and plants the biosynthesis of proline is a necessary biological response to water deficits, salinity stress and cold stress. It serves as “compatible” or “counteracting solutes”. In animal species and in man, the pathway glutamate $\Rightarrow$ P5C $\Rightarrow$ ornithine is also used in enterocytes in order to produce ornithine, in the absence of exogenous ornithine, for a correct urea cycle homeostasis (and also citrulline and arginine biosynthesis). The cloning of cDNAs encoding key enzymes of proline biosynthesis, such as P5CS from animal species and man, may prove to be as useful as the specific probes for these genes and their products will clarify our understanding of

urea cycle homeostasis and collagen metabolism. Hepatocytes and retina cells use the ornithine $\Rightarrow$ P5C $\Rightarrow$ proline pathway in order to adjust the ornithine levels available to urea cycle and to supply proline to neuronal cells.

References

- Adams E (1970) Metabolism of proline and of hydroxyproline. *Int Rev Connect Tissue Res* 5: 1–91
- Adams E, Frank L (1980) Metabolism of proline and the hydroxyprolines. *Annu Rev Biochem* 49: 1005–1061
- Adams E, Goldstone A (1960) Hydroxyproline metabolism. IV. Enzymatic synthesis of γ -hydroxyglutamate from Δ^1 -pyrroline-3-hydroxy-5-carboxylate. *J Biol Chem* 235: 3504–3512
- Aral B, Schlenzig JS, Liu G, Kamoun P (1996) Database cloning human Δ^1 -pyrroline-5-carboxylate synthetase (P5CS) cDNA. *C R Acad Sci* 319: 171–178
- Aral B, Rabier D, Poggi F, Saudubray JM, Kamoun P (1997) A mutation of Δ^1 -pyrroline 5-carboxylate synthetase associated with hyperammonemia, hypocitrullinemia, hypornithinemia and hypoprolinemia (submitted to publication)
- Baich A (1969) Proline synthesis in *Escherichia coli*. A proline-inhibitable glutamic acid kinase. *Biochim Biophys Acta* 192: 462–467
- Baich A (1971) The biosynthesis of proline in *Escherichia coli*. Phosphate-dependent glutamate γ -semialdehyde dehydrogenase (NADP), the second enzyme in the pathway. *Biochim Biophys Acta* 244: 129–134
- Baich A (1977) Alternative pathways for proline synthesis in mammalian cells. *Somatic Cell Genet* 3: 529–538
- Ballantyne JS, Chamberlin ME (1994) Regulation of cellular amino acid levels. In: Strange K (ed) *Cellular and molecular physiology of cell volume regulation*. CRC Press, Boca Raton, FL, pp 111–122
- Barrett DJ, Bateman JB, Sparkes RS, Mohandas T, Klisak I, Inana G (1987) Chromosomal localization of human ornithine aminotransferase gene sequences to 10q26. *Invest Ophthalmol Vis Sci* 28: 1037–1042
- Berthold HK, Reeds PJ, Klein PD (1995) Isotopic evidence for the differential regulation of arginine and proline synthesis in man. *Metabolism* 44: 466–473
- Boggess SE, Stewart CR (1980) The relationship between water stress induced proline accumulation and inhibition of protein synthesis in tobacco leaves. *Plant Sci Lett* 17: 245–252
- Boggess SE, Aspinall D, Paleg LG (1976) Stress metabolism. IX. The significance of end-product inhibition of proline biosynthesis and of compartmentation in relation to stress-induced proline accumulation. *Aust J Plant Physiol* 3: 513–525
- Brandriss MC (1979) Isolation and preliminary characterization of *Saccharomyces cerevisiae* proline auxotrophs. *J Bacteriol* 138: 816–822
- Brandriss MC, Magasanik B (1980) Proline: an essential intermediate in arginine degradation in *Saccharomyces cerevisiae*. *J Bacteriol* 143: 1403–1410
- Brandriss MC, Magasanik B (1981) Subcellular compartmentation in control of converging pathways for proline and arginine. *J Bacteriol* 145: 1359–1364
- Brandriss MC, Falvey DA (1992) Proline biosynthesis in *Saccharomyces cerevisiae*: analysis of the PRO3 gene, which encodes Δ^1 pyrroline-5-carboxylate reductase. *J Bacteriol* 174: 3782–3788
- Conn EE, Stumpf PK, Bruening G, Doi R (1987) The metabolism of ammonia and nitrogen-containing compounds. In: *Outlines of biochemistry*, 5th edn. John Wiley & Sons, New York, p 530
- Csonka LN (1981) Proline over-production results in enhanced osmotolerance in *Salmonella typhimurium*. *Mol Gen Genet* 182: 82–86

- Csonka LN (1989) Physiological and genetic responses of bacteria to osmotic stress. *Microbiol Rev* 53: 121–147
- Csonka LN, Baich A (1983) Proline biosynthesis. In: Herrmann KM, Somerville RL (eds) *Amino acids, biosynthesis and genetic regulation*. Addison-Wesley, Reading, pp 35–51
- Csonka LN, Hanson AD (1991) Prokaryotic osmoregulation: genetics and physiology. *Annu Rev Microbiol* 45: 569–606
- Delauney AJ, Verma DPS (1990) A soybean Δ^1 -pyrroline-5-carboxylate reductase gene was isolated by functional complementation in *Escherichia coli* and is found to be osmoregulated. *Mol Gen Genet* 221: 299–305
- Delauney AJ, Verma DPS (1993) Proline biosynthesis and osmoregulation in plants. *Plant J* 4: 215–223
- Delauney AJ, Hu C-AA, Kisher PBK, Verma DPS (1993) Cloning of ornithine δ -aminotransferase cDNA from *Vigna aconitifolia* by trans-complementation in *Escherichia coli* and regulation of proline synthesis. *J Biol Chem* 268: 18673–18678
- Deutch AH, Rushlow KE, Smith CJ, Kretschmer PJ (1982) *Escherichia coli* (1-pyrroline-5-carboxylate reductase: gene sequence, protein overproduction and purification. *Nucleic Acids Res* 10: 7701–7714
- Deutch AH, Smith CJ, Rushlow KE (1984) Analysis of the *Escherichia coli* proBA locus by DNA and protein sequencing. *Nucleic Acids Res* 12: 6337–6355
- Dorffling K, Dorffling H, Lesselich G (1993) In vitro-selection and regeneration of hydroxyproline-resistant lines of winter wheat with increased proline content and increased frost tolerance. *J Plant Physiol* 142: 222–225
- Dougherty KM, Brandriss MC, Valle D (1992) Cloning human pyrroline-5-carboxylate reductase cDNA by complementation in *Saccharomyces cerevisiae*. *J Biol Chem* 267: 871–875
- Elthon TE, Stewart CR (1982) Proline oxidation in corn mitochondria. *Plant Physiol* 70: 567–572
- Fleming GA, Hagedorn CH, Granger AS, Phang JM (1984) Pyrroline-5-carboxylate in human plasma. *Metabolism* 33: 739–742
- Fleming GA, Granger A, Rogers QR, Prosser M, Ford DB, Phang JM (1989) Fluctuations in plasma pyrroline-5-carboxylate concentrations during feeding and fasting. *J Clin Endocrinol Metab* 69: 448–452
- Forte McRobbie CM, Pietruszko R (1986) Purification and characterization of human liver “high Km” aldehyde dehydrogenase and its identification as glutamic γ -semialdehyde dehydrogenase. *J Biol Chem* 261: 2154–2163
- Fukutoku Y, Yamada Y (1984) Sources of proline-nitrogen in water-stressed soybean (*Glycine max*). II. Fate of ^{15}N -labelled protein. *Physiol Plant* 61: 622–628
- Gamper H, Moses V (1974) Enzyme organization in the proline biosynthetic pathway of *Escherichia coli*. *Biochim Biophys Acta* 354: 75–87
- Garcia-Rios MG, LaRosa PC, Bressan RA, Csonka LN, Hanquier JM (1991) Cloning by complementation of the γ -glutamyl kinase gene from a tomato expression library. In: Hallick RB (ed) *Abstracts of the Third International Congress of Plant Molecular Biology*, Tucson, Arizona, Oct. 6–11, 1991. Abstract #1507.
- Garcia-Rios MG, Csonka LN, Bressan RA, LaRosa PC, Haliquier J (1994) Cloning of a DNA fragment encoding γ -glutamyl kinase and γ -glutamyl phosphate reductase from a tomato cDNA library. In: Cherry JH (ed) *Cell biology: biochemical and cellular mechanisms of stress tolerance in plants*, NATO ASI Series H. Springer, Berlin Heidelberg New York Tokyo
- Hamilton PT, Reeve JN (1985) Structure of genes and an insertion element in the methane producing archaeobacterium. *Mol Gen Genet* 200: 47–59
- Hanson AD, Hitz WD (1982) Metabolic responses of mesophytes to plant water deficits. *Annu Rev Plant Physiol* 33: 163–203
- Hauser ER, Finkelstein JE, Valle D, Brusilow SW (1990) Allopurinol-induced

- orotidinuria. A test for mutations at the ornithine carbamoyltransferase. *N Engl J Med* 322: 1641–1645
- Hayzer DJ, Leisinger T (1980) The gene-enzyme relationships of proline biosynthesis in *Escherichia coli*. *J Gen Microbiol* 118: 287–293
- Hayzer DJ, Leisinger T (1981) Proline biosynthesis in *Escherichia coli*. Stoichiometry and end-product identification of the reaction catalysed by glutamate semialdehyde dehydrogenase. *Biochem J* 197: 269–274
- Hayzer DJ, Leisinger T (1982) Proline biosynthesis in *Escherichia coli*. Purification and characterization of glutamate-semialdehyde dehydrogenase. *Eur J Biochem* 121: 561–565
- Hayzer DJ, Moses V (1978a) Proline biosynthesis by cell-free extracts of *Escherichia coli* and potential errors arising from the use of a bioradiological assay procedure. *Biochem J* 173: 207–217
- Hayzer DJ, Moses V (1978b) The enzymes of proline biosynthesis in *Escherichia coli*. Their molecular weights and the problem of enzyme aggregation. *Biochem J* 173: 219–228
- Hensless JG, Jones ME (1982) Ornithine synthesis from glutamate in rat small intestinal mucosa. *Arch Biochem Biophys* 219: 186–197
- Herzfeld A, Mezl VA, Knox WE (1977) Enzymes metabolizing Δ^1 -pyrroline-5-carboxylate in rat tissues. *Biochem J* 166: 95–103
- Heyser JW, De Bruin D, Kincard ML, Johnson RY, Rodriguez MM, Robinson NJ (1989a) Inhibition of NaCl-induced proline biosynthesis by exogenous proline in halophytic *Distichlis spicata* suspension cultures. *J Exp Bot* 40: 225–232
- Heyser JW, Chacon MJ, Warren RC (1989b) Characterization of L-[5- ^{13}C]-proline biosynthesis in halophytic and nonhalophytic suspension cultures by ^{13}C NMR. *J Plant Physiol* 135: 459–466
- Hu C-AA, Delauney AJ, Verma DPS (1992) A bifunctional enzyme (Δ^1 -pyrroline-5-carboxylate synthetase) catalyzes the first two steps in proline biosynthesis in plants. *Proc Natl Acad Sci USA* 89: 9354–9358
- Hu CA, Geraghty M, Lin W-W, Valle D (1996) Type II hyperprolinemia: cloning and expression in yeast of cDNAs encoding human Δ^1 -pyrroline-5-carboxylate (P5C) dehydrogenase and delineation of mutations responsible for type II hyperprolinemia (HP II). In: Byers PH (ed) Abstracts of the 46th Annual Meeting of the American Society of Human Genetics, San Francisco, California, Oct. 29–Nov. 2, 1996. Abstract #1146
- Huang AHC, Cavalieri AJ (1979) Proline oxidase and water-stress induced proline accumulation in spinach leaves. *Plant Physiol* 63: 531–535
- Inana G, Totsuka S, Redmond M, Dougherty T, Nagle J, Shiono T, Ohura T (1986) Molecular cloning of human ornithine aminotransferase mRNA. *Proc Natl Acad Sci USA* 83: 1203–1207
- Isseroff H (1979) In: Simeon V, Vitale LJ (eds) Proceedings of the FEBS Special Meeting on Enzymes. Pergamon, Oxford, pp 303–314
- Jernigan HM Jr (1990) Metabolism of glutamine and glutamate in human lenses. *Exp Eye Res* 50: 597–601
- Johnson AB, Strecker HJ (1962) The interconversion of glutamic acid and proline. IV. The oxidation of proline by rat liver mitochondria. *J Biol Chem* 237: 1876–1882
- Ketchum REB, Warren RC, Klima LJ, Lopez-Gutierrez F, Nabors MW (1991) The mechanism and regulation of proline accumulation in suspension cultures of the halophytic grass *Distichlis spicata* L. *J Plant Physiol* 137: 368–374
- Kowaloff EM, Granger AS, Phang JM (1976) Alterations in proline metabolic enzymes with mammalian development. *Metabolism* 25: 1087–1094
- Kowaloff EM, Phang JM, Granger AS, Downing SJ (1977) Regulation of proline oxidase activity by lactate. *Proc Natl Acad Sci USA* 74: 5368–5371
- Kowaloff EM, Phang JM, Granger AS, Downing SJ (1978) Glucocorticoid induction of proline oxidase in LLC-RK1 cells. *J Cell Physiol* 97: 153–159

- Kramer JJ, Henslee JG, Wakabayashi Y, Jones ME (1985) Δ^1 -pyrroline-5-carboxylate synthase from rat intestinal mucosa. *Methods Enzymol* 113: 113–120
- Krishna RV, Leisinger T (1979) Biosynthesis of proline in *Pseudomonas aeruginosa*. Partial purification and characterization of γ -glutamyl kinase. *Biochem J* 181: 215–222
- Krishna RV, Beilstein P, Leisinger T (1979) Biosynthesis of proline in *Pseudomonas aeruginosa*. Properties of γ -glutamyl phosphate reductase and Δ^1 -pyrroline-5-carboxylate reductase. *Biochem J* 181: 223–230
- Krueger R, Jager H-J, Hintz M, Pahlich E (1986) Purification to homogeneity of pyrroline-5-carboxylate reductase of barley. *Plant Physiol* 80: 142–144
- LaRosa RC, Rhodes D, Rhodes JC, Bressati RA, Csonka LN (1991) Elevated accumulation of proline in NaCl-adapted tobacco cells is not due to altered Δ^1 -pyrroline-5-carboxylate reductase. *Plant Physiol* 96: 245–250
- Lee EC, Walker RI, Guerry P (1985) Expression of *Campylobacter* genes for proline biosynthesis in *Escherichia coli*. *Can J Microbiol* 31: 1064–1067
- Li W, Brandriss MC (1992) Proline biosynthesis in *Saccharomyces cerevisiae*: molecular analysis of the PRO1 gene, which encodes γ -glutamyl kinase. *J Bacteriol* 174: 4148–4156
- Liu G, Maunoury C, Kamoun P, Aral B (1996) Assignment of the human gene encoding the Δ^1 -pyrroline-5-carboxylate synthetase (P5CS) to 10q24.3 by in situ hybridization. *Genomics* 37: 145–146
- Lodato RF, Smith RJ, Valle D, Phang JM, Aoki TT (1981) Regulation of proline biosynthesis: the inhibition of pyrroline-5-carboxylate synthase activity by ornithine. *Metabolism* 30: 908–913
- Lorans G, Phang JM (1981) Proline synthesis and redox regulation: differential functions of pyrroline-5-carboxylate. *Biochem Biophys Res Commun* 101: 1018–1025
- Lorans G, Rosenfeld C, Petitou M, Phan-Dinh-Tuy F, Mathe G (1978) Metabolism of proline in a human leukemic lymphoblastoid cell line. *Cancer Res* 38: 3950–3953
- Lundgren D, Ogur M (1972) Amino acid inhibition of a Δ^1 -pyrroline-5-carboxylate dehydrogenase preparation from beef kidney mitochondria. *Biochem Biophys Res Commun* 49: 147–149
- Lundgren DW, Ogur M (1973) Inhibition of yeast Δ^1 -pyrroline-5-carboxylate dehydrogenase by common amino acids and the regulation of proline catabolism. *Biochim Biophys Acta* 297: 246–257
- Lyons RT, Pitot HC (1976) The regulation of ornithine aminotransferase synthesis by glucagon in the rat. *Arch Biochem Biophys* 174: 262–272
- Mahan MJ, Csonka LN (1983) Genetic analysis of the proBA genes of *Salmonella typhimurium*: physical and genetic analyses of the cloned proB⁺A⁺ genes of *Escherichia coli* and of a mutant allele that confers proline overproduction and enhanced osmotolerance. *J Bacteriol* 156: 1249–1262
- Matsuzawa T (1982a) Disease of the ornithine-proline pathway: Δ^1 -pyrroline-5-carboxylate reductase. *Metab Pediatr Syst Ophthalmol* 6: 123–128
- Matsuzawa T (1982b) Purification and characterization of pyrroline-5-carboxylate reductase from bovine retina. *Biochim Biophys Acta* 717: 215–219
- Matsuzawa T, Ishiguro I (1980) Δ^1 -Pyrroline-5-carboxylate reductase from Baker's yeast. Purification, properties and its application in the assays of L- δ^1 -pyrroline-5-carboxylate and L-ornithine in tissue. *Biochim Biophys Acta* 613: 318–323
- Matsuzawa T, Obara Y (1987) Amino acid synthesis from ornithine: enzymes and quantitative comparison in brain slices and detached retinas from rats and chicks. *Brain Res* 413: 314–319
- Matsuzawa T, Iwasaki K, Hiraiwa N, Inagaki E, Ishiguro I (1982a) Disease of ornithine-proline pathway: a Δ^1 -pyrroline-5-carboxylate reductase deficiency in the retina of retinal degeneration mice. *Adv Exp Med Biol* 153: 361–370
- Matsuzawa T, Iwasaki K, Hiraiwa N, Inagaki E, Ishiguro I (1982b) In: Lowenthal A, Mori A, Marescau B (eds) *Urea cycle disease*. Plenum Press, New York

- Mazelis M (1979) ATP-dependent hydrolysis of 5-oxoproline to glutamate in higher plants. In: Hewitt EJ, Cutting CV (eds) *Nitrogen assimilation of plants*. Academic Press, London, pp 407–417
- McGivan JD, Bradford NM, Beavis AD (1977) Factors influencing the activity of ornithine aminotransferase in isolated rat liver. *Biochem J* 162: 147–156
- Meister A, Radhakrishnan AN, Buckley SD (1957) Enzymatic synthesis of L-pipecolic acid and L-proline. *J Biol Chem* 229: 789–800
- Mestichelli LJJ, Gupta RN, Spenser ID (1979) The biosynthetic route from ornithine to proline. *J Biol Chem* 254: 640–647
- Meyer J (1977) Proline transport in rat liver mitochondria. *Arch Biochem Biophys* 178: 387–395
- Mezl VA, Knox WE (1977) Metabolism of arginine in lactating rat mammary gland. *Biochem J* 166: 105–113
- Mixson AJ, Granger AN, Phang JM (1991) An assay for pyrroline-5-carboxylate based on its interaction with cysteine. *Anal Letters* 24: 625–641
- Mifflin BJ, Lea PJ (1977) Amino acid metabolism. *Annu Rev Plant Physiol* 28: 299–329
- Mitchell GA, Looney JE, Brody LC, Steel G, Suchanek M, Engelhardt JF (1988) Human ornithine- δ -aminotransferase. cDNA cloning and analysis of the structural gene. *J Biol Chem* 263: 14288–14295
- Morris CJ, Thompson JF, Johnson CM (1969) Metabolism of glutamic acid and N-acetylglutamic acid in leaf discs and cell-free extracts of higher plants. *Plant Physiol* 44:1023–1026
- Mueckler MM, Merrill MJ, Pitot HC (1983) Translational and pretranslational control of ornithine aminotransferase synthesis in rat liver. *J Biol Chem* 258: 6109–6114
- Oaks A (1992) The function of roots in the synthesis of amino acids and amides. In: Singh BK, Flores HE, Shannon JC (eds) *Biosynthesis and molecular regulation of amino acids in plants*. American Society of Plant Physiologists, Rockville, MD, pp 111–120
- Oaks A, Mitchell DJ, Barnard RA, Johnson FJ (1970) The regulation of proline biosynthesis in maize roots. *Can J Bot* 48: 2249–2258
- Orser CS, Goodner BW, Johnston M, Gelvin SB, Csonka LN (1988) The *Escherichia coli* proB gene corrects the proline auxotrophy of *Saccharomyces cerevisiae* PRO1 mutants. *Mol Gen Genet* 212: 124–128
- Phang JM (1985) The regulatory functions of proline and pyrroline-5-carboxylic acid. *Curr Top Cell Regul* 25: 91–132
- Phang JM, Downing SJ, Valle D (1973) A radioisotopic assay for Δ^1 -pyrroline-5-carboxylate reductase. *Anal Biochem* 55: 266–271
- Phang JM, Downing SJ, Yeh GC, Smith RJ, Williams JA (1979) Stimulation of the hexose-monophosphate pentose pathway by Δ^1 -pyrroline-5-carboxylic acid in human fibroblasts. *Biochem Biophys Res Commun* 87: 363–370
- Phang JM, Downing SJ, Yeh GC, Smith RJ, Williams JA, Hagedorn CH (1982) Stimulation of the hexosemonophosphate-pentose pathway by pyrroline-5-carboxylate in cultured cells. *J Cell Physiol* 110: 255–261
- Phang JM, Yeh GC, Scriver CR (1995) Disorders of proline and hydroxyproline metabolism. In: Scriver C, Beaudet A, Sly W, Valle D (eds) *The metabolic and molecular bases of inherited disease*. McGraw Hill, New York, pp 1125–1146
- Pritchard JB, O'Connor N, Oliver JM, Berlin RD (1975) Uptake and supply of purine compounds by the liver. *Am J Physiol* 229: 967–972
- Rabier D, Nuttin C, Poggi F, Padovani JP, Abdo K, Bardet J, Parvy P, Kamoun P, Saudubray JM (1992) Familial joint hyperlaxicity, skin hyperelasticity, cataract and mental retardation with hyperammonemia and low citrulline, ornithine and proline. A new disorder of collagen metabolism? In: SSIEM (ed) *Abstracts of the 30th Annual Symposium of the The Society for the Study of Inborn Errors of Metabolism (SSIEM)*, Leuven, September 8th–11th, 1992. Abstract #P61
- Ramesh V, Shaffer MM, Allaire JM, Shih VE, Gusella JF (1986) Investigation of gyrate atrophy using a cDNA clone for human ornithine aminotransferase. *DNA* 5: 493–501

- Ramesh V, Eddy R, Bruns GA, Shih VE, Shows TB, Gusella JF (1987) Localization of the ornithine aminotransferase gene and related sequences on two human chromosomes. *Hum Genet* 76: 121–126
- Rayapati PJ, Stewart CR (1991) Solubilization of a proline dehydrogenase from maize (*Zea mays* L.) mitochondria. *Plant Physiol* 95: 787–791
- Riby JE, Hurwitz RE, Kretchmer N (1990) Development of ornithine metabolism in the mouse intestine. *Pediatr Res* 28: 261–265
- Rhodes D, Handa S (1989) Amino acid metabolism in relation to osmotic adjustment in plant cells. In: Cherry JH (ed) *Environmental stress in plants: biochemical and physiological mechanisms*, NATO ASI Series, vol G19. Springer, Berlin Heidelberg New York Tokyo, pp 41–62
- Rhodes D, Handa S, Dressait RA (1986) Metabolic changes associated with adaptation of plant cells to water stress. *Plant Physiol* 82: 890–903
- Rogers QR, Phang JM (1985) Deficiency of pyrroline-5-carboxylate synthase in the intestinal mucosa of the cat. *J Nutr* 115: 146–150
- Ross G, Dunn D, Jones ME (1978) Ornithine synthesis from glutamate in rat intestinal mucosa homogenates: evidence for the reduction of glutamate to gamma-glutamyl semialdehyde. *Biochem Biophys Res Commun* 85: 140–147
- Rossi JJ, Vender J, Berg CM, Coleman WH (1977) Partial purification and some properties of Δ^1 -pyrroline-5-carboxylate reductase from *Escherichia coli*. *J Bacteriol* 129: 108–114
- Rushlow KE, Deutch AH, Smith CJ (1984) Identification of a mutation that relieves γ -glutamyl kinase from allosteric feedback inhibition by proline. *Gene* 39: 109–112
- Savioz A, Jeenes DJ, Kocher HP, Haas D (1990) Comparison of proC and other house-keeping genes of *Pseudomonas aeruginosa* with their counterparts in *Escherichia coli*. *Gene* 86: 107–111
- Seddon AP, Zhao KY, Meister A (1989) Activation of glutamate by γ -glutamate kinase: formation of γ -cis-cycloglutamyl phosphate, an analog of γ -glutamyl phosphate. *J Biol Chem* 264: 11326–11335
- Semon BA, Phang JM (1991) Accumulation of pyrroline 5-carboxylic acid in conditioned medium of cultured fibroblast: stimulatory effects of serum, insulin, and IGF-1. *In Vitro Cell Dev Biol* 27A: 665–669
- Shiono T, Kador PF, Kinoshita JJ (1986) Purification and characterization of rat lens pyrroline-5-carboxylate reductase. *Biochim Biophys Acta* 881: 72–78
- Small C, Jones ME (1987) A specific radiochemical assay for pyrroline-5-carboxylate dehydrogenase. *Anal Biochem* 161: 380–386
- Small WC, Jones ME (1990) Pyrroline 5-carboxylate dehydrogenase of the mitochondrial matrix of rat liver. *J Biol Chem* 265: 18668–18672
- Smith CJ, Deutch AH, Rushlow KE (1984) Purification and characteristics of a γ -glutamyl kinase involved in *Escherichia coli* proline biosynthesis. *J Bacteriol* 157: 545–551
- Smith LT (1985) Characterization of a γ -glutamyl kinase from *Escherichia coli* that confers proline overproduction and osmotic tolerance. *J Bacteriol* 164: 1088–1093
- Smith ME, Greenberg DM (1957) Preparation and properties of partially purified glutamic semialdehyde reductase. *J Biol Chem* 226: 317–327
- Smith RJ, Phang JM (1978) Proline metabolism in cartilage: the importance of proline biosynthesis. *Metabolism* 27: 685–694
- Smith RJ, Phang JM (1979) The importance of ornithine as a precursor for proline in mammalian cells. *J Cell Physiol* 98: 475–481
- Smith RJ, Downing SJ, Phang JM (1977) Enzymatic synthesis and purification of L-pyrroline-5-carboxylic acid. *Anal Biochem* 82: 170–176
- Smith RJ, Reddi AH, Phang JM (1979) Changes in proline synthetic and degradative enzymes during matrix-induced cartilage and bone formation. *Calcif Tissue Int* 27: 275–279

- Smith RJ, Downing SJ, Phang JM, Lodato RF, Aoki TT (1980) Pyrroline-5-carboxylate synthase activity in mammalian cells. *Proc Natl Acad Sci USA* 77: 5221–5225
- Smith RJ, Lodato RF, Valle DL, Kazakis A (1981) Mutant cell lines resistant to azetidine carboxylic acid: quantitative and qualitative differences in pyrroline-5-carboxylate synthase activities. *Biochem Biophys Res Commun* 99: 789–795
- Splittstoesser WE, Fowden L (1973) Ornithine transaminase from *Cucurbita maxima* cotyledons. *Phytochemistry* 12: 785–790
- Stetten MR (1951) Mechanism of the conversion of ornithine into proline and glutamic acid in vivo. *J Biol Chem* 189: 499
- Stewart CR (1981) Proline accumulation: biochemical aspects. In: Paleg LG, Aspinall D (eds) *The physiology and biochemistry of drought resistance in plants*. Academic, Sydney, pp 243–259
- Stewart CR, Boggess SE, Aspinall D, Paleg G (1977) Inhibition of proline oxidation by water stress. *Plant Physiol* 59: 930–932
- Stewart CR, Larher E (1980) Accumulation of amino acids and related compounds in relation to environmental stress. In: Mifflin BJ (ed) *The biochemistry of plants*, vol 5. Academic Press, New York, pp 609–635
- Stewart CR, Boggess SE (1978) Metabolism of [5-3H]proline by barley leaves and its use in measuring the effects of water stress on proline oxidation. *Plant Physiol* 61: 654–657
- Stewart PM, Batshaw M, Valle D, Walser M (1981) Effects of arginine-free meals on ureagenesis in cats. *Am J Physiol* 241: E310–E315
- Strecker HJ (1960a) The interconversion of glutamic acid and proline II. The preparation and properties of Δ^1 -pyrroline-5-carboxylic acid. *J Biol Chem* 235: 2045–2050
- Strecker HJ (1960b) The interconversion of glutamic acid and proline III. Δ^1 -pyrroline 5-carboxylic acid dehydrogenase. *J Biol Chem* 235: 3218–3223
- Strom AR, Le Rudulier D, Jakowec MW, Bunnell RC, Valentine RC (1983) Osmoregulatory (Osm) genes and osmoprotective compounds. In: Kosuge T, Meredith CP, Hollaender A (eds) *Genetic engineering of plants. An agricultural perspective*. Plenum Press, New York, pp 39–59
- Szoke A, Miao G-H, Hong Z, Verma DPS (1992) Subcellular localization of Δ^1 -pyrroline-5-carboxylate reductase in root/nodule and leaf of soybean. *Plant Physiol* 99: 1642–1649
- Thompson JE (1980) Arginine synthesis, proline synthesis, and related processes. In: Mifflin BJ (ed) *The biochemistry of plants*, vol 5. Academic Press, New York, pp 375–402
- Tomenchok DM, Brandriss MC (1987) Gene-enzyme relationships in the proline biosynthetic pathway of *Saccharomyces cerevisiae*. *J Bacteriol* 169: 5364–5372
- Treichel S (1986) The influence of NaCl on Δ^1 -pyrroline-5-carboxylate reductase in proline-accumulating cell suspension cultures of *Mesembryanthemum nodiflorum* and other halophytes. *Plant Physiol* 67: 173–181
- Valle D, Simell O (1995) The hyperornithinaemias. In: Scriver C, Beaudet A, Sly W, Valle D (eds) *The metabolic and molecular bases of inherited disease*. McGraw Hill, New York, pp 1147–1185
- Valle D, Downing SJ, Harris SC, Phang JM (1973) Proline biosynthesis: multiple defects in Chinese hamster ovary cells. *Biochem Biophys Res Commun* 53: 1130–1136
- Valle D, Blaese RM, Phang JM (1975) Increased sensitivity of lymphocyte Δ^1 -pyrroline-5-carboxylate reductase to inhibition by proline with transformation. *Nature* 253: 214–216
- Valle D, Goodman SI, Applegarth DA, Shih VE, Phang JM (1976) Type II hyperprolinemia. Δ^1 -pyrroline-5-carboxylic acid dehydrogenase deficiency in cultured skin fibroblasts and circulating lymphocytes. *J Clin Invest* 58: 598–603
- Valle D, Kupfer-Kaiser MI, Del-Valle LA (1977) Gyrate atrophy of the choroid and retina: deficiency of ornithine aminotransferase in transformed lymphocytes. *Proc Natl Acad Sci USA* 74: 5159–5161

- Valle D, Walser M, Brusilow SW, Kupfer-Kaiser MI (1980) Gyrate atrophy of the choroid and retina: amino acid metabolism and correction of hyperornithinemia with an arginine-deficient diet. *J Clin Invest* 65: 371–378
- Verbruggen N, Villarroel R, Van Montagu M (1993) Osmoregulation of a pyrroline-5-carboxylate reductase gene in *Arabidopsis thaliana*. *Plant Physiol* 103: 771–781
- Verma DPS, Hu C-AA, Delauney AJ, Miao G-H, Hong Z (1992) Deciphering proline biosynthesis pathways in plants by direct, trans-, and co-complementation in bacteria. In: Singh BK, Flores HE, Shannon JC (eds) *Biosynthesis and molecular regulation of amino acids in plants*. American Society of Plant Physiologists, Rockville, MD, pp 128–138
- Volpe P, Sawamura R, Strecker HJ (1969) Control of ornithine Δ -transaminase in rat liver and kidney. *J Biol Chem* 244: 719–726
- Wakabayashi Y, Jones ME (1983) Pyrroline-5-carboxylate synthesis from glutamate by rat intestinal mucosa. *J Biol Chem* 258: 3865–3872
- Wakabayashi Y, Henslee JG, Jones ME (1983) Pyrroline-5-carboxylate synthesis from glutamate by rat intestinal mucosa. Subcellular localization and temperature stability. *J Biol Chem* 258: 3873–3882
- Wakabayashi Y, Yamada R, Iwashima A (1985) Temperature- and time-dependent inactivation of pyrroline-5-carboxylate synthase: suggestive evidence for an allosteric regulation of the enzyme. *Arch Biochem Biophys* 238: 469–475
- Wakabayashi Y, Yamada E, Hasegawa T, Yamada R (1991) Enzymological evidence for the indispensability of small intestine in the synthesis of arginine from glutamate. *Arch Biochem Biophys* 291: 1–8
- Wang T, Lawler AM, Steel G, Sipila I, Milam AH, Valle D (1995) Mice lacking ornithine- δ -amino-transferase have paradoxical neonatal hypoornithinemia and retinal degeneration. *Nature Genet* 11: 185–190
- Williamson CL, Slocum RD (1992) Molecular cloning and evidence for osmoregulation of the Δ 1-pyrroline-5-carboxylate (proC) gene in pea (*Pisum sativum* L.). *Plant Physiol* 100: 1464–1470
- Windmueller HG, Spaeth AE (1975) Intestinal metabolism of glutamine and glutamate from the lumen as compared to glutamine. *Arch Biochem Biophys* 171: 662–672
- Wu G, Knabe DA, Flynn NE (1994) Synthesis of citrulline from glutamine in pig enterocytes. *Biochem J* 299: 115–121
- Wu G, Flynn NE, Yan W, Barstow DG Jr (1995a) Glutamine metabolism in chick enterocytes: absence of pyrroline-5-carboxylase synthase and citrulline synthesis. *Biochem J* 306: 717–721
- Wu G, Knabe DA, Yan W, Flynn NE (1995b) Glutamine and glucose metabolism in enterocytes of the neonatal pig. *Am J Physiol* 268: R334–R342
- Yancey PH (1994) Compatible and counteracting solutes. In: Strange K (ed) *Cellular and molecular physiology of cell volume regulation*. CRC Press, Boca Raton, FL, pp 81–109
- Yamada E, Wakabayashi Y (1991) Development of pyrroline-5-carboxylate synthase and N-acetylglutamate synthase and their changes in lactation and aging. *Arch Biochem Biophys* 291: 15–23
- Yeh GC, Phang JM (1980) The function of pyrroline-5-carboxylate reductase in human erythrocytes. *Biochem Biophys Res Commun* 94: 450–457
- Yeh GC, Phang JM (1981) The stimulation of purine nucleotide production by pyrroline-5-carboxylic acid in human erythrocytes. *Biochem Biophys Res Commun* 103: 118–124
- Yeh GC, Phang JM (1983) Pyrroline-5-carboxylate stimulates the conversion of purine antimetabolites to their nucleotide forms by a redox-dependent mechanism. *J Biol Chem* 258: 9774–9779
- Yeh GC, Harris SC, Phang JM (1981) Pyrroline-5-carboxylate reductase in human erythrocytes. *J Clin Invest* 67: 1042–1046

- Yeh GC, Roth EF Jr, Phang JM, Harris SC, Nagel RL, Rinaldi A (1984) The effect of pyrroline-5-carboxylic acid on nucleotide metabolism in erythrocytes from normal and glucose-6-phosphate dehydrogenase-deficient subjects. *J Biol Chem* 259: 5454–5458
- Yip MC, Knox WE (1972) Function of arginase in lactating mammary gland. *Biochem J* 127: 893–899
- Zhang CS, Lu Q, Verma DP (1995) Removal of feedback inhibition of Δ^1 -pyrroline-5-carboxylate synthetase, a bifunctional enzyme catalyzing the first two steps of proline biosynthesis in plants. *J Biol Chem* 270: 20491–20496

Authors' address: Pr. P. Kamoun, Laboratoire de Biochimie B, Hôpital Necker-Enfants Malades, 149, Rue de Sèvres, F-75015 Paris, France.

Received February 1, 1997