

Generation of reactive oxygen species by beta amyloid fibrils and oligomers involves different intra/extracellular pathways

Giovanna Cenini · Cristina Cecchi · Anna Pensalfini ·
Sara Anna Bonini · Giulia Ferrari-Toninelli ·
Gianfranco Liguri · Maurizio Memo · Daniela Uberti

Published online: 1 September 2009
© Springer-Verlag 2009

Erratum to: Amino Acids (2009)
DOI 10.1007/s00726-009-0319-7

The original version of this article unfortunately contained a mistake.

The presentation of the first author's name was incorrect.

The corrected presentation is given below.

Giovanna Cenini, Cristina Cecchi, Anna Pensalfini, Sara Anna Bonini, Giulia Ferrari-Toninelli, Gianfranco Liguri, Maurizio Memo and Daniela Uberti.

The online version of the original article can be found under doi:[10.1007/s00726-009-0319-7](https://doi.org/10.1007/s00726-009-0319-7).

G. Cenini · S. A. Bonini · G. Ferrari-Toninelli · M. Memo ·
D. Uberti (✉)
Department of Biomedical Sciences and Biotechnologies,
University of Brescia, Viale Europa 11, 25124 Brescia, Italy
e-mail: uberti@med.unibs.it

C. Cecchi · A. Pensalfini · G. Liguri
Department of Biochemical Sciences, University of Florence,
Florence, Italy