

Dietary arginine supplementation enhances antioxidative capacity and improves meat quality of finishing pigs

Xianyong Ma · Yingcai Lin · Zongyong Jiang · Chuntian Zheng ·
Guilian Zhou · Deqian Yu · Ting Cao · Jun Wang · Fang Chen

Received: 1 September 2008 / Accepted: 11 November 2008 / Published online: 5 December 2008
© Springer-Verlag 2008

Abstract The present study was conducted to test the hypothesis that dietary arginine supplementation may improve meat quality of finishing pigs. Beginning at ~60 kg body weight, pigs were fed a corn- and soybean meal-based diet supplemented with 0, 0.5 or 1% L-arginine until they reached a body weight of ~110 kg. On the last day of the experiment, pigs were food-deprived for 16 h before blood samples were obtained for analysis of amino acids, insulin, and other metabolites. Immediately thereafter, pigs were slaughtered for determination of carcass composition, muscle biochemical parameters, and meat quality. The result showed that arginine did not affect pig growth performance or carcass traits. However, 1% arginine decreased drip loss of pork muscle at 48 h postmortem, while increasing intramuscular fat content ($P < 0.05$). Supplementing 0.5 or 1% arginine to the diet increased arginine concentration and decreased cortisol level in serum, while enhancing antioxidative capacity and glutathione peroxidase activity in serum ($P < 0.05$). Additionally, 1% arginine increased antioxidative capacity in skeletal muscle ($P < 0.05$). Furthermore, 0.5 or 1% arginine decreased the cortisol receptor mRNA level in muscle ($P < 0.05$). Collectively, these results indicate that supplemental arginine improved meat quality and attenuated oxidative stress of finishing pigs.

Keywords Arginine · Finishing pig · Meat quality · Antioxidant capacity · Stress

Abbreviations

CAT	Catalase
CR	Cortisol receptor
GSH-Px	Glutathione peroxidase
IGF-IR	Insulin-like growth factor-I receptor
MDA	Malondialdehyde
PPAR γ	Peroxisome proliferators-activated receptor γ
SOD	Superoxide dismutase
T-AOC	Total antioxidant capacity

Introduction

There is increasing interest in improving pork meat quality among producers and consumers. A number of studies have shown that dietary supplementation with certain nutrients may enhance water-holding capacity, color, and anti-oxidative capacity of pork muscle. These nutrients have antioxidant function and include magnesium, selenium, vitamin E, vitamin C, tryptophan, creatine, and conjugated linoleic acids (Buckley et al. 1995; Swigert et al. 2004). Available evidence shows that oxidative, heat, or slaughter stress can result in the oxidation of fat, protein and metal ions to generate malondialdehyde (MDA), carbonyl protein complexes, aldehydes, ketones, and phenolic compounds (Galli 2007; Fang et al. 2002). These products are known to affect the quality of pork and poultry meats (Lefaucheur et al. 1991; Sandercock et al. 2001). L-Arginine is a nutritionally important amino acid and plays multiple physiologic functions in animals (Jobgen et al. 2006;

X. Ma · Y. Lin · Z. Jiang (✉) · C. Zheng · G. Zhou · D. Yu ·
J. Wang
Institute of Animal Science, Guangdong Academy
of Agricultural Science, 510640 Guangzhou, Guangdong, China
e-mail: jiangz41@163.com

X. Ma · Y. Lin · Z. Jiang · T. Cao · F. Chen
Guangdong Public Laboratory of Animal Breeding
and Nutrition, 510640 Guangzhou, Guangdong, China

Mateo et al. 2007; Wang et al. 2008a; Wu et al. 2007a). One of these functions is to increase antioxidant ability, reduce superoxide release, and ameliorate lipid peroxidation (Galli 2007; Petrovic et al. 2008).

In animals, arginine was metabolized to nitric oxide, proline, glutamine and polyamines with enormous biological importance (Liao et al. 2008; Wu and Morris 1998). For example, physiological levels of these metabolites can attenuate the stress response (Hamasu et al. 2008; Suenaga et al. 2008a, b), enhance the immune function (Li et al. 2007b; Tan et al. 2008a), regulate protein synthesis (Tan et al. 2008b; Yao et al. 2008), and promote wound healing (Flynn et al. 2002). In view of the foregoing, we hypothesized that dietary arginine supplementation may improve meat quality of finishing pigs through increasing anti-oxidative capacity and ameliorating stress. The present study was conducted to test this hypothesis.

Materials and methods

Pigs and diet

A total of 60 Du × (Chang × Da) finishing pigs (male, castrated) of approximately 60 kg body weight (90 days of age) were used in this experiment. All pigs were housed in the animal facilities of the Institute of Animal Science, Guangdong Academy of Agricultural Science. They were allotted randomly to one of three dietary treatments on the basis of body weight in a completely randomized design. Each treatment consisted of five replicates with four pigs per replicate. Pigs in Group 1 (the control) were fed a corn- and soybean meal-based diet (Table 1), whereas pigs in Groups 2 and 3 were fed the basal diet supplemented with 0.5 and 1% L-arginine (>98% purity), respectively (Table 1). L-Alanine was used as isonitrogenous control, as described by Kim and Wu (2004). L-Arginine and L-alanine were purchased from Guangzhou Weijian Medicine and Healthy Foods Import–Export Company (Guangdong, China).

Feeding and slaughter procedure

The experiment was initiated when the pigs weighted ~60 kg (90 days of age) and ended when the pigs reached ~110 kg (170 days of age) body weight. At the beginning and the termination of the experiment, each pig was weighed. The amount of feed consumed was recorded during the experiment to determine average daily gain, average daily feed intake, and the gain to feed ratio. At the end of the experiment, 14 h after the last feeding, two pigs were selected randomly from each replicate to obtain blood

samples at 7:00 a.m. via the anterior vena cava using vacuum tubes (containing no anticoagulant), followed immediately by slaughter. Carcasses were weighed and split. The loin eye area and backfat thickness were measured. This study was carried out in accordance with the Chinese guidelines (Science and Technology Ministry of China 2006) for animal welfare and approved by Guangdong Academy of Agricultural Science.

Sample collection and analysis of biochemical parameters

Blood samples were centrifuged for 5 min at 3,000g, 4°C, and the serum was stored at −20°C for assays of biochemical parameters. Following slaughter, ~2 cm³ longissimus muscle (the caudal end of the muscle) was collected immediately and stored at −20°C for the determination of hydroxy radical, MDA, total antioxidant capacity (T-AOC), as well as activities of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and catalase (CAT), using commercial kits (Jiancheng Bioengineering Institution, Nanjing, China). Amino acids in serum were analyzed by Amino Acid Analyzer (L-8900, HITACHI, Japan), as described by Kong et al. (2008) and Yin et al. (2008).

Meat quality measurements

Muscle samples were taken at the last thoracic vertebra and the pH value was recorded at 45 min, 24 and 48 h postmortem using a pH meter (HI8242C, Beijing Hanna Instruments Science & Technology Co. Ltd., Beijing, China). Drip loss was estimated according to the method described by Carlo et al. (1999). Briefly, three cylindrical muscle cores were taken from each slice with a cork borer (25 mm diameter), weighed, wrapped in a plastic bag, and stored at 4°C. Samples were taken from the container and weighed after stored for 24 and 48 h. Drip loss was estimated by calculating the difference between the initial and the final weight of the muscle sample. Meat color CIE LAB value (L*, relative lightness; a*, relative redness; b*, relative yellowness) was assessed immediately on the fresh surface of meat samples using a colorimeter (CR-410, Minolta Co. Ltd, Suita-shi, Osaka, Japan) according to the method described by Mason et al. (2005). Each sample was a complete cross section of thoraco-lumbar vertebra, about 2-cm thickness. Shear force was determined using an Instron Universal Mechanical Machine (Instron model 4411, Instron Corp., Canton) as follows. Approximately 250 g longissimus muscle sample from each pig was refrigerated overnight at 4°C and then brought to room temperature before

Table 1 Dietary composition and nutrient levels

	Control	Supplementation with 0.5% arginine	Supplementation with 1% arginine
Diet ingredients			
Corn meal	70.82	71.3	71.8
Soybean meal	17.6	17.6	17.6
Wheat bran	3	3	3
Rap meal	4	4	4
Soybean oil	0.2	0.2	0.2
L-Ala	1.84	0.97	0
L-Arg	0	0.5	1
Premix ^a	1	1	1
L-Lysine HCl	0.12	0.12	0.12
Lime powder	1	1.01	1.02
Salt	0.3	0.3	0.29
Dicalcium phosphate	0.3	0.3	0.3
Total	100	100	100
Nutrient levels			
Digestible energy (Mcal/kg)	3.17	3.19	3.20
CP (%)	15.4	15.4	15.4
Lys (%)	0.8	0.8	0.8
Met (%)	0.24	0.24	0.24
Met + Cys (%)	0.50	0.50	0.50
Thr (%)	0.58	0.58	0.58
Trp (%)	0.17	0.17	0.17
Arg (%)	0.96	1.46	1.96
Val (%)	0.70	0.70	0.70
Ca (%)	0.55	0.55	0.55
Total P (%)	0.42	0.42	0.42
Available P (%)	0.19	0.19	0.19
Na (%)	0.15	0.15	0.15
Cl (%)	0.24	0.24	0.24

^a Providing 1,750 IU/kg vitamin A, 220 IU/kg vitamin D₃, 3 IU/kg vitamin E, 0.55 mg/kg vitamin K₃, 0.25 mg/kg vitamin B₁, 1.0 mg/kg vitamin B₂, 0.7 mg/kg vitamin B₆, 3 µg/kg vitamin B₁₂, 4 mg/kg niacin, 1.6 mg/kg calcium pantothenate, 0.1 mg/kg folic acid, 7 µg/kg biotin, 0.08 g/kg choline chloride, 6.5 mg/kg manganese, 15 mg/kg iron, 15 mg/kg zinc, 1.5 mg/kg copper, 0.07 mg/kg iodine(I₂), 0.03 mg/kg selenium, and 1 g/kg sodium chloride

cooking. The temperature of the water bath was set at 80°C and meat samples were cooked for 1 h to an internal temperature of 70°C, and then cooled to room temperature. About 10 slices of 1 cm × 1.5 cm were cut perpendicular to the fiber orientation of the muscle. The marbling of meat sample (thoraco-lumbar vertebra section) was assessed by a marbling testing standard (NPPC 1991) after being stored at 4°C for 24 h. Intramuscular fat (IMF) content was determined as follows. Approximately 10 g longissimus muscle sample from each pig was cut into pieces and dried in a freezing drier, ground into powder, and extracted for lipids with petroleum ether extract using the Soxtec 2055 fat extraction system (Foss Tecator AB, Box 70, S-26321 Höganäs, Sweden).

Real-time PCR analysis

Total RNA was isolated from fresh longissimus muscle tissue using TRIZOL reagent (Invitrogen Co., USA) according to the manufacturer's instructions. The RNA concentration was determined at 260/280 nm and the cDNA was produced using a commercial kit containing Reverse Transcriptase XL (AMV) and RNasin (Invitrogen Co., USA). Real-time PCR was performed using an ABI 7500 Mastercycler (Applied Biosystems, USA) with qPCR Mix (TaKaRa, BIOINC, Japan) according to the manufacturers' instructions. The DNA was sequenced by Sangon Technical Co. Ltd. (Shanghai, China). Primers and probes (Table 2) of genes [β -actin, cortisol receptor (CR), leptin, IGF-IR, and PPAR- γ]

Table 2 PCR primers for genes

Gene	Primers	
CR	Forward primer	5'-cctgccctggtagtcacg-3'
	Reverse primer	5'-agcgtcttgagggtttgtg-3'
PPAR γ	Forward primer	5'-ctgaaccagagtctgctgac-3'
	Reverse primer	5'-ggtgattgtctgtggtcttcc-3'
IGF-IR	Forward primer	5'-acctgccctggtagtcacg-3'
	Reverse primer	5'-tagcgtcttgagggtttgtg-3'
Leptin	Forward primer	5'-ggctcccaggtgctgtt-3'
	Reverse primer	5'-gaacctggcccttcgagatc-3'
β -Actin	Forward primer	5'-gacctgaccgactacatc-3'
	Reverse primer	5'-gccgatggtgatgacctggc-3'

were designed according to the sequences published in GenBank. The real-time PCR cycling conditions used were 95°C for 40 s, followed by 40 cycles at 95°C for 5 s and 58°C for 20 s, then 1 cycle at 72°C for 30 s.

Statistical analysis

Values are expressed as mean $\pm$ SEM. Data were analyzed by one-way analysis of variance and the Student–Newman–Keuls multiple comparison test. All statistical analyses were performed using SAS (v6.12, SAS Institute Inc., Cary, NC, USA). $P < 0.05$ was taken to indicate statistical significance.

Results

Growth performance and carcass traits

The effects of dietary arginine supplementation on growth performance and carcass traits of the pigs are summarized in (Table 3). Supplementation with arginine did not affect ($P > 0.05$) average daily gain, average daily feed intake, feed to gain ratio, final body weight, carcass yield, lean meat, percentage of lean tissue, loin eye area, or backfat thickness.

Meat quality

There was no difference in meat quality traits (e.g., pH value and meat color) between the control and arginine-supplemented pigs (Table 4). The drip loss at 48 h post-mortem was lower ($P < 0.05$) in pigs supplemented with 1% arginine when compared with the control pigs, but there was no difference ($P > 0.05$) between the control group and the pigs supplemented with 0.5% arginine. The shear force of meat did not differ ($P > 0.05$) among the three groups of pigs, but had a trend to decrease in

Table 3 Effect of dietary arginine supplementation on growth performance and carcass traits of finishing pigs

Treatment	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
ADG (g/d)	786 $\pm$ 3	792 $\pm$ 6	764 $\pm$ 10
ADFI (kg/d)	3.06 $\pm$ 0.13	2.89 $\pm$ 0.10	2.98 $\pm$ 0.10
F:G	3.90 $\pm$ 0.12	3.64 $\pm$ 0.18	3.90 $\pm$ 0.17
Live weight (kg)	118 $\pm$ 2	120 $\pm$ 3	121 $\pm$ 3
Carcass yield (%)	74.7 $\pm$ 0.6	74.1 $\pm$ 0.4	74.3 $\pm$ 0.4
Lean meat (kg)	26.3 $\pm$ 1.4	26.8 $\pm$ 0.7	27.2 $\pm$ 1.0
Lean percentage (%)	63.8 $\pm$ 1.7	65.0 $\pm$ 1.5	63.5 $\pm$ 1.8
Loin eye area (mm ²)	5375 $\pm$ 259	5461 $\pm$ 184	5459 $\pm$ 87
Backfat depth (cm)			
First rib	3.87 $\pm$ 0.35	4.00 $\pm$ 0.15	4.10 $\pm$ 0.22
Tenth rib	1.99 $\pm$ 0.15	2.06 $\pm$ 0.22	2.14 $\pm$ 0.02
Last rib	1.21 $\pm$ 0.16	1.24 $\pm$ 0.18	1.29 $\pm$ 0.06

Data are mean $\pm$ SEM, $n = 20$ for ADG, ADFI and F:G and $n = 10$ for other parameters

ADG average daily gain, ADFI average daily feed intake, F:G feed to gain ratio

Table 4 Effect of dietary arginine supplementation on meat quality of finishing pigs

Treatment	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
pH			
45 min	6.44 $\pm$ 0.07	6.52 $\pm$ 0.07	6.61 $\pm$ 0.06
24 h	5.57 $\pm$ 0.1	5.58 $\pm$ 0.07	5.65 $\pm$ 0.04
48 h	5.46 $\pm$ 0.07	5.46 $\pm$ 0.08	5.50 $\pm$ 0.05
Drip loss (%)			
24 h	1.21 $\pm$ 0.09	1.10 $\pm$ 0.10	1.02 $\pm$ 0.07
48 h	2.32 $\pm$ 0.14 ^a	2.02 $\pm$ 0.11 ^{ab}	1.67 $\pm$ 0.14 ^b
L			
0	45.4 $\pm$ 1.5	46.2 $\pm$ 0.3	46.7 $\pm$ 0.9
24 h	53.6 $\pm$ 1.0	53.6 $\pm$ 0.7	55.3 $\pm$ 1.1
48 h	53.8 $\pm$ 0.4	54.3 $\pm$ 0.7	55.0 $\pm$ 0.8
a*			
0	14.9 $\pm$ 0.2	16.1 $\pm$ 0.9	14.7 $\pm$ 0.7
24 h	15.0 $\pm$ 0.3	16.1 $\pm$ 0.4	14.9 $\pm$ 0.5
48 h	15.6 $\pm$ 0.3	16.8 $\pm$ 0.6	15.0 $\pm$ 0.8
b*			
0	3.60 $\pm$ 0.42	4.02 $\pm$ 0.37	3.67 $\pm$ 0.29
24 h	2.86 $\pm$ 0.29	3.22 $\pm$ 0.41	3.14 $\pm$ 0.28
48 h	3.05 $\pm$ 0.24	3.59 $\pm$ 0.47	3.06 $\pm$ 0.31
Shear force (Nf)	47.8 $\pm$ 2.7	42.7 $\pm$ 1.9	42.8 $\pm$ 3.1
Intramuscular fat (%)	1.31 $\pm$ 0.14 ^b	1.99 $\pm$ 0.18 ^a	1.74 $\pm$ 0.17 ^a

Data are mean $\pm$ SEM, $n = 10$. Within a row, means with different superscript letters differ ($P < 0.05$)

L lightness, a* redness, b* yellowness

Table 5 Effect of dietary arginine supplementation on concentrations of biochemical parameters in blood of finishing pigs

Treatment	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
Cholesterol (mmol/L)	2.10 ± 0.10	2.03 ± 0.09	2.02 ± 0.04
Free fatty acid (mmol/L)	0.27 ± 0.02	0.26 ± 0.03	0.25 ± 0.02
Urea N (mmol/L)	5.40 ± 0.32	5.52 ± 0.28	5.67 ± 0.22
Glucose (mmol/L)	3.73 ± 0.22	4.05 ± 0.30	3.99 ± 0.23
Low density lipoprotein (mmol/L)	0.29 ± 0.01	0.28 ± 0.01	0.28 ± 0.01
Triglycerides (mmol/L)	0.46 ± 0.04	0.49 ± 0.03	0.47 ± 0.03
Insulin (μIU/ml)	28.6 ± 2.2	24.0 ± 1.0	27.3 ± 2.5
Phosphokinase(IU/L)	41.3 ± 1.8	35.2 ± 2.6	35.0 ± 3.0
Alkaline phosphatase (IU/L)	86.0 ± 6.8	86.8 ± 6.4	88.6 ± 6.2
Lactate dehydrogenase (IU/L)	494 ± 25	448 ± 15	430 ± 24
Cortisol (ng/ml)	89.4 ± 0.8 ^a	56.3 ± 0.6 ^b	53.5 ± 0.5 ^b

Data are mean ± SEM, $n = 10$. Within a row, means with different superscript letters differ ($P < 0.05$)

arginine-supplemented pigs. The IMF content was higher ($P < 0.05$) in pigs supplemented with 0.5 and 1% arginine than in the control group and did not differ ($P > 0.05$) between the 0.5 and 1% arginine groups (Table 4).

Biochemical parameters in serum

Dietary supplementation with arginine decreased ($P < 0.05$) the concentration of cortisol in serum but did not affect ($P > 0.05$) concentrations of cholesterol, free fatty acid, glucose, urea, low density lipoprotein, triglycerides or insulin in serum (Table 5). The activities of phosphokinase, alkaline phosphatase, and lactate dehydrogenase in serum did not differ ($P > 0.05$) between control and arginine-supplemented pigs. Compared with the control group, supplementing 0.5 and 1% arginine to the basal diet increased ($P < 0.05$) the concentrations of arginine, proline, as well as glutamine plus glutamate in the serum of pigs, but had no effect ($P > 0.05$) on the concentration of lysine in serum (Table 6).

MDA production and enzyme activities

Dietary arginine supplementation did not affect ($P > 0.05$) the concentration of MDA or the activities of SOD and CAT in serum or skeletal muscle of pigs (Table 7). However, supplementation with 1% arginine decreased the level of hydroxy radical in the serum and muscle. Supplementation with 0.5 and 1% arginine increased ($P < 0.05$) the activities of T-AOC and GSH-Px in serum, while supplementation with 1% arginine augmented ($P < 0.05$) T-AOC level in the muscle.

Gene expression

The results of gene sequencing indicated 100% homology of the fragments obtained from the present study with the sequences published in GenBank. The mRNA levels for

Table 6 Effect of dietary arginine supplementation on concentrations of amino acids in serum of finishing pigs

Treatment	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
Arginine	0.20 ± 0.01 ^b	0.26 ± 0.03 ^a	0.29 ± 0.02 ^a
Glutamine	0.51 ± 0.01 ^b	0.60 ± 0.02 ^a	0.62 ± 0.04 ^a
Glutamate	0.22 ± 0.01 ^b	0.29 ± 0.01 ^a	0.26 ± 0.01 ^{ab}
Proline	0.23 ± 0.02 ^b	0.25 ± 0.02 ^{ab}	0.28 ± 0.03 ^a
Lysine	0.21 ± 0.03	0.23 ± 0.02	0.24 ± 0.01

Data are mean ± SEM (mmol/L), $n = 10$. Within a row, means with different superscript letters differ ($P < 0.05$)

cortisol receptor (CR), insulin-like growth factor-I receptor (IGF-IR), leptin, and peroxisome proliferators-activated receptor γ (PPAR γ) genes in skeletal muscle are summarized in Table 8. The CR mRNA level in the muscle of pigs supplemented with 0.5 and 1% arginine was lower ($P < 0.05$) than that in the control group. The PPAR γ mRNA level in the skeletal muscle of pigs supplemented with 1% arginine was higher ($P < 0.05$) than that in the control group and the pigs supplemented with 0.5% arginine. The level of leptin mRNA was not different ($P > 0.05$) among the three groups of pigs.

Discussion

L-Arginine is the physiological substrate of nitric oxide, polyamines, creatine, agmatine, glutamate, and proline with enormous biological importance (Krane 2008; Montanez et al. 2008; Wu and Morris 1998). Increasing evidence indicates that arginine regulates the partitioning of dietary energy in favor of muscle protein accretion and fat reduction in animals (Fu et al. 2005; Wu et al. 2007c, 2008a, b). Additionally, either dietary supplementation

Table 7 Effect of dietary arginine supplementation on antioxidant enzyme activities and MDA content in serum and skeletal muscle of finishing pigs

Treatment	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
Serum			
Hydroxy radical (U/ml)	265 ± 31 ^a	211 ± 16 ^{ab}	180 ± 20 ^b
MDA (nmol/ml)	3.94 ± 0.28	3.46 ± 0.59	3.52 ± 0.41
T-AOC (U/ml)	2.78 ± 0.24 ^b	3.54 ± 0.76 ^a	3.98 ± 0.71 ^a
SOD (U/ml)	120 ± 8	130 ± 10	133 ± 9
GSH-Px (U/ml)	580 ± 31 ^b	664 ± 46 ^a	658 ± 27 ^a
CAT (U/ml)	563 ± 31	629 ± 34	635 ± 49
Muscle			
Hydroxy radical (U/mg protein)	95.5 ± 1.7 ^a	93.8 ± 2.2 ^a	79.0 ± 1.6 ^b
MDA (nmol/mg protein)	3.34 ± 0.17	2.68 ± 0.22	2.72 ± 0.27
T-AOC (U/mg protein)	1.29 ± 0.10 ^b	1.61 ± 0.09 ^{ab}	1.90 ± 0.06 ^a
SOD (U/mg protein)	56.3 ± 3.1	62.8 ± 3.0	63.5 ± 4.7
GSH-Px (U/mg protein)	51.3 ± 4.5	62.8 ± 4.1	59.3 ± 2.2
CAT (U/mg protein)	21.5 ± 1.0	24.6 ± 1.3	23.0 ± 1.7

Data are mean ± SEM, $n = 10$.

Within a row, means with different superscript letters differ ($P < 0.05$)

Table 8 Effect of dietary arginine supplementation on genes expression in skeletal muscle of finishing pigs

Item	Dietary supplementation with L-arginine (%)		
	0.0	0.5	1.0
CR/ β -actin	6.58 ± 0.33 ^a	1.78 ± 0.15 ^b	1.46 ± 0.12 ^b
IGF-IR/ β -actin	235 ± 26	376 ± 45	455 ± 47
Leptin/ β -actin	2.91 ± 0.34	4.24 ± 0.47	3.89 ± 0.63
PPAR γ / β -actin	183 ± 15 ^b	263 ± 21 ^{ab}	461 ± 39 ^a

Data are mean ± SEM, $n = 10$. Within a row, means with different superscript letters differ ($P < 0.05$)

with L-arginine (Kim and Wu 2004, 2008; Yao et al. 2008) or activation of endogenous arginine synthesis (Frank et al. 2007; Wu et al. 2004b) enhanced average daily weight gain in young pigs, compared with control piglets. Of particular interest, increasing the dietary arginine:lysine ratio increased the growth performance of heat-stressed broilers (Chen et al. 2005). Thus, it is important to determine whether dietary supplementation with arginine could improve the meat quality of pigs.

This study was not designed to determine the effect of dietary arginine supplementation on growth performance in swine. Thus, when body weight was used as the criterion to terminate the experiment, no significant differences in pig growth performance or carcass traits were observed (Table 3). However, when growing-finishing pigs were fed a diet supplemented with 1% L-arginine between 110 and 170 days of age, daily weight gain was enhanced and body fat content reduced, compared with the control group of pigs without arginine supplementation (Tan et al. 2008b). This discrepancy between the present study and the work of Tan et al. (2008b) may be explained by the difference in

experimental design, the length of arginine supplementation, and the breed of the pigs used.

Arginine is synthesized from glutamine and proline in most mammals, including pigs (Wu and Knabe 1995; Wu 1997) and humans (Hu et al. 2008a, b). However, increasing extracellular concentrations of arginine regulate the metabolism of protein, glucose and lipids in favor of lean tissue gain and white-fat reduction in animals (Jobgen et al. 2006; Wu et al. 2008a, b). Therefore, as reported by Tan et al. (2008b), we found that supplementing arginine to the diet of growing-finishing pigs decreased early muscle pH decline postmortem and drip losses. This beneficial result can be explained by lower rates of glycogenolysis and glycolysis in skeletal muscle of arginine-treated pigs compared with the control group (Tan et al. 2008b). On the basis of the previous reports that drip loss was positively related with oxidation (Swigert et al. 2004) and that intramuscular administration of zinc metallothionein to preslaughter-stressed pigs improved anti-oxidative status and pork quality (Li et al. 2007a), we surmise that the reduction of muscle drip loss in arginine-supplemented pigs may result from an improved anti-oxidative status in the tissue.

In support of the above proposition, available evidence shows that physiological levels of arginine and nitric oxide have antioxidative function (Wu and Meininger 2008). For example, Wascher et al. (1997) reported that administration of L-arginine diminished superoxide release and copper-induced lipid peroxidation in rats. Recently, Petrovic et al. (2008) demonstrated that L-arginine increased the antioxidative defense system in rats in response to cold acclimation. Importantly, dietary supplementation with arginine decreased the hydroxyl radical level in serum, while increasing T-AOC concentrations in serum and

skeletal muscle as well as the activity of GSH-Px [an antioxidative enzyme (Wu et al. 2004a)] in serum of pigs (Table 7), indicating the enhancement of whole-body antioxidative function. In addition, proline, glutamine, creatine and polyamines were important metabolites of arginine (Hu et al. 2008a, b; Phang et al. 2008; Wu and Morris 1998). Accordingly, dietary supplementation with 1% arginine increased serum concentrations of proline and glutamine + glutamate (Table 6), as well as creatine (He et al. 2008). It is noteworthy that proline attenuates the stress response in the central nervous system of chicks (Hamasu et al. 2008), while reducing inflammatory responses and oxidative stress in mammals (Bassit et al. 2008; Gualano et al. 2008). Similarly, glutamine (Wang et al. 2008b) and polyamines (Rider et al. 2007) enhance anti-oxidative function in cells.

Cortisol, the major circulating glucocorticoid in pigs, regulates protein and amino acid metabolism (Flynn et al. 2008). Another novel and important finding of the present study is that dietary arginine supplementation decreased the concentration of cortisol in serum and the mRNA level of glucocorticoid receptor in skeletal muscle (Table 8). This result suggests a reduction of stress levels in arginine-supplemented pigs, as concluded by other investigators (Han et al. 2008). Similarly, Suenaga et al. (2008a, b) reported that arginine reduced stress and induced sedative effects in chickens.

As reported by Tan et al. (2008b), we found that dietary supplementation with 1% arginine increased IMF (Table 4) but had no effect on serum concentrations of glucose, lipids or insulin (Table 5). Interestingly, the arginine treatment enhanced the PPAR γ mRNA level in skeletal muscle (Table 8). This finding can be explained by the fact that PPAR γ stimulates fat synthesis in this tissue (Hsu and Ding 2003), which can explain the effect of arginine observed from the current study. In contrast, leptin in muscle has been reported to decrease the activity of acetyl-CoA carboxylase, a key regulatory enzyme in fatty acid synthesis (Steinberg et al. 2003). Consistent with this view, we found that leptin mRNA level in muscle was lowered in arginine-supplemented pigs (Table 8). Clearly, future studies are needed to determine changes of fatty acid synthesis and oxidation, as well as the signaling actions of leptin and nitric oxide [another regulator of lipid metabolism (Jobgen et al. 2006)] in skeletal muscle of arginine-supplemented pigs.

Edmonds and Baker (1987) reported that arginine had an antagonistic effect on lysine absorption, but other investigators did not observe such a phenomenon (Tan et al. 2008a; Yao et al. 2008). This discrepancy may be explained by the differences in the contents of amino acids, including basic amino acids (Wu et al. 2008a, b). Studies with pigs have shown that dietary supplementation with

<2% arginine (on dry matter basis) is generally safe and does not result in an antagonism among basic amino acids (Wu et al. 2007b). In support of this view, we found that supplementing 0.5 and 1% arginine to the basal diet increased the concentration of arginine in the serum of pigs but had no effect on that of lysine (Table 6).

In summary, results of the present study demonstrate that dietary arginine supplementation reduced stress levels, enhanced whole-body antioxidative function, and improved the meat quality of finishing pigs. The arginine treatment increased IMF content and reduced muscle drop loss. Further research is required to elucidate the underlying molecular mechanisms.

Acknowledgments This study was supported by a grant (2004CB117500) from the “973” plan of China. We thank the staff of Institute of Animal Science, Guangdong Academy of Agricultural Science the Guangdong and Public Laboratory of Animal Breeding and Nutrition for technical assistance.

References

- Bassit RA, Curi R, Costa Rosa LFBP (2008) Creatine supplementation reduces plasma levels of pro-inflammatory cytokines and PGE2 after a half-ironman competition. *Amino Acids* 35:425–431
- Buckley DJ, Morrissey PA, Gray JI (1995) Influence of dietary vitamin E on the oxidative stability and quality of pig meat. *J Anim Sci* 73:3122–3130
- Carlo C, Giovannangelo O, Luigina P et al (1999) Influence of dietary vitamin e supplementation on “heavy” pig carcass characteristics, meat quality, and vitamin e status. *J Anim Sci* 77:1755–1761
- Chen J, Li X, Balnave D, Brake J (2005) The Influence of dietary sodium chloride, arginine: lysine ratio, and methionine source on apparent ileal digestibility of arginine and lysine in acutely heat-stressed broilers. *Poult Sci* 84:294–297
- Edmonds MS, Baker DH (1987) Failure of excess dietary lysine to antagonize arginine in young pigs. *J Nutr* 117:1396–1401
- Fang YZ, Yang S, Wu G (2002) Free radicals, antioxidants, and nutrition. *Nutrition* 18:872–879
- Flynn NE, Meiningner CJ, Haynes TE, Wu G (2002) The metabolic basis of arginine nutrition and pharmacotherapy. *Biomed Pharmacother* 56:427–438
- Flynn NE, Bird JG, Guthrie AS (2008) Glucocorticoid regulation of amino acid and polyamine metabolism in the small intestine. *Amino Acids*. doi:10.1007/s00726-008-0206-7
- Frank JW, Escobar J, Hguyen HV et al (2007) Oral *N*-carbamylglutamate supplementation increases protein synthesis in skeletal muscle of piglets. *J Nutr* 137:315–319
- Fu WJ, Haynes TE, Kohli R et al (2005) Dietary L-arginine supplementation reduces fat mass in Zucker diabetic fatty rats. *J Nutr* 135:714–721
- Galli F (2007) Amino acid and protein modification by oxygen and nitrogen species. *Amino Acids* 32:497–499
- Gualano B, Novaes RB, Artioli GG et al (2008) Effects of creatine supplementation on glucose tolerance and insulin sensitivity in sedentary healthy males undergoing aerobic training. *Amino Acids* 34:245–250
- Hamasu K, Haraguchi T, Kabuki Y et al. (2008) L-Proline is a sedative regulator of acute stress in the brain of neonatal chicks. *Amino Acids*. doi:10.1007/s00726-008-0164-0

- Han J, Liu YL, Fan W et al. (2008) Dietary L-arginine supplementation alleviates immunosuppression induced by cyclophosphamide in weaned pigs. *Amino Acids*. doi:[10.1007/s00726-008-0184-9](https://doi.org/10.1007/s00726-008-0184-9)
- He QH, Kong XF, Wu G et al. (2008) Metabolomic analysis of the response of growing pigs to dietary L-arginine supplementation. *Amino Acids*. doi:[10.1007/s00726-008-0192-9](https://doi.org/10.1007/s00726-008-0192-9)
- Hsu JM, Ding ST (2003) Effect of polyunsaturated fatty acids on the expression of transcription factor adipocyte determination and differentiation-dependent factor 1 and of lipogenic and fatty acid oxidation enzymes in porcine differentiating adipocyte. *Br J Nutr* 90:507–513
- Hu CA, Khalil S, Zhaorigetu S et al (2008a) Human Δ^1 -pyrroline-5-carboxylate synthase: function and regulation. *Amino Acids* 35:665–672
- Hu CA, Williams DB, Zhaorigetu S et al (2008b) Functional genomics and SNP analysis of human genes encoding proline metabolic enzymes. *Amino Acids* 35:655–664
- Jobgen WS, Fried SK, Fu WJ et al (2006) Regulatory role for the arginine-nitric oxide pathway in metabolism of energy substrates. *J Nutr Biochem* 17:571–588
- Kim SW, Wu G (2004) Dietary arginine supplementation enhances the growth of milk-fed young pigs. *J Nutr* 134:625–630
- Kim SW, Wu G (2008) Regulatory role for amino acids in mammary gland growth and milk synthesis. *Amino Acids*. doi:[10.1007/s00726-008-0151-5](https://doi.org/10.1007/s00726-008-0151-5)
- Kong XF, Yin YL, He QH et al. (2008) Dietary supplementation with Chinese herbal powder enhances ileal digestibilities and serum concentrations of amino acids in young pigs. *Amino Acids*. doi:[10.1007/s00726-008-0176-9](https://doi.org/10.1007/s00726-008-0176-9)
- Krane SM (2008) The importance of proline residues in the structure, stability and susceptibility to proteolytic degradation of collagens. *Amino Acids* 35:703–710
- Leflaucheur L, Dividich JL, Mourou J et al (1991) Influence of environmental temperature on growth, muscle and adipose tissue metabolism, and meat quality in swine. *J Anim Sci* 69:2844–2854
- Li LL, Hou ZP, Yin YL et al (2007a) Intramuscular administration of zinc metallothionein to preslaughter stress pigs improves anti-oxidative status and pork quality. *Asian Aust J Anim Sci* 20:761–767
- Li P, Yin YL, Li DF et al (2007b) Amino acids and immune function. *Br J Nutr* 98:237–252
- Liao XH, Majithia A, Huang XL, Kimmel AR (2008) Growth control via TOR kinase signaling, an intracellular sensor of amino acids and energy availability, with crosstalk potential to proline metabolism. *Amino Acids* 35:761–770
- Mateo RD, Wu G, Bazer FW et al (2007) Dietary L-arginine supplementation enhances the reproductive performance of gilts. *J Nutr* 137:652–656
- Montanez RC, Rodriguez-Caso C, Sanchez-Jimenez F, Medina MA (2008) In silico analysis of arginine catabolism as a source of nitric oxide or polyamines in endothelial cells. *Amino Acids* 34:223–229
- NPPC (1991) Procedures to evaluate market hogs. The 3rd National Pork Producers Council Des Moines, USA
- Petrovic V, Buzadzic B, Korac A et al (2008) Antioxidative defence alterations in skeletal muscle during prolonged acclimation to cold: role of L-arginine/NO-producing pathway. *J Exp Biol* 211:114–120
- Phang JM, Donald SP, Pandhare J, Liu YM (2008) The metabolism of proline, a stress substrate, modulates carcinogenic pathways. *Amino Acids* 35:681–690
- Rider JE, Hacker A, Mackintosh CA et al (2007) Spermine and spermidine mediate protection against oxidative damage caused by hydrogen peroxide. *Amino Acids* 33:231–240
- Sandercock DA, Hunter RR, Nute GR et al (2001) Acute heat stress-induced alterations in blood acid-base status and skeletal muscle membrane integrity in broiler chickens at two ages: implications for meat quality. *Poult Sci* 80:418–425
- Science and Technology Ministry of China (2006) The guiding suggestion about treating experimental animals amicably. Document no. 398
- Steinberg GR, Rush JW, Dyck DJ (2003) AMPK expression and phosphorylation are increased in rodent muscle after chronic leptin treatment. *Am J Physiol Endocrinol Metab* 284:E648–E654
- Suenaga R, Tomonaga S, Yamane H et al (2008a) Intracerebroventricular injection of L-arginine induces sedative and hypnotic effects under an acute stress in neonatal chicks. *Amino Acids* 35:139–146
- Suenaga R, Yamane H, Tomonaga S et al (2008b) Central L-arginine reduced stress responses are mediated by L-ornithine in neonatal chicks. *Amino Acids* 35:107–113
- Swigert KS, Mckeith FK, Carr TC et al (2004) Effects of dietary vitamin D3, vitamin E, and magnesium supplementation on pork quality. *Meat Sci* 67:81–86
- Tan BE, Li XG, Kong XF et al. (2008a) Dietary L-arginine supplementation enhances the immune status in early-weaned piglets. *Amino Acids*. doi:[10.1007/s00726-008-0155-1](https://doi.org/10.1007/s00726-008-0155-1)
- Tan BE, Yin YL, Liu ZQ et al. (2008b) Dietary L-arginine supplementation increases muscle gain and reduces body fat mass in growing-finishing pigs. *Amino Acids*. doi:[10.1007/s00726-008-0148-0](https://doi.org/10.1007/s00726-008-0148-0)
- Wang WW, Qiao SY, Li DF (2008a) Amino acids and gut function. *Amino Acids*. doi:[10.1007/s00726-008-0152-4](https://doi.org/10.1007/s00726-008-0152-4)
- Wang JJ, Chen LX, Li P et al (2008b) Gene expression is altered in piglet small intestine by weaning and dietary glutamine supplementation. *J Nutr* 138:1025–1032
- Wascher TC, Posch K, Wallner S (1997) Vascular effects of L-arginine: anything beyond a substrate for the NO-synthase? *Biochem Biophys Res Commun* 234:35–38
- Wu G (1997) Synthesis of citrulline and arginine from proline in enterocytes of postnatal pigs. *Am J Physiol Gastrointest Liver Physiol* 272:G1382–G1390
- Wu G, Knabe DA (1995) Arginine synthesis in enterocytes of neonatal pigs. *Am J Physiol Regulatory Integrative Comp Physiol* 269:R621–R629
- Wu G, Morris SM (1998) Arginine metabolism: nitric oxide and beyond. *Biochem J* 336:1–17
- Wu G, Meininger CJ (2008) Nitric oxide and vascular insulin resistance. *BioFactors* (in press)
- Wu G, Fang YZ, Yang S et al (2004a) Glutathione metabolism and its implications for health. *J Nutr* 134:489–492
- Wu G, Knabe DA, Kim SW (2004b) Arginine nutrition in neonatal pigs. *J Nutr* 134:2390S–2783S
- Wu G, Bazer FW, Davis TA et al (2007a) Important roles for the arginine family of amino acids in swine nutrition and production. *Livest Sci* 112:8–22
- Wu G, Bazer FW, Cudd TA et al (2007b) Pharmacokinetics and safety of arginine supplementation in animals. *J Nutr* 137:1673S–1680S
- Wu G, Collins JK, Perkins-Veazie P et al (2007c) Dietary supplementation with watermelon pomace juice enhances arginine availability and ameliorates the metabolic syndrome in Zucker diabetic fatty rats. *J Nutr* 137:2680–2685
- Wu G, Bazer FW, Datta S et al (2008a) Proline metabolism in the conceptus: Implications for fetal growth and development. *Amino Acids* 35:691–702
- Wu G, Bazer FW, Davis TA et al. (2008b) Arginine metabolism and nutrition in growth, health and disease. *Amino Acids*. doi:[10.1007/s00726-008-0210-y](https://doi.org/10.1007/s00726-008-0210-y)
- Yao K, Yin YL, Chu WY et al (2008) Dietary arginine supplementation increases mTOR signaling activity in skeletal muscle of neonatal pigs. *J Nutr* 138:867–872
- Yin FG, Liu YL, Yin YL et al. (2008) Dietary supplementation with Astragalus polysaccharide enhances ileal digestibilities and serum concentrations of amino acids in early weaned piglets. *Amino Acids*. doi:[10.1007/s00726-008-0142-6](https://doi.org/10.1007/s00726-008-0142-6)