

Reduction of protein radicals by GSH and ascorbate: potential biological significance

Janusz M. Gebicki · Thomas Nauser ·
Anastasia Domazou · Daniel Steinmann ·
Patricia L. Bounds · Willem H. Koppenol

Received: 30 November 2009 / Accepted: 23 April 2010 / Published online: 9 June 2010
© Springer-Verlag 2010

Abstract The oxidation of proteins and other macromolecules by radical species under conditions of oxidative stress can be modulated by antioxidant compounds. Decreased levels of the antioxidants glutathione and ascorbate have been documented in oxidative stress-related diseases. A radical generated on the surface of a protein can: (1) be immediately and fully repaired by direct reaction with an antioxidant; (2) react with dioxygen to form the corresponding peroxy radical; or (3) undergo intramolecular long range electron transfer to relocate the free electron to another amino acid residue. In pulse radiolysis studies, *in vitro* production of the initial radical on a protein is conveniently made at a tryptophan residue, and electron transfer often leads ultimately to residence of the unpaired electron on a tyrosine residue. We review here the kinetics data for reactions of the antioxidants glutathione, selenocysteine, and ascorbate with tryptophanyl and tyrosyl radicals as free amino acids in model compounds and proteins. Glutathione repairs a tryptophanyl radical in lysozyme with a rate constant of $(1.05 \pm 0.05) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$, while ascorbate repairs tryptophanyl and tyrosyl radicals ca. 3 orders of magnitude faster. The *in vitro* reaction of glutathione with these radicals is too slow to prevent formation of peroxy radicals, which become reduced by glutathione to

hydroperoxides; the resulting glutathione thiyl radical is capable of further radical generation by hydrogen abstraction. Although physiologically not significant, selenogluthione reduces tyrosyl radicals as fast as ascorbate. The reaction of protein radicals formed on insulin, β -lactoglobulin, pepsin, chymotrypsin and bovine serum albumin with ascorbate is relatively rapid, competes with the reaction with dioxygen, and the relatively innocuous ascorbyl radical is formed. On the basis of these kinetics data, we suggest that reductive repair of protein radicals may contribute to the well-documented depletion of ascorbate in living organisms subjected to oxidative stress.

Keywords Protein oxidation · Protein repair · Ascorbate · Glutathione · Selenocysteine · Long range electron transfer

Introduction

Although radical production is part of normal cellular function, excess exposure of macromolecules to oxidising radicals may play a role in ageing and many diseases (Halliwell and Gutteridge 2007). Proteins form by far the largest mass of oxidisable organic components of living matter, and there is a large body of evidence showing that proteins are significant targets of partially reduced oxygen species (PROS) *in vivo*. Increased levels of protein carbonyls are early markers of damage in biological systems subjected to oxidative stress, even in the presence of endogenous antioxidants. The carbonyls are derived from amino acid residues, principally Arg, Pro and Lys, oxidised in reactions with physiological PROS, where the first products are, in some cases, carbon- and oxygen-centred radicals (Requena et al. 2003; von Sonntag 1987). Additional identified products of reaction between PROS and proteins are protein

J. M. Gebicki (✉)
Department of Biological Sciences,
Macquarie University, Sydney 2109, Australia
e-mail: jgebicki@cbms.mq.edu.au

T. Nauser · A. Domazou · D. Steinmann ·
P. L. Bounds · W. H. Koppenol (✉)
Department of Chemistry and Applied Biosciences,
Institute of Inorganic Chemistry, ETH Zürich,
8093 Zurich, Switzerland
e-mail: koppenol@inorg.chem.ethz.ch

When Hasc^- or GSH reacts with a protein radical, hydrogen is abstracted to form the $\text{asc}^{\bullet-}$ or $\text{GS}^{\bullet}$ radical. While $\text{asc}^{\bullet-}$ by all accounts is relatively harmless, $\text{GS}^{\bullet}$ is not, as is indicated by the electrode potentials at pH 7, $E^\circ(\text{asc}^{\bullet-}, \text{H}^+/\text{Hasc}^-) = +0.28 \text{ V}$ (Williams and Yandell 1982) and $E^\circ(\text{GS}^{\bullet}, \text{H}^+/\text{GSH}) = +0.92 \text{ V}$ (Madej and Wardman 2007). Further, the cysteinyl radical can abstract hydrogen from the α -carbon of an amino acid (Nauser and Schöneich 2003).

In this review, we examine the evidence supporting the hypothesis that proteins activated by reactions with PROS may be responsible for much of the depletion of Hasc^- and GSH observed under oxidative stress. We focus on these antioxidants because of their importance in vivo and report largely on our own recent work, because earlier literature on the kinetics of the reactions of protein radicals is limited. It is hoped that this work will provide stimulus for further quantitative studies of biologically relevant reactions of protein radicals.

History

In 1978, it was reported that random attack of $\text{HO}^{\bullet}$ with an amino acid on the surface of cytochrome *c* leads to ca. 50% reduction of Fe^{3+} to Fe^{2+} (van Leeuwen et al. 1978). A tunnelling model was developed to explain the transfer of an electron from the surface to the haem iron buried in the interior of the protein. In this case, electron transfer is from a reducing radical to Fe^{3+} . At the end, the amino acid on the surface loses two electrons, most likely via formation of a double bond, although this was never demonstrated.


In landmark papers published during the 1980s, Prütz and co-workers (Butler et al. 1982; Prütz et al. 1980, 1982) showed that radical attack on a protein leads to a cascade of intramolecular electron transfer reactions that usually results in the formation of a tyrosyl radical. Similar intramolecular electron transfers were studied by Bobrowski and coworkers in the 1990s (Bobrowski et al. 1990, 1992), but only a few studies were dedicated to the repair of protein radicals (Hoey and Butler 1984; Jovanovic and Simic 1985; Santus et al. 2000).

Generation of protein radicals




We use radiolysis and photolysis as tools to generate radicals that are identical ($\text{HO}^{\bullet}$) or similarly oxidising ($\text{N}_3^{\bullet}$) as radicals produced in vivo, e.g. $\text{NO}_2^{\bullet}$ and $\text{CO}_3^{\bullet-}$ from the reaction of ONOO^- with CO_2 (Bonini et al. 1999; Lymar and Hurst 1995; Meli et al. 1999).

Earlier studies of the kinetics of formation of amino acid and protein radicals demonstrated that the short lifetimes of

the radicals require the use of nanosecond laser flash photolysis or pulse radiolysis techniques (von Sonntag 1987). These approaches are capable of generating the radicals commonly produced in organisms subjected to oxidative stress. In flash photolysis, the energy selectively excites the Tyr and Trp residues, converting them to the corresponding Tyr phenoxyl ($\text{TyrO}^{\bullet}$) and Trp indolyl radicals ($\text{Trp}^{\bullet}$), respectively (Bensasson et al. 1983), and, because the absorbance maxima are sufficiently widely separated ($\text{TyrO}^{\bullet}$ at 405 nm and $\text{Trp}^{\bullet}$ at 510 nm), the kinetics of formation and decay of these radicals can be studied by spectrophotometry. In pulse radiolysis of dilute aqueous solutions of amino acids and proteins, virtually all of the absorbed energy of the MeV electrons leads to the decomposition of the water:



When the sample solutions are previously saturated with N_2O , the e_{aq}^- is converted to $\text{HO}^{\bullet}$ (Eq. 2), a powerful oxidant that reacts with virtually all organic compounds. To prevent the generation of too wide a range of protein radical residues, each with its own reactivity, we also include N_3^- to convert $\text{HO}^{\bullet}$ to $\text{N}_3^{\bullet}$ (Eq. 3), which selectively oxidises Trp and Tyr residues (Eqs. 4, 5) (Land and Prütz 1979):



The rate constants of these reactions at near neutral pH are $k_3 = 1.2 \times 10^{10}$, $k_4 \approx 2 \times 10^9$ and $k_5 = 1.0 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ (Butler et al. 1982, 1984; Buxton et al. 1988). These values were used as a guide to choose the concentrations of solutes in the irradiated solutions to ensure completion of Eqs. 3–5 and to exclude competing processes, such as radical–radical reactions.

Reaction of free and lysozyme-bound $\text{TyrO}^{\bullet}$ and $\text{Trp}^{\bullet}$ with glutathione

The absorbance of solutions of 140 μM lysozyme (Lyz), 10 mM phosphate buffer pH 7.4 and 0.1 M sodium azide saturated with N_2O after a 20-ns radiation pulse in the absence and presence of glutathione (GSH) is shown as a function of time in Fig. 1; second-order rate constants for the reactions of GSH are collected in Table 1. The 510 nm absorbance due to the $\text{Lyz}(\text{Trp}^{\bullet})$ develops within 200 μs after the pulse, while the subsequent decay of the radical is slower. In contrast, the $\text{Lyz}(\text{TyrO}^{\bullet})$ (405 nm peak)




Fig. 1 Time-dependent absorbance changes in pulse-irradiated lysozyme (LZ). The peaks at 510 and 410 nm are due to the $\text{Lyz}(\text{Trp}^\bullet)$ and $\text{Lyz}(\text{TyrO}^\bullet)$ radicals, respectively (reproduced from Nauser et al. 2005 with permission of the authors and the Biochemical Society)

generated by intramolecular long-range electron transfer (LRET) to $\text{Lyz}(\text{Trp}^\bullet)$ forms more slowly, as reported in earlier studies (Butler et al. 1982). Irradiation in the presence of 2.5 mM GSH leads to complete loss of the absorbance at 510 nm due to $\text{Trp}^\bullet$ within 2 ms, which indicates that all $\text{Lyz}(\text{Trp}^\bullet)$ radicals are effectively repaired by H transfer. Under conditions of excess GSH, the reaction follows pseudo-first-order kinetics, and the first-order rate constant (k_{obs}) is linearly dependent on the concentration of GSH, which allows the second-order rate constant, $k_6 = (1.05 \pm 0.05) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ for the bimolecular reaction to be derived



thereby preventing formation of $\text{Lyz}(\text{TyrO}^\bullet)$ by intramolecular LRET:



which proceeds with a first-order rate constant $k_7 = (5.3 \pm 1.3) \times 10^2 \text{ s}^{-1}$. Measurements of the kinetics of the reaction between GSH and $\text{Lyz}(\text{Trp}^\bullet)$ show that the reaction is complete within 1.5 ms. GSH is also effective in vitro at accelerating the decay of the $\text{Lyz}(\text{Trp}^\bullet)$ generated by flash photolysis of Lyz (28 μM in 10 mM phosphate buffer at pH 7.4 in the presence of 100 mM *t*-butanol). Table 1 also lists the rate constants for reaction of GSH with the protein amino acid analogues *N*-Ac-Trp amide and *N*-Ac-TyrOH amide at pH 7.4 measured in similar experiments.

The value of the rate constant for LRET in Lyz agrees well with an earlier published rate constant (Bobrowski et al. 1997). It should be noted that, although the reactions of $\text{Lyz}(\text{Trp}^\bullet)$ radicals with GSH are only moderately fast, the result indicates that reaction with oxidised protein

Table 1 Rate constants for reactions of GSH, Sec and GSeH with proteins and protein amino acid analogues

Reaction	Rate constant ($\text{M}^{-1} \text{ s}^{-1}$)	Reference
$\text{GSH} + \text{Lyz}(\text{Trp}^\bullet)$	$(1.05 \pm 0.05) \times 10^5$	Nauser et al. (2005)
$\text{GSH} + \text{N-Ac-Trp}^\bullet$ amide	$(1.1 \pm 0.3) \times 10^5$	Nauser et al. (2005)
$\text{GSH} + \text{N-Ac-TyrO}^\bullet$ amide	$(2.4 \pm 1.6) \times 10^2$	Nauser et al. (2005)
$\text{Sec} + \text{N-Ac-TyrO}^\bullet$	$(8 \pm 2) \times 10^8$	Steinmann et al. (2008)
$\text{Sec} + \text{Ins}(\text{TyrO}^\bullet)$	$(1.6 \pm 0.4) \times 10^8$	Steinmann et al. (2008)
$\text{GSeH} + \text{N-Ac-TyrO}^\bullet$	$(5 \pm 2) \times 10^8$	Steinmann et al. (2008)

could account for much or for at least some of the documented loss of GSH under conditions of oxidative stress in vivo. Further, when residues other than Trp and Tyr become oxidised to form C-centred radicals, these radicals subsequently form peroxy radicals, which greatly enhances the amount of GSH oxidised; under air, enhancement of GSH oxidation can be as high as 20-fold, and we have suggested that the peroxy radicals and hydroperoxides generated can constitute a new source of biological damage (Nauser et al. 2005).


Repair of tyrosyl radicals by selenocysteine and selenogluthione

We have also found that selenocysteine (Sec) repairs $\text{TyrO}^\bullet$ radicals (Steinmann et al. 2008). Sec reduces *N*-Ac-TyrO[•]-amide with a rate constant of $(8 \pm 2) \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ and $\text{TyrO}^\bullet$ in insulin (Ins) with $(1.6 \pm 0.4) \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ (Table 1). The rate constant for the reduction of *N*-Ac-TyrO[•]-amide by selenogluthione (GSeH) is slightly lower $(5 \pm 2) \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ than that for Sec. In contrast, Cys and GSH react 3 and 5 orders of magnitude, respectively, more slowly with *N*-Ac-TyrO[•]-amide than Sec and GSeH; Cys and GSH react 3 and 2 orders of magnitude, respectively, more slowly with $\text{TyrO}^\bullet$ in Ins than their selenium analogues.



Reactions of free and protein-bound $\text{TyrO}^\bullet$ and $\text{Trp}^\bullet$ radicals with ascorbate

The kinetics for repair of the $\text{Trp}^\bullet$ and $\text{TyrO}^\bullet$ radicals by ascorbate (Hasc^-) were studied by pulse radiolysis with the *N*-acetylated amino acid amides and with the proteins Pep, Ins, chymotrypsin (Chym) and β -lactoglobulin (β -Lact) (Domazou et al. 2009). Radicals were generated by reaction with N_3^- ; repair of $\text{Trp}^\bullet$ was followed at 510 and of $\text{TyrO}^\bullet$ at 405 nm. The conditions were chosen to ensure

Fig. 2 The spontaneous decay (red) of **a** Pep(Trp[•]) and **b** Pep(TyrO[•]). The radicals were generated by pulse radiolysis of solutions that contained pepsin and sodium azide, and were saturated with N₂O. The blue curves show the accelerated decay of Pep(Trp[•]) and Pep(TyrO[•]) in the presence of 0.2 mM Hasc⁻



conversion of e_{aq}^- to $HO^\bullet$ (Eq. 2) and complete scavenging of $HO^\bullet$ by N_3^- (Eq. 3) to guarantee formation of stoichiometric amounts of $N_3^\bullet$. Furthermore, low radiation doses were used to limit radical–radical recombination, and the amino acid or protein was present in excess to prevent direct reaction of $N_3^\bullet$ with Hasc⁻. In all cases, spontaneous rates of decay of radicals were accelerated in the presence of Hasc⁻. The overall result is the formation of the relatively inert ascorbyl radical and repair of the amino acid. The results for pepsin, as summarised in Eqs. 8 and 9, are representative of all results with proteins and are shown in Fig. 2.



The collected rate constants for Eqs. 8 and 9 and for the corresponding reactions in proteins other than pepsin are listed in Table 2. The rate constants for reaction of Hasc⁻ with *N*-Ac-TyrO[•] amide are comparable to those for the reduction of TyrO[•] by Sec and GSeH. It can be seen that, in general, Hasc⁻ reacts 10–50 times faster with protein-bound Trp[•] than with protein-bound TyrO[•] radicals. Interestingly, the protein environment apparently does not affect the kinetics of Eq. 8, but lowers the rate constant for protein-bound TyrO[•] (Eq. 9) by about a factor of 10, compared to free *N*-Ac-Trp[•] amide. Presumably, these effects depend on variables such as the nature of the protein, the location of the reacting residue and potential stabilisation of the radical by neighbouring amino acids. Preliminary evidence suggests that peroxy radicals of *N*-Ac-Pro-amide, *N*-Ac-Ala-amide and *N*-Ac-Gly-Ala-amide react with Hasc⁻ with rate constants of ca. $10^7 \text{ M}^{-1} \text{ s}^{-1}$ (Zelenay and Domazou 2009, unpublished).

Potential biological significance of radical oxidation of proteins and long-range electron transfer

We have presented evidence that the antioxidants GSH and Hasc⁻ react with TyrO[•] and Trp[•] as free amino acids, protein residue analogues and protein radicals. Both

Table 2 Rate constants of reactions ascorbate with Trp[•] and TyrO[•] as free amino acids, peptide models and proteins

Amino acid or protein radical	Rate constant	Reference
TyrO [•]	9.3×10^7	Hoey and Butler (1984)
<i>N</i> -Ac-TyrO [•] amide	$(2.6 \pm 0.2) \times 10^8$	Domazou et al. (2009)
Trp [•]	1×10^8	Jovanovic and Simic (1985)
<i>N</i> -Ac-Trp [•] amide	$(1.4 \pm 0.1) \times 10^8$	Domazou et al. (2009)
Ins(TyrO [•])	$(2.9 \pm 0.1) \times 10^7$	Domazou et al. (2009)
Chym(TyrO [•])	4×10^7	Domazou et al. (2009)
Chym(Trp [•])	$(1.6 \pm 0.1) \times 10^8$	Domazou et al. (2009)
Pep(TyrO [•])	$(3.5 \pm 0.9) \times 10^7$	Domazou et al. (2009)
Pep(Trp [•])	$(1.8 \pm 0.1) \times 10^8$	Domazou et al. (2009)
β -Lact(TyrO [•])	$(4.0 \pm 0.6) \times 10^5$	Domazou et al. (2009)
β -Lact(Trp [•])	$(2.2 \pm 0.2) \times 10^7$	Domazou et al. (2009)
Lyz(TyrO [•])	1.1×10^7	Hoey and Butler (1984)
Lyz(Trp [•])	8×10^7	Santus et al. (2000)
Lyz(Trp [•])	8.3×10^7	Hoey and Butler (1984)

antioxidants inhibit the transfer of electrons from Tyr to Trp[•] in proteins in vitro, but GSH does not prevent the formation of protein peroxy radicals or hydroperoxides. The repair of protein C-centred radicals by Hasc⁻ competes with the reaction with O₂. Both reactions may contribute to the loss of GSH and Hasc⁻ in living organisms under conditions of oxidative stress. It should be noted that, while the radicals formed in a protein can be efficiently repaired by GSH and Hasc⁻, repair does not necessarily restore biological function, which may be irreversibly altered by the initial radical reaction and/or subsequent formation of other radical species by LRET.

Certain cellular signalling pathways, e.g. the signal transducers and activators of transcription (STAT) family of transcription factors, function via reversible phosphorylation of tyrosine residues by tyrosine kinases and phosphatases (Ihle 2001), which are modulated by the redox status of the cell (Monteiro and Stern 1996), likely due to the impact of LRET on ultimate formation of TyrO[•] radicals in proteins.

Finally, radical reactions in proteins are not uniformly harmful. Numerous enzyme reaction mechanisms rely upon transient formation of an organic radical, e.g. the relatively stable tyrosyl radical formed during the reaction of ribonuclease reductase I (Stubbe and van der Donk 1998). Iodination of tyrosine residues on thyroglobulin during thyroid hormone biosynthesis synthesis also involves transient tyrosyl radical formation (Nunez 1984). Phosphorylation of free tyrosine has been shown to block tyrosine iodination (Gmeiner and Seelos 1996) and dimerisation, both of which are radical reactions.

Acknowledgments Supported by the Eidgenössische Technische Hochschule, Zürich, Switzerland, the Swiss Nationalfonds, Bern, Switzerland, and Macquarie University, Sydney, Australia.

References

- Bensasson RV, Land EJ, Truscott TG (1983) Flash photolysis and pulse radiolysis. Pergamon, Oxford
- Bisby RH, Parker AW (1995) Reaction of ascorbate with the α -tocopheroxyl radical in micellar and bilayer membrane systems. *Arch Biochem Biophys* 317:170–178
- Bobrowski K, Wierchowski KL, Holcman J, Ciurak M (1990) Intramolecular electron transfer in model peptides containing methionine, tryptophan and tyrosine: a pulse radiolysis study. *Int J Radiat Biol* 57:919–932
- Bobrowski K, Wierchowski KL, Holcman J, Ciurak M (1992) Pulse radiolysis studies of intramolecular electron transfer in model peptides and proteins. IV. Met/S-Br $\rightarrow$ tyr/O radical transformation in aqueous solution of H-Tyr-(Pro)n-Met-OH peptides. *Int J Radiat Biol* 62:507–516
- Bobrowski K, Holcman J, Poznanski J, Wierchowski KL (1997) Pulse radiolysis studies of intramolecular electron transfer in model peptides and proteins. 7. Trp $\rightarrow$ TyrO radical transformation in hen egg-white lysozyme. Effects of pH, temperature, Trp62 oxidation and inhibitor binding. *Biophys Chem* 63:153–166
- Bonini MG, Radi R, Ferrer-Sueta G, Ferreira AMD, Augusto O (1999) Direct EPR detection of the carbonate radical anion produced from peroxynitrite and carbon dioxide. *J Biol Chem* 274:10802–10806
- Buettner GR (1993) The pecking order of free radicals and antioxidants. Lipid peroxidation, α -tocopherol, and ascorbate. *Arch Biochem Biophys* 300:535–543
- Buffington GD, Doe WF (1995) Altered ascorbic acid status in the mucosa from inflammatory bowel disease patients. *Free Radic Res* 22:131–143
- Burton GW, Wronska U, Stone L, Foster DO, Ingold KU (1990) Biokinetics of dietary *RRR*- α -tocopherol in the male guinea pig at three dietary levels of vitamin C and two levels of vitamin E. Evidence that vitamin C does not “spare” vitamin E in vivo. *Lipids* 25:199–210
- Butler J, Land EJ, Prütz WA, Swallow AJ (1982) Charge transfer between tryptophan and tyrosine in proteins. *Biochim Biophys Acta* 705:150–162
- Butler J, Land EJ, Swallow AJ, Prütz WA (1984) The azide radical and its reaction with tryptophan and tyrosine. *Radiat Phys Chem* 23:265–270
- Buxton GV, Greenstock CL, Helman WP, Ross AB (1988) Critical review of rate constants for reactions of hydrated electrons. Hydrogen atoms and hydroxyl radicals ($^{\bullet}\text{OH}/^{\bullet}\text{O}^-$) in aqueous solution. *J Phys Chem Ref Data* 17:513–886
- Davies MJ (2005) The oxidative environment and protein damage. *Biochim Biophys Acta* 1703:93–109
- Davies MJ, Dean RT (1997) Radical-mediated protein oxidation: from chemistry to medicine. Oxford University Press, New York
- Domazou AS, Koppenol WH, Gebicki JM (2009) Efficient repair of protein radicals by ascorbate. *Free Radic Biol Med* 46:1049–1057
- Drechsel DA, Patel M (2008) Role of reactive oxygen species in the neurotoxicity of environmental agents implicated in Parkinson's disease. *Free Radic Biol Med* 44:1873–1886
- Flohé L, Günzler WA, Schock HH (1973) Glutathione peroxidase: a selenoenzyme. *FEBS Lett* 32:132–134
- Foy CJ, Passmore AP, Vahidassr MD, Young IS, Lawson JT (1999) Plasma chain-breaking antioxidants in Alzheimer's disease, vascular dementia and Parkinson's disease. *Q J Med* 92:39–45
- Garrison WM (1987) Reaction mechanisms in the radiolysis of peptides, polypeptides, and proteins. *Chem Rev* 87:381–398
- Gebicki JM (1997) Protein hydroperoxides as new reactive oxygen species. *Redox Rep* 3:99–110
- Gmeiner BMK, Seelos CCC (1996) Tyrosine phosphorylation blocks tyrosine free radical formation and, hence, the hormonogenic iodination reaction. *Free Radic Biol Med* 21:349–351
- Halliwell B, Gutteridge JMC (2007) Free radicals in biology and medicine, 4th edn. Oxford University Press, Oxford
- Hoey BM, Butler J (1984) The repair of oxidized amino acids by antioxidants. *Biochim Biophys Acta* 791:212–218
- Ihle JN (2001) The Stat family in cytokine signaling. *Curr Opin Cell Biol* 13:211–217
- Jovanovic SV, Simic MG (1985) Repair of tryptophanyl radicals by antioxidants. *J Free Radic Biol Med* 1:125–129
- Kume-Kick J, Rice ME (1998) Estrogen-dependent modulation of rat brain ascorbate levels and ischemia-induced ascorbate loss. *Brain Res* 803:105–113
- Land EJ, Prütz WA (1979) Reaction of azide radicals with amino acids and proteins. *Int J Radiat Biol* 36:75–83
- Liu R-M, Pavia KAG (2010) Oxidative stress and glutathione in TGF- β -mediated fibrogenesis. *Free Radic Biol Med* 48:1–15
- Locatelli F, Canaud B, Eckardt K-U, Stenvinkel P, Wanner C, Zoccali C (2003) Oxidative stress in end-stage renal disease: an emerging threat to patient outcome. *Nephrol Dial Transplant* 18:1272–1280
- Lymar SV, Hurst JK (1995) Rapid reaction between peroxynitrite ion and carbon dioxide: implications for biological activity. *J Am Chem Soc* 117:8867–8868
- Madej E, Wardman P (2007) The oxidizing power of the glutathione thiyl radical as measured by its electrode potential at physiological pH. *Arch Biochem Biophys* 462:94–102
- Meli R, Nauser T, Koppenol WH (1999) Direct observation of intermediates in the reaction of peroxynitrite with carbon dioxide. *Helv Chim Acta* 82:722–725
- Michiels C, Raes M, Toussaint O, Remacle J (1994) Importance of Se-glutathione peroxidase, catalase and Cu/Zn-SOD for cell survival against oxidative stress. *Free Radic Biol Med* 17:235–248
- Misso NLA, Brooks-Wildhaber J, Ray S, Vally H, Thompson PJ (2005) Plasma concentrations of dietary and nondietary antioxidants are low in severe asthma. *Eur Respir J* 26:257–264
- Monteiro HP, Stern A (1996) Redox modulation of tyrosine phosphorylation-dependent signal transduction pathways. *Free Radic Biol Med* 21:323–333
- Moor E, Shohami E, Kanevsky E, Grigoriadis N, Symeonidou C, Kohen R (2006) Impairment of the ability of the injured aged brain in elevating urate and ascorbate. *Exp Gerontol* 41:303–311
- Nauser T, Schöneich C (2003) Thiyl radicals abstract hydrogen atoms from the $^2\text{C-H}$ bonds in model peptides: absolute rate constants

- and effect of amino acid structure. *J Am Chem Soc* 125:2042–2043
- Nauser T, Koppenol WH, Gebicki JM (2005) The mechanism and kinetics of the oxidation of GSH by protein free radicals. *Biochem J* 392:693–701
- Neta P, Huie RE, Ross AB (1990) Rate constants for reactions of peroxy radicals in fluid solutions. *J Phys Chem Ref Data* 19:413–513
- Nunez J (1984) Thyroid hormones: mechanism of phenoxy ether formation. *Methods Enzymol* 107:476–488
- Pietzsch J (ed) (2006) Protein oxidation and disease. In: Protein oxidation and disease. Research Signpost, Trivandrum, pp 1–6
- Prütz WA, Butler J, Land EJ, Swallow AJ (1980) Direct demonstration of electron transfer between tryptophan and tyrosine in proteins. *Biochem Biophys Res Commun* 96:408–414
- Prütz WA, Siebert F, Butler J, Land EJ, Menez A, Montenay-Garestier T (1982) Intramolecular radical transformations involving methionine, tryptophan and tyrosine. *Biochim Biophys Acta* 705:139–149
- Pryor WA (1986) Cancer and free radicals. In: Shankel DM, Hollaender A, Hartman PE, Kada T (eds) Antimutagenesis and anticarcinogenesis mechanisms. Basic Books, New York, pp 45–59
- Requena JR, Levine RL, Stadtman ER (2003) Recent advances in the analysis of oxidized proteins. *Amino Acids* 25:221–226
- Rubbo H, Radi R, Anselmi D, Kirk M, Barnes S, Butler J, Eiserich JP, Freeman BA (2000) Nitric oxide reaction with lipid peroxy radicals spares α -tocopherol during lipid peroxidation—greater oxidant protection from the pair nitric oxide/ α -tocopherol than α -tocopherol/ascorbate. *J Biol Chem* 275:10812–10818
- Rush JD, Koppenol WH (1990) Reactions of Fe(II)-ATP and Fe(II)-citrate complexes with *t*-butyl hydroperoxide and cumylhydroperoxide. *FEBS Lett* 275:114–116
- Santus R, Patterson LK, Hug GL, Bazin M, Mazière JC, Morlière P (2000) Interactions of superoxide anion with enzyme radicals: kinetics of reaction with lysozyme tryptophan radicals and corresponding effects on tyrosine electron transfer. *Free Radic Res* 33:383–391
- Simpson JA, Narita S, Gieseg S, Gebicki S, Gebicki JM, Dean RT (1992) Long-lived reactive species on free-radical-damaged proteins. *Biochem J* 282:621–624
- Steinmann D, Nauser T, Beld J, Tanner M, Günther D, Bounds PL, Koppenol WH (2008) Kinetics of tyrosyl radical reduction by selenocysteine. *Biochemistry* 47:9602–9607
- Stubbe J, van der Donk WA (1998) Protein radicals in enzyme catalysis. *Chem Rev* 98:705–762
- van Leeuwen JW, Raap A, Koppenol WH, Nauta H (1978) A tunneling model to explain the reduction of ferricytochrome *c* by H and OH radicals. *Biochim Biophys Acta* 503:1–9
- von Sonntag C (1987) The chemical basis of radiation biology. Taylor & Francis, London
- Williams MH, Yandell JK (1982) Outer-sphere electron transfer reactions of ascorbate anions. *Aust J Chem* 35:1133–1144
- Winterbourn CC (2008) Reconciling the chemistry and biology of reactive oxygen species. *Nat Chem Biol* 4:278–286