

(0.170 g, 52 % yield). Elemental analysis calcd (%) for $C_{28}H_{44}O_{14}NKTi$: C 47.66, H 6.28, N 1.99; found: C 47.43, H 6.27, N 2.06; m.p. 124–125 °C (decomp); IR (THF): $\bar{\nu}$ = 1930 (m), 1793 cm^{-1} (s) (CO); 1H NMR (300 MHz, $[D_8]THF$, 0 °C, TMS): δ = 6.16 (m, 2H, pyrrolyl), 5.40 (m, 2H, pyrrolyl), 3.62 (s, 40H, [15]crown-5); $^{13}C\{^1H\}$ (75 MHz, $[D_8]THF$, 0 °C, TMS): δ = 289 (s, CO), 114 (s, pyrrolyl), 99 (s, pyrrolyl), 69 (s, [15]crown-5).

Received: July 30, 1999 [Z13810]

- [1] a) T. J. Marks, J. R. Kolb, *Chem. Rev.* **1977**, 77, 263; b) B. D. James, M. G. H. Wallbridge, *Prog. Inorg. Chem.* **1970**, 11, 99.
- [2] S. W. Kirtley, M. A. Andres, R. Bau, G. W. Grynkeiwich, T. J. Marks, D. L. Tipton, B. R. Whittlesey, *J. Am. Chem. Soc.* **1977**, 99, 7154.
- [3] a) M. Y. Darensbourg, R. Bau, M. W. Marks, R. R. Burch, Jr., J. C. Deaton, S. Slater, *J. Am. Chem. Soc.* **1982**, 104, 6961; b) K.-M. Chi, S. R. Frerichs, S. B. Philson, J. E. Ellis, *J. Am. Chem. Soc.* **1988**, 110, 303.
- [4] a) Previously it was shown that the dinuclear hydrido complex $[HMo_2(CO)_{10}]^-$ in the presence of I^- reacted with equimolar amounts of $BH_3 \cdot THF$ to give $[Mo(CO)_4(\eta^2-BH_4)]^-$ and $[Mo(CO)_5]^-$; [2] b) G. W. Parshall, *J. Am. Chem. Soc.* **1964**, 86, 361.
- [5] a) G. Medford, S. G. Shore, *J. Am. Chem. Soc.* **1978**, 100, 3953; b) T. J. Coffy, G. Medford, J. Plotkin, G. J. Long, J. C. Huffman, S. G. Shore, *Organometallics* **1989**, 8, 2404; c) T. J. Coffy, S. G. Shore, *J. Organomet. Chem.* **1990**, 394, C27.
- [6] In a related reaction, $[Fe(CO)_4]^{2-}$ was shown to react with $BH_3 \cdot THF$ to give equimolar amounts of $[Fe(CO)_4(\eta^2-B_2H_5)]^-$ and $K[BH_4]$. [5b] Although anionic $M-BH_3$ complexes [4b] are likely intermediates in both this reaction and that involving the titanate, in neither case were any intermediates observed. The two reactions provide quite a contrast, however, since in the case of Fe free $[BH_4]^-$ is formed, while the Ti-mediated reaction affords coordinated $[BH_4]^-$.
- [7] a) B. A. Kelsey, J. E. Ellis, *J. Am. Chem. Soc.* **1986**, 108, 5016; b) G. Binsch in *Dynamic Nuclear Magnetic Resonance Spectroscopy* (Eds.: L. M. Jackman, F. A. Cotton), Academic Press, New York, **1975**, chap. 3.
- [8] a) Crystal data for $[K([18]crown-6)]_2$: $C_{42}H_{60}K_2N_{12}O_{22}Ti_2$, M_r = 1118.92, monoclinic, space group $P2_1/c$, a = 15.496(3), b = 15.374(3), c = 11.245(2) Å, β = 91.02(3)°, V = 2678.4(0) Å³, Z = 2, ρ_{calcd} = 1.387 g cm⁻³, $F(000)$ = 1168, $\mu(Mo_{K\alpha})$ = 0.530 mm⁻¹, λ = 0.71073 Å, T = 173(2) K; 9448 reflections were measured on a Siemens SMART CCD area detector diffractometer and yielded 4499 unique data ($2\theta_{max}$ = 50°, R_{int} = 0.0309, semi-empirical absorption corrections). The structure was solved by direct methods and refined by full-matrix least-squares methods on F^2 with statistical weighting, anisotropic displacement parameters, and constrained isotropic H atoms to give $wR2 = \{\sum[w(F_o^2 - F_c^2)]^2 / \sum[w(F_o^2)]^2\}^{1/2} = 0.1411$ on all data, conventional $R = 0.0611$ on F values of 4424 reflections having $F_o^2 > 2\sigma(F_o^2)$, goodness of fit $S = 0.983$ for all F^2 values and 558 refined parameters. Largest difference peak/hole 0.345/−0.330 e Å⁻³. All calculations were performed on SGI INDY R4400-SC or Pentium computers with the SHEXTL-Plus program suite; b) Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-132492. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [9] a) H. Nöth, M. Schmidt, *Organometallics* **1995**, 14, 4601; b) F. Corazza, C. Floriani, A. Chiesi-Villa, C. Guastini, *Inorg. Chem.* **1991**, 30, 145.
- [10] a) M. Mancini, P. Bougeard, R. C. Burns, M. Mlekuz, B. G. Sayer, J. I. A. Thompson, M. J. McGlinchey, *Inorg. Chem.* **1984**, 23, 1072; b) Y. Oishi, T. A. Albright, H. Fujimoto, *Polyhedron* **1995**, 14, 2603. This paper discusses the electronic properties of the unknown $[Mn(CO)_4(BH_4)]$. It is noteworthy that $[Mn(CO)_3(\eta^3-BH_4)]$, which would be analogous to $[Mn(CO)_3(\eta^5-C_5H_5)]$, is also unknown.
- [11] J. K. Seaburg, P. J. Fischer, V. G. Young, Jr., J. E. Ellis, *Angew. Chem.* **1998**, 110, 163; *Angew. Chem. Int. Ed.* **1998**, 37, 155.
- [12] a) P. J. Fischer, K. A. Ahrendt, V. G. Young, Jr., J. E. Ellis, *Organometallics* **1998**, 17, 13; b) K.-M. Chi, S. R. Frerichs, B. K. Stein, D. W. Blackburn, J. E. Ellis, *J. Am. Chem. Soc.* **1988**, 110, 163; $HBPz_3$ = hydridotris(1-pyrazolyl)borate.
- [13] J. E. Ellis, A. J. DiMaio, A. L. Rheingold, B. S. Haggerty, *J. Am. Chem. Soc.* **1992**, 114, 10676; $tacn$ = 1,4,7-triazacyclononane; Me_3tacn = 1,4,7-trimethyl-1,4,7-triazacyclononane.
- [14] a) M. L. Luetkens, J. C. Huffman, A. P. Sattelberger, *J. Am. Chem. Soc.* **1985**, 107, 3361; b) J. E. Gozum, G. S. Girolami, *J. Am. Chem. Soc.* **1991**, 113, 3829, and references therein.
- [15] Details of the structure and structural characterization will be described elsewhere: P. J. Fischer, V. G. Young, Jr., J. E. Ellis, unpublished results. Crystal data for $[(Ph_3P)_2N]^-3$: $C_{44}H_{34}N_2O_4P_2Ti$, M_r = 764.57, monoclinic, space group $P2_1$, red plate, a = 10.6547(2), b = 12.6892(3), c = 13.9628(4) Å, β = 98.483(1)°, Z = 2, R = 0.0502, with GOF on F^2 of 1.066.
- [16] a) M. Rakowski Du Bois, K. G. Parker, C. Ohman, B. C. Noll, *Organometallics* **1997**, 16, 2325, and references therein; b) see also ref. [8] of J. R. Perera, M. J. Heeg, H. B. Schlegel, C. H. Winter, *J. Am. Chem. Soc.* **1999**, 121, 4536.
- [17] R. V. Bynum, W. E. Hunter, R. D. Rogers, J. L. Atwood, *Inorg. Chem.* **1980**, 19, 2368.
- [18] a) K. K. Joshi, P. L. Pauson, A. R. Qazi, W. H. Stubbs, *J. Organomet. Chem.* **1964**, 1, 471; b) R. B. King, A. Efraty, *J. Organomet. Chem.* **1969**, 20, 264.

A Metamagnetic Two-Dimensional Molecular Material with Nickel(II) and Azide**

Montserrat Monfort,* Immaculada Resino, Joan Ribas, and Helen Stoeckli-Evans

Until a few years ago, the main aim of magnetochemical research was to show that molecular magnetism could lead to the development of new magnetic materials.^[1] One strategy for the design of molecular-based magnets involves assembling transition metal ions in one-, two-, or three-dimensional networks. We are currently studying materials formed from the interaction of transition metal ions with the azido ligand. The azido bridge is a versatile and efficient magnetic coupler; it may transmit either antiferro- or ferromagnetic interactions. Working with Ni^{II} salts and diamines as blocking ligands, for a Ni/diamine ratio of 1/1 four positions in the coordination sphere of the nickel are free. These positions can be occupied by four azido bridges. The most remarkable species obtained to date are the neutral polymeric compounds $[Ni(\mu-N_3)_2(\text{diamine})]_n$: three one-dimensional systems showing global ferromagnetic behavior and metamagnetism at low temperature,^[2] five one-dimensional complexes with alternating end-on (EO) and end-to-end (EE) azido bridges in different sequences showing global anti-ferromagnetic behavior,^[3–6]

[*] Prof. M. Monfort, Dra. I. Resino, Prof. J. Ribas
Departament de Química Inorgànica
Universitat de Barcelona, Diagonal 647, 08028 Barcelona (Spain)
Fax: (+34) 93-490-77-25
E-mail: mmonfort@kripto.ubi.es
Prof. H. Stoeckli-Evans
Institut de Chimie, Université de Neuchâtel
Av. de Bellevaux 51, 2000 Neuchâtel (Switzerland)

[**] This work was supported by the Spanish Dirección General de Investigación Científica y Técnica (project PB96-0163).

and one two-dimensional sheet showing global antiferromagnetic behavior and a canting phenomenon at low temperature (50 K) that gives the complex the characteristics of a molecular magnet.^[7] Here, we report the synthesis, structure, and magnetic properties of $[\text{Ni}(\mu\text{-N}_3)_2(\text{N,N}\text{-Et}_2\text{-N'}\text{-Me-en})]_n$ (**1**; $\text{N,N}\text{-Et}_2\text{-N'}\text{-Me-en} = \text{N,N}$ -diethyl- N' -methylethylenediamine), which is the first example of a metamagnetic two-dimensional molecular material with Ni^{II} and azide, featuring short-range, intralayer, ferromagnetic behavior and weak interlayer antiferromagnetic interactions. The crystal structure analysis^[8] reveals parallel sheets of nickel atoms linked by EO and EE azido bridges (Figure 1). Each nickel atom is in a




Figure 1. Layer structure of **1**. Dinuclear nickel units, bridged by two EO azido bridges, are linked to four equivalent units by means of four single EE azido bridges. For clarity, only the N atoms of the amine ligands are shown.

distorted octahedral NiN_6 environment and is linked to three other nickel atoms: through two EO azido bridges to one nickel atom and through two EE bridges in a *cis* position to two nickel atoms. Each Ni^{II} ion completes its coordination by binding one bidentate $\text{N,N}\text{-Et}_2\text{-N'}\text{-Me-en}$ ligand. The Ni–N bond lengths range between 2.09 and 2.15 Å, which is characteristic of these complexes. The Ni–N–Ni angle of 100.46° lies within the range of reported values.^[2, 4, 5] All the EE fragments are equivalent and the coordination of the azido bridge is asymmetric; the two Ni–N–N angles are 136.9° and 144.2° , and the Ni– N_3 –Ni torsion angle is 110.44° .

The cryomagnetic behavior of **1** is shown in Figure 2. At room temperature $\chi_{\text{M}}T$ is the same ($1.44 \text{ cm}^3 \text{ K mol}^{-1}$) for all values of the applied field. This value increases monotonically with cooling to 60 K. As the temperature is lowered further $\chi_{\text{M}}T$ increases more and more rapidly, reaching a maximum value around 16 K, before decreasing at lower temperatures. The maximum value of $\chi_{\text{M}}T$ is field dependent, but the temperature for the maximum $\chi_{\text{M}}T$ remains unchanged. The magnetic behavior, except for the decreasing $\chi_{\text{M}}T$ below 16 K, suggests an intramolecular ferromagnetic interaction between adjacent Ni^{II} ions. These results can be explained by considering the structure. The interaction between the nickel atoms




Figure 2. Plot of $\chi_{\text{M}}T$ versus temperature at three values of the applied field ($\square$: 500 G, $\circ$: 1000 G, $\triangle$: 2000 G). Inset: Plot of χ_{M} versus temperature.

through the EO azido bridges, according to the structural parameters and literature data,^[9, 10] must be ferromagnetic. For the interaction through the EE azido ligand, previous studies^[11, 12] show that the value of the superexchange coupling depends mainly on the Ni–N–N angle α and on the Ni– N_3 –Ni torsion angle θ . Considering the Ni–N–N angle, the maximum antiferromagnetic coupling would be expected for $\alpha = 108^\circ$. For greater values, the interaction must decrease and reach accidental orthogonality at $\alpha = 165^\circ$. Taking into account the torsion angle, the maximum superexchange is found for $\theta = 180^\circ$ or 0° and diminishes gradually to an expected minimum of zero for a torsion angle of $\theta = 90^\circ$. To our knowledge all compounds with an EE azido bridge and a Ni^{II} ion are antiferromagnetic with the exception of two compounds: one is a dinuclear nickel(II) cryptate,^[13] in which the Ni–N–N angles are 165.8° and 157.6° ; the other is a one-dimensional compound,^[14] in which the torsion angle is 75.7° . For **1** the values of the Ni–N–N angles deviate from 108° , and the torsion angle is close to 90° , in the range where the antiferromagnetic contribution is minimized to favor the ferromagnetic contribution.

The magnetic behavior of **1**, with the abrupt increase in the value of $\chi_{\text{M}}T$ on cooling, a maximum around 16 K, and decreasing $\chi_{\text{M}}T$ below this temperature, suggests a three-dimensional antiferromagnetic ordering of the ferromagnetic layers.^[15] Further support for this comes from magnetization versus applied field plots at three temperatures (Figure 3). All curves have the sigmoidal shape (more markedly at lower temperatures) expected for a metamagnet: initially the magnetization increases slightly with increasing field strength because of antiferromagnetic interlayer interactions, subsequently it increases abruptly for a phase transition to a ferromagnetic state, and finally it reaches a saturation magnetization plateau.

The magnetization was also measured in the range of 3–20 K (Figure 4). The field-cooled magnetization (FCM), measured under an applied field of 10 G, increases rapidly on cooling to 16 K and then diminishes. This behavior is typical of metamagnetic ordering. When the applied field was switched off at 3 K, there remained a remnant magnetization (RM), that decreased upon warming. The zero-field-cooled magnetization (ZFCM), measured upon warming under an




Figure 3. Magnetization versus applied field at three temperatures.




Figure 4. Plots of FCM (○), ZFCM (◻), and RM (Δ) versus temperature at an applied field of 10 G.

applied field of 10 G, shows a maximum at 16 K and decreases rapidly as the temperature is increased further.

The present example is a new demonstration of the versatility of the azido bridge for building molecular-based materials. The magnetic measurement shows, for the first time in a Ni^{II} azido compound, an intralayer ferromagnetic interaction. Finally, the interlayer antiferromagnetic interaction gives rise to metamagnetic behavior. In order to complete the magnetic studies, measurements at alternating current, and of the FCM at different field strengths are now in progress.

Experimental Section

A solution of *N,N*-Et₂-*N'*-Me-en (0.26 g, 2 mmol) in DMF/EtOH (1/1, 10 mL) was added to a solution of Ni(NO₃)₂·6H₂O (0.58 g, 2 mmol) in DMF/EtOH (20 mL). Solid NaN₃ (0.26 g, 2 mmol) was added slowly. The clear dark green solution was filtered to remove any impurities and left to stand undisturbed at room temperature. Single green crystals of **1** were collected after two weeks.

- [1] *Molecular-Based Magnetic Materials: Theory, Techniques and Applications* (Eds.: M. M. Turnbull, T. Sugimoto, L. K. Thompson) ACS Symp. Ser. **1996**, 644.
- [2] J. Ribas, M. Monfort, C. Diaz, C. Bastos, X. Solans, *Inorg. Chem.* **1994**, 33, 484–489.
- [3] J. Ribas, M. Monfort, B. Kumar-Gosh, X. Solans, M. Font-Bardía, *J. Chem. Soc. Chem. Commun.* **1995**, 2375–2376.
- [4] J. Ribas, M. Monfort, I. Resino, X. Solans, P. Rabu, F. Maingot, M. Drillon, *Angew. Chem.* **1996**, 108, 2671–2673; *Angew. Chem. Int. Ed. Engl.* **1996**, 35, 2520–2522.
- [5] G. Viau, M. G. Lombardi, G. De Munno, M. Julve, F. Lloret, J. Faus, A. Caneschi, J. M. Clemente-Juan, *Chem. Commun.* **1997**, 1195–1196.
- [6] J. Ribas, M. Monfort, B. Kumar-Gosh, X. Solans, *Angew. Chem.* **1994**, 106, 2177–2179; *Angew. Chem. Int. Ed. Engl.* **1994**, 33, 2087–2089.
- [7] J. Ribas, M. Monfort, X. Solans, M. Drillon, *Inorg. Chem.* **1994**, 33, 742–745.
- [8] Crystal data (C₇H₁₇N₈Ni): monoclinic, space group *P*2₁/*c*, *a* = 10.0375(13), *b* = 8.6285(6), *c* = 13.7710(14) Å, β = 95.941(14)°, *V* = 1186.3(2) Å³, ρ_{calcd} = 1.523 g cm⁻³; Stoe Image Plate diffractometer, MoKα radiation λ = 0.71073 Å, *T* = 293 K; 8174 measured (2.8 ≤ θ ≤ 25°) and 2092 independent reflections used in the refinement; Lorentz polarization but no absorption corrections (μ = 1.551 mm⁻¹). The structure was solved by direct methods by using SHELXS97 and refined with SHELXL97, 145 parameters. H computed and refined isotropically, *R*(on *F*) = 0.017, ω*R*(on *F*²) = 0.009, residual electron density 0.066 e Å⁻³. Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-121920. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [9] O. Castell, R. Caballol, V. M. García, K. Handrick, *Inorg. Chem.* **1996**, 35, 1609–1615.
- [10] E. Ruiz, J. Cano, S. Alvarez, P. Alemany, *J. Am. Chem. Soc.* **1998**, 120, 11122–11129.
- [11] A. Escuer, R. Vicente, J. Ribas, M. Salah el Fallah, X. Solans, M. Font-Bardía, *Inorg. Chem.* **1994**, 33, 1842–1847.
- [12] R. Cortés, K. Urtiaga, L. Lezama, J. L. Pizarro, A. Goñi, M. I. Arriortua, T. Rojo, *Inorg. Chem.* **1994**, 33, 4009–4015.
- [13] A. Escuer, C. J. Harding, Y. Dussart, J. Nelson, V. McKee, R. Vicente, *J. Chem. Soc. Dalton Trans.* **1999**, 223–227.
- [14] C. S. Hong, Y. Do, *Angew. Chem.* **1999**, 111, 153–155; *Angew. Chem. Int. Ed.* **1999**, 38, 193–194.
- [15] H. Miyasaka, N. Matsumoto, H. Okawa, N. Re, E. Gallo, C. Floriani, *J. Am. Chem. Soc.* **1996**, 118, 981–994.

Received: May 27, 1999

Revised: September 13, 1999 [Z13474]