


- [8] a) J.-M. Lehn, A. Rigault, J. Siegel, J. Harrowfield, B. Chevrier, D. Moras, *Proc. Natl. Acad. Sci. USA* **1987**, *84*, 2565–2569; b) J.-M. Lehn, A. Rigault, *Angew. Chem.* **1988**, *100*, 1121–1122; *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 1095–1097.
- [9] R. Krämer, J.-M. Lehn, A. De Cian, J. Fischer, *Angew. Chem.* **1993**, *105*, 764–767; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 703–706.
- [10] a) B. Hasenknopf, J.-M. Lehn, B. O. Kneisel, G. Baum, D. Fenske, *Angew. Chem.* **1996**, *108*, 1987–1990; *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 1838–1840; b) B. Hasenknopf, J.-M. Lehn, N. Boumediene, A. Dupont-Gervais, A. Van Dorsselaer, B. Kneisel, D. Fenske, *J. Am. Chem. Soc.* **1997**, *119*, 10956–10962.
- [11] a) T. Gulik-Krzywicki, C. Fouquey, J.-M. Lehn, *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 163–167; b) N. Kimizuka, S. Fujikawa, H. Kuwahara, T. Kunitake, A. Marsh, J.-M. Lehn, *J. Chem. Soc. Chem. Commun.* **1995**, 2103–2104.
- [12] a) G. S. Hanan, J.-M. Lehn, N. Kyritsakas, J. Fischer, *J. Chem. Soc. Chem. Commun.* **1995**, 765–766; b) G. S. Hanan, U. S. Schubert, D. Volkmer, E. Rivière, J.-M. Lehn, N. Kyritsakas, J. Fischer, *Can. J. Chem.* **1997**, *75*, 169–182.
- [13] a) D. M. Bassani, J.-M. Lehn, G. Baum, D. Fenske, *Angew. Chem.* **1997**, *109*, 1931–1933; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1845–1847; b) D. M. Bassani, J.-M. Lehn, *Bull. Soc. Chim. Fr.* **1997**, *134*, 897–906.
- [14] M. Ohkita, J.-M. Lehn, G. Baum, D. Fenske, *Chem. Eur. J.* **1999**, *12*, 3471–3481.
- [15] a) K. T. Potts, *Bull. Soc. Chim. Belg.* **1990**, *99*, 741–768; b) K. T. Potts, K. A. Gheysen Raiford, M. Keshavarz-K., *J. Am. Chem. Soc.* **1993**, *115*, 2793–2807.
- [16] A. J. Majeed, Ø. Antonsen, T. Benneche, K. Undheim, *Tetrahedron* **1989**, *45*, 993–1006.
- [17] S. T. Howard, *J. Am. Chem. Soc.* **1996**, *118*, 10269–10274.
- [18] F. R. Heitzler, M. Neuburger, M. Zehnder, E. C. Constable, *Liebigs Ann.* **1997**, 297–301.
- [19] a) Assignment of the C4 and C5 protons can be deduced from the presence of multiple 3J coupling only possible for these protons. b) The association constants calculated from the shifts of the C4 and C5 proton resonances differ from that obtained from the remaining aromatic protons, assuming a simple monomer–dimer equilibrium; this may be due to the fact that intermolecular self-association should increase intramolecular helical self-organization. A value in the 10^3 M^{-1} range may be grossly estimated.
- [20] a) R. H. Martin, N. Defay, H. P. Figeys, M. Flammang-Barbieux, J. P. Cosyn, M. Gelbecke, J. J. Schurter, *Tetrahedron* **1969**, *25*, 4985–4998; b) H. Wynberg, *Acc. Chem. Res.* **1971**, *4*, 65–73.
- [21] A. S. Shetty, J. Zhang, J. S. Moore, *J. Am. Chem. Soc.* **1996**, *118*, 1019–1027.
- [22] J. van Esch, F. Schoonbeek, M. De Loos, E. M. Veen, R. M. Kellogg, B. L. Feringa in *Supramolecular Science: Where It Is and Where It Is Going* (Eds.: R. Ungaro, E. Dalcanele), Kluwer, Dordrecht, **1999**, pp. 233–259.
- [23] H. Engelkamp, S. Middelbeek, R. J. M. Nolte, *Science* **1999**, *284*, 785–788; for another case, see: W. Yang, X. Chai, L. Chi, X. Liu, Y. Cao, R. Lu, Y. Jiang, X. Tang, H. Fuchs, T. Li, *Chem. Eur. J.* **1999**, *5*, 1144–1149.
- [24] a) J.-M. Lehn, J. Malthête, A.-M. Levelut, *J. Chem. Soc. Chem. Commun.* **1985**, 1794–1795; b) N. Khazanovich, J. R. Granja, D. E. McRee, R. A. Milligan, R. Ghadiri, *J. Am. Chem. Soc.* **1994**, *116*, 6011–6012.
- [25] J.-M. Lehn, *Supramolecular Chemistry. Concepts and Perspectives*, VCH, Weinheim, **1995**, chap. 8.

A New Enantioselective Approach to Total Synthesis of the Securinega Alkaloids: Application to (–)-Norsecurinine and Phyllanthine**

Gyoonhee Han, Matthew G. LaPorte, James J. Folmer, Kim M. Werner, and Steven M. Weinreb*

Dedicated to the memory of Professor George Buchi

The securinega alkaloids are a family of approximately 20 tetracyclic compounds produced by several *Securinega* and *Phyllanthus* species of the Euphorbiaceae plant family.^[1] The major securinega alkaloid is securinine (**1**), isolated from the leaves of *Securinega suffruticosa*. Its enantiomer, virosecurinine, is also naturally occurring, as are the antipodes of



several other alkaloids of this class. Allosecurinine (**3**), which is the C2 epimer of **1**, is also a common alkaloid of this group, as is securitinine (**4**). The enantiomer of **3** (viroallosecurinine) is a natural product as well. Some additional common alkaloids of this group are (–)-norsecurinine (**5**), whose (+)-enantiomer is also naturally occurring, and phyllanthine (**2**).

These alkaloids are associated with a wide range of biological activity. Securinine causes central nervous system stimulation and clonic–tonic convulsions in laboratory animals. This activity is due to the alkaloid acting as a γ -amino butyric acid (GABA) receptor antagonist.^[2] These alkaloids have found a sporadic use clinically for such diseases as poliomyelitis, ALS (amyotrophic lateral sclerosis) and chronic aplastic anemia.^[1] Securinine has also been found to be an antimalarial agent^[3] and to have antibacterial activity.^[4]


To date there has been only limited synthetic work in this area. A nonstereoselective total synthesis of racemic securinine (**1**) was described by Horii et al. in 1966.^[5] More recently, Heathcock and coworkers devised a clever route to racemic norsecurinine (**5**).^[6] In this synthesis, both the A-ring and the configuration of the C2 atom of **5** were derived from L-proline, although unfortunately optical activity was lost by the racemization of an intermediate. Jacobi et al.^[7] have developed an innovative route to both (+)- and (–)-norsecurinine,

[*] Prof. Dr. S. M. Weinreb, Dr. G. Han, M. G. LaPorte, Dr. J. J. Folmer, Dr. K. M. Werner
Department of Chemistry
The Pennsylvania State University
University Park, PA 16802 (USA)
Fax: (+1) 814-865-3314
E-mail: smw@chem.psu.edu

[**] We are grateful to the National Institutes of Health (GM-32299) for financial support of this research.

starting from L- and D-proline, respectively, once again as a source of the five-membered A-ring and the absolute configuration of the C2 atom. Finally, the Magnus group^[8] has reported an elegant synthesis of racemic norsecurinine via the related securinega alkaloid nirurine, starting from 3-hydroxypyridine.


We have been attempting to develop a novel, general strategy for construction of the securinega alkaloids and have established two fundamental design criteria: 1) since most members of this family differ only in the A-ring size and functionality, as well as the C2 configuration (compare **1–5**), the strategy should be flexible enough to allow for the variations in these areas of the molecules; 2) the approach should provide access to both enantiomeric series of alkaloids. With these points in mind, our basic strategy (Scheme 1) was to use readily available *trans*-4-hydroxy-L-proline (**8**) (or the



Scheme 1. Retrosynthetic pathway for securinega alkaloids **6** using 4-hydroxy-L-proline (**8**) as a starting material.

commercially available D-enantiomer) as a source of the alkaloid B-ring and the C7 configuration.^[9] The intent was to first elaborate **8** into a BC-ring fragment like **7**, followed by annulation of the desired A-ring having the appropriate stereogenicity at C2. Finally, the butenolide D-ring would be appended at a late stage to afford the alkaloids **6**. Herein we describe the successful application of this concept to a new enantioselective total synthesis of (–)-norsecurinine (**5**) and the first total synthesis of phyllanthine (**2**).


The enantiomerically pure ester **9**, prepared from *trans*-4-hydroxy-L-proline (**8**),^[10] was converted via the α,β -unsaturated nitrile **10** to ketonitrile **11** (Scheme 2). We were pleased to find that exposure of **11** to SmI₂, followed by acidic



Scheme 2. a) TBSCl, imidazole, CH₂Cl₂, 88%; b) DIBAL-H, PhMe, –78 °C → rt, 98%; c) DMSO, (COCl)₂, NEt₃, CH₂Cl₂, –78 °C/Ph₃P=CHCN, CH₂Cl₂, –78 °C → rt, 90% (7:2 *E/Z* mixture); d) 1 atm H₂, 10% Pd-C, EtOAc, 98%; e) Bu₄NF, THF, rt, 100%; f) Jones reagent, acetone, rt, 84%; g) SmI₂, MeOH, THF, –78 °C → rt/H₃O⁺, 78%; h) TBSOTf, iPr₃NEt, CH₂Cl₂, 0 °C → rt, 91%; i) (CH₂O)₂, *p*-TsOH, PhH, reflux, 100%.

hydrolysis, provided the key scalemic bicyclic BC-ring system **12** in good yield.^[11] Hydroxyketone **12** was then transformed into protected ketal silyl ether derivative **13**.


Intermediate **13** was converted to the (–)-norsecurinine ABC-ring synthon as outlined in Scheme 3. The *N*-tosyl group of **13** was removed to afford amine **14**, which was acylated with isatoic anhydride to yield the *ortho*-aminobenzamide



Scheme 3. a) Na, naphthalene, DME, –78 °C, 88%; b) isatoic anhydride, DMAP, MeCN, rt, 87% from **13**; c) NaNO₂, HCl, CuCl, MeOH, rt; d) allylmagnesium bromide, BF₃·Et₂O, THF, –78 → 0 °C; e) (Boc)₂O, NEt₃, CH₂Cl₂, reflux, 68%; f) disiamyl borane, THF, 0 °C → rt/H₂O₂, NaOH; g) TsCl, DMAP, NEt₃, CH₂Cl₂, 71% from **20**; h) 3% HCl, MeOH, 60 °C, 81%.

derivative **15**. In a key step which utilizes our recently reported method,^[12] amide **15** could be regioselectively converted to a mixture of α -chloro- and α -methoxybenzamides **16** and **17**, respectively, in a good combined yield. Subsequent treatment of this mixture with excess allylmagnesium bromide/BF₃·Et₂O stereoselectively afforded the allyl amine **19**, which was transformed without purification into the *tert*-butoxycarbonyl (Boc) derivative **20**. Formation of alkylation product **19** undoubtedly proceeds via the *N*-acyliminium species **18**, which is exclusively *exo*-attacked by the Grignard reagent, followed by debenzoylation by the excess organometallic reagent. The configuration of **20** was established by X-ray analysis of the derivative **25a** (see below). To complete formation of the requisite tricyclic ABC-species, olefin **20** underwent hydroboration, followed by treatment of the resulting primary alcohol with tosylate, giving sulfonate **21**. Removal of the Boc group from **21** resulted in direct in situ cyclization to ketal silyl ether **22**.


Mild acidic hydrolysis of **22** then led to the pivotal tricyclic α -hydroxy ketone **23a** (Scheme 4). This compound could be reacted with the ketene ylide reagent of Bestmann,^[13] a reaction which is best conducted in this case at high pressure (12 kbar), to afford (+)-14,15-dihydronorsecurinine (**25a**), a naturally occurring minor alkaloid isolated along with (–)-norsecurinine from the root bark of *Securinega virosa*.^[14] The structure of the synthetic alkaloid **25a** was firmly established by an X-ray analysis of its hydrochloride salt.^[15]



Scheme 4. a) 3 N HCl, 95 °C, 74 %; b) PhSeCl, EtOAc, NEt₃, reflux, 60 %; c) PhSeCl, EtOAc, py, reflux, 67 %; d) Ph₃P=C=O, PhMe, CH₂Cl₂, 12 kbar, rt, 89 %; e) (EtO)₂POCH₂CO₂H, CMC, CH₂Cl₂, 0 °C → rt, 88 %; f) K₂CO₃, [18]crown-6, PhMe, 0 °C → rt, 95 %; g) dimethyldioxirane, CH₂Cl₂/Me₂CO, -78 °C, 39 %.

In order to prepare (–)-norsecurinine itself, we next investigated conversion of hydroxy ketone **23a** to the corresponding enone **24b** (Scheme 4). Despite extensive effort we were unable to effect the α -phenylselenation of **23a**, or any hydroxyl-protected derivatives, using standard enolate selenation methods.^[16] On the other hand, we did find that application of an alternate method based on the work of Liotta^[17] (PhSeCl, EtOAc, pyridine, reflux) to substrate **23a** did not afford the anticipated α -selenophenylketone **23b**, but surprisingly gave α -selenophenylene **24a** as the only discernible product. However, we have been unable to remove the selenophenyl group from **24a** to produce the enone **24b**. Interestingly, we also discovered that simply changing the selenation base from pyridine to triethylamine allowed conversion of ketone **23a** to the desired α -phenylselenoketone **23b**. Disappointingly, all attempts to effect elimination of the selenophenyl group from **23b** via the corresponding selenoxide species^[16a] led to a number of decomposition products, probably due to the instability of the resulting enone **24b**. Despite these problems, it was possible to successfully complete the alkaloid total synthesis by converting the keto selenide **23b** to the phosphonate species **26**, which could then be cyclized to butenolide derivative **25b**.^[18] Finally, treatment of the butenolide **25b** with dimethyldioxirane at low temperature led directly to (–)-norsecurinine (**5**) having spectral data identical with those of authentic material.^[16c, 19]

In order to test the generality of our strategy for synthesis of the securinega alkaloids, we have extended the approach to phyllanthine (**2**). Thus, the bicyclic amine **14** could be cleanly oxidized with iodosobenzene^[20] to afford the imine **27** (Scheme 5). We were pleased to find that this imine undergoes an *exo*-stereoselective hetero Diels–Alder reaction with Danishefsky's diene under high pressure (yield ca. 70 %), or more conveniently and efficiently upon catalysis by Yb(OTf)₃,^[21, 22] to afford the tricyclic vinylogous amide **28** after aqueous workup (yield 84 %).^[23] Treatment of **28** with two equivalents of L-selectride effected both conjugate



Scheme 5. a) PhIO, CH₂Cl₂, rt, 87 %; b) Danishefsky's diene, Yb(OTf)₃, MeCN, 84 %; c) L-selectride (2 equiv.), THF, -78 °C, 85 %; d) NaH, MeI, THF, 0 °C → rt, 87 %; e) 3 N HCl, reflux, 78 %; f) (PhSe)₂, SeO₂, MsOH, CH₂Cl₂, 0 °C → rt, 58 %; g) NaI, BF₃·Et₂O, MeCN, 0 °C, 84 %; h) (EtO)₂POCH₂CO₂H, DCC, THF, 0 °C → rt, 67 %; i) K₂CO₃, [18]crown-6, PhMe, 0 °C → rt, 84 %.

reduction and subsequent stereoselective reduction of the ketone group, yielding the axially oriented alcohol **29**. This compound could then be O-methylated and the protecting groups removed by acid hydrolysis to afford hydroxyketone **30**. Attempted application to ketone **30** of the phenylselenation method^[17] used for **23a** (see above) resulted in a complex mixture of products. However, using the selenation procedure reported by Sonoda's group (PhSeSePh, MsOH, SeO₂),^[24] **30** was transformed to the α -phenylselenoenone **31**, which was deselenated in good yield with NaI/BF₃·Et₂O^[25] to provide the required enone **32**. To complete the synthesis, hydroxyenone **32** was esterified to give phosphonate **33**.^[26] Cyclization of **33**, promoted by potassium carbonate/[18]crown-6, then afforded phyllanthine (**2**) which had ¹H and ¹³C NMR data in good accord with those reported for the natural product.^[27]

In summary, we have successfully implemented a new general strategy for enantioselective construction of the securinega alkaloids. Key steps in the total synthesis of (–)-norsecurinine (**5**) include an intramolecular SmI₂-catalyzed ketonitrile coupling and application of our new method for generation of *N*-acylimines.^[12] The total synthesis of phyllanthine (**2**) features a stereoselective imino Diels–Alder reaction for construction of the A-ring.

Received: August 27, 1999 [Z13932]

- [1] For reviews see: a) V. Snieckus, *The Alkaloids*, **1975**, *14*, 425; b) J. A. Beutler, A. N. Brubaker, *Drugs Future* **1987**, *12*, 957.
- [2] D. Rognan, T. Boulanger, R. Hoffmann, D. P. Vercauteren, J.-M. Andre, F. Durant, C.-G. Wermuth, *J. Med. Chem.* **1992**, *35*, 1969.
- [3] H. Weenen, M. H. H. Nkunya, D. H. Bray, L. B. Mwasumbi, L. S. Kinabo, V. A. Kilimali, J. B. Wijnberg, *Planta Med.* **1990**, *56*, 371.
- [4] J. L. Mensah, I. Lagarde, C. Ceschin, G. Michel, J. Gleye, I. Fouraste, *J. Ethnopharmacol.* **1990**, *28*, 129.
- [5] a) S. Saito, H. Yoshikawa, Y. Sato, H. Nakai, N. Sugimoto, Z. Horii, M. Hanaoka, Y. Tamura, *Chem. Pharm. Bull.* **1966**, *14*, 313; b) Z. Horii, T. Imanishi, T. Tanaka, I. Mori, M. Hanaoka, C. Iwata, *Chem. Pharm.*


- Bull.* **1972**, *20*, 1768; c) Z. Horii, T. Imanishi, M. Hanaoka, C. Iwata, *Chem. Pharm. Bull.* **1972**, *20*, 1774.
- [6] a) C. H. Heathcock, R. A. Jennings, T. W. von Geldern, *J. Org. Chem.* **1983**, *48*, 3428; b) C. H. Heathcock, T. W. von Geldern, *Heterocycles* **1987**, *25*, 75.
- [7] P. A. Jacobi, C. A. Blum, R. W. DeSimone, U. E. S. Udodong, *J. Am. Chem. Soc.* **1991**, *113*, 5384.
- [8] P. Magnus, J. Rodriguez-Lopez, K. Mulholland, I. Matthews, *Tetrahedron* **1993**, *49*, 8059.
- [9] Interestingly, the Heathcock^[6] and Jacobi^[7] strategies leading to the more common (–)-enantiomer of norsecurinine (**5**) require D-proline whereas our approach to (–)-**5** uses the less-expensive L-amino acid.
- [10] R. H. Andreatta, V. Nair, A. V. Robertson, W. R. J. Simpson, *Aust. J. Chem.* **1967**, *20*, 1493.
- [11] Compare: G. A. Molander, C. N. Wolfe, *J. Org. Chem.* **1998**, *63*, 9031, and references therein. See also: Y. Yamamoto, D. Matsumi, R. Hattori, K. Itoh, *J. Org. Chem.* **1999**, *64*, 3224.
- [12] a) G. Han, M. C. McIntosh, S. M. Weinreb, *Tetrahedron Lett.* **1994**, *35*, 5813; b) S. M. Weinreb, *J. Heterocycl. Chem.* **1996**, *33*, 1437; c) G. Han, M. G. LaPorte, M. C. McIntosh, S. M. Weinreb, *J. Org. Chem.* **1996**, *61*, 9483.
- [13] a) H. J. Bestmann, D. Sandmeier, *Chem. Ber.* **1980**, *113*, 274; b) K. Nickisch, W. Klose, F. Bohlmann, *Chem. Ber.* **1980**, *113*, 2038.
- [14] a) S. Saito, T. Tanaka, K. Kotera, H. Nakai, N. Sugimoto, Z.-I. Horii, M. Ikeda, Y. Tamura, *Chem. Pharm. Bull.* **1964**, *12*, 1520; b) S. Saito, T. Tanaka, K. Kotera, H. Nakai, N. Sugimoto, Z.-I. Horii, M. Ikeda, Y. Tamura, *Chem. Pharm. Bull.* **1965**, *13*, 786.
- [15] We thank Dr. M. Parvez (University of Calgary) for this crystal structure determination. We also are indebted to Dr. C. J. Easton and J. Kelly (Australian National University) for obtaining an ORD spectrum of **25a**, which established that our synthetic material is in the same enantiomeric series as the natural compound.^[14]
- [16] a) H. J. Reich, S. Wollowitz, *Org. React.* **1993**, *44*, 1; b) attempted DDQ or Saegusa oxidations of **23a** and **25a** also failed; c) other selenide oxidants such as *m*-CPBA, O₃, or NaIO₄ gave significantly lower yields of **5**.
- [17] A similar reaction leading directly to an α -selenophenyleneone was described: D. Liotta, C. Barnum, R. Puleo, G. Zima, C. Bayer, H. S. Kezar III, *J. Org. Chem.* **1981**, *46*, 2920. See also: K. B. Sharpless, R. F. Lauer, A. Y. Teranishi, *J. Am. Chem. Soc.* **1973**, *95*, 6137.
- [18] a) Condensation of α -phenylselenoketone **23b** with the Bestmann ketene ylide^[13] gave only a low yield of butenolide **25b**; b) selenides **23b**, **25b** and **26** are mixtures of epimers at C14.
- [19] We are grateful to Professor Peter Jacobi for providing copies of the IR, NMR, CD and mass spectra of authentic norsecurinine.^[7]
- [20] a) P. Muller, D. M. Gilabert, *Tetrahedron* **1988**, *44*, 7171; b) J. Larsen, K. A. Jorgensen, *J. Chem. Soc. Perkin Trans. 2* **1992**, 1213.
- [21] For reviews and lead references to imino Diels–Alder reactions, see: a) D. L. Boger, S. M. Weinreb, *Hetero Diels–Alder Methodology in Organic Synthesis*, Academic Press, Orlando, **1987**, ch. 2; b) S. M. Weinreb in *Comprehensive Organic Synthesis*, Vol. 4 (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, p. 401.
- [22] Compare: S. Kobayashi, S. Nagayama, *J. Am. Chem. Soc.* **1997**, *119*, 10049.
- [23] The crystal structure and configuration of cycloadduct **28** were confirmed by X-ray analysis. We thank Dr. M. Shang (University of Notre Dame) for determining this structure.
- [24] N. Miyoshi, T. Yamamoto, N. Kambe, S. Murai, N. Sonoda, *Tetrahedron Lett.* **1982**, *23*, 4813. See also: M. Watanabe, B. Z. Awen, M. Kato, *J. Org. Chem.* **1993**, *58*, 3923. This procedure is useful for forming an α -phenylselenoketone from a ketone, although direct formation of an α -phenylselenoone as occurred with **30** has apparently not previously been observed.
- [25] This protocol was previously used to dehalogenate an α -bromoone: A. K. Mandal, S. W. Mahajan, *Tetrahedron* **1988**, *44*, 2293.
- [26] Reaction of hydroxy enone **31** with the Bestmann ketene ylide^[13] does not give phyllanthine.
- [27] a) J. Parelo, *Bull. Soc. Chim. Fr.* **1968**, 1117; b) D. Arbain, A. A. Birkbeck, L. T. Byrne, M. V. Sargent, B. W. Skelton, A. H. White, *J. Chem. Soc. Perkin Trans. 1* **1991**, 1863.

Formal Three-Electron Reduction by an f-Element Complex: Formation of [(C₅Me₅)(C₈H₈)U]₂(C₈H₈) from Cyclooctatetraene and [(C₅Me₅)₃U]**

William J. Evans,* Gregory W. Nyce, and Joseph W. Ziller

Reduction reactions involving more than two electrons are not common for metal complexes containing just one metal.^[1] In f-element chemistry, multielectron reductions from mono-metallic species are particularly rare^[2] since the common low-valent ions Th^{III}, U^{III}, Eu^{II}, Yb^{II}, Sm^{II}, and Tm^{II} typically react as one-electron An^{IV}/An^{III} or Ln^{III}/Ln^{II} redox couples.^[3–5] Recently however, we have shown that there is a new way of accomplishing reductive chemistry with f-element complexes by steric crowding^[6,7] and we report here how this method can be coupled with a traditional one-electron redox couple to accomplish multielectron reduction with a mono-metallic uranium complex.

It has recently been discovered that the sterically crowded [(C₅Me₅)₃Ln] complexes (Ln = Sm,^[6] Nd^[7]) can act as one-electron reductants as shown in Equation (1). The reductive reactivity of the [(C₅Me₅)₃Ln] complexes can be rationalized



by the fact that the C₅Me₅[–] ions are not well-stabilized electrostatically in these compounds due to the long Ln–C(C₅Me₅) distances caused by steric congestion. If the redox reaction in Equation (1) is coupled to a one-electron metal-based reduction, for example, the U^{IV}/U^{III} couple^[8] of uranium, then a complex such as [(C₅Me₅)₃U]^[9] could be a two-electron reducing agent. This concept was tested by reacting [(C₅Me₅)₃U] with 1,3,5,7-cyclooctatetraene (C₈H₈). It had been shown that U^{III} complexes would act as one-electron reductants with cyclooctatetraene.^[10] Our efforts to couple that reactivity with sterically induced reduction are reported here.

Reaction of [(C₅Me₅)₃U] with C₈H₈ in a 1:1 stoichiometry formed (C₅Me₅)₂ and a new product, **1**, but residual [(C₅Me₅)₃U] remained at the end of the reaction. Subsequently, it was determined that a 2:3 [(C₅Me₅)₃U]:C₈H₈

[*] Prof. W. J. Evans, G. W. Nyce, Dr. J. W. Ziller
Department of Chemistry, University of California
Irvine, CA 92697 (USA)
Fax: (+1) 949-824-2210
E-mail: wevans@uci.edu

[**] We thank the US National Science Foundation for support of this research.