


- [4] a) H.-F. Hsu, J. R. Shapley, *J. Am. Chem. Soc.* **1996**, *118*, 9192–9193; b) K. Lee, H.-F. Hsu, J. R. Shapley, *Organometallics* **1997**, *16*, 3876–3877; c) K. Lee, J. R. Shapley, *Organometallics* **1998**, *17*, 3020–3026.
- [5] a) J. T. Park, J.-J. Cho, H. Song, *J. Chem. Soc. Chem. Commun.* **1995**, 15–16; b) J. T. Park, J.-J. Cho, H. Song, C.-S. Jun, Y. Son, J. Kwak, *Inorg. Chem.* **1997**, *36*, 2698–2699; c) J. T. Park, H. Song, J.-J. Cho, M.-K. Chung, J.-H. Lee, I.-H. Suh, *Organometallics* **1998**, *17*, 227–236; d) H. Song, K. Lee, J. T. Park, M.-G. Choi, *Organometallics* **1998**, *17*, 4477–4488.
- [6] J. M. Fernandez, B. F. G. Johnson, J. Lewis, P. R. Raithby, *J. Chem. Soc. Dalton Trans.* **1981**, 2250–2257.
- [7] P. F. Jackson, B. F. G. Johnson, J. Lewis, J. N. Nicholls, M. McPartlin, W. J. H. Nelson, *J. Chem. Soc. Chem. Commun.* **1980**, 564–566.
- [8] B. F. G. Johnson, J. Lewis, J. N. Nicholls, J. Puga, P. R. Raithby, M. J. Rosales, M. McPartlin, W. Clegg, *J. Chem. Soc. Dalton Trans.* **1983**, 277–290.

Preparation, Structure, and Redox Reactions of Nine-Membered Cyclic Peroxides: A Novel Electrochromic System Undergoing Reversible Extrusion and Trapping of O₂**

Takanori Suzuki,* Jun-ichi Nishida, Masakazu Ohkita, and Takashi Tsuji

Cyclic peroxides having a medium-size ring skeleton have recently attracted much attention from a pharmacological standpoint^[1] and the mechanistic studies of their electron-transfer reactions^[2,3] have provided the basis for a better understanding of their biological activities. The nine-membered peroxides, however, are so rare^[4] that their chemistry has not been well developed. During the course of our study on molecular systems in which the structures and properties can be controlled by electron transfer (ET)^[5] we have found that the dihydrotrioxonin derivatives **1**, a new class of nine-membered cyclic peroxides having a hitherto unknown 1,2,6-trioxacyclononane skeleton, could be generated very easily. Here we report the preparation and X-ray structure of **1** along with their unique electron-transfer behavior. It is worth noting that ET induces the oxidative deoxygenation of peroxides **1** to form the dicationic dyes **2**²⁺, from which compounds **1** are

recovered upon 2e reduction in the presence of O₂ (Scheme 1). This system could be used to electrochemically control O₂ concentration.



Scheme 1. Interconversion between **1** and **2**²⁺ by the oxidative-deoxygenation and reductive-oxygenation reactions.

Diols **3a** and **3b**^[6] were prepared in 54 and 41 % yield, respectively, by the reaction of 2,2'-dilithiodiphenyl ether^[7] with xanthone and thioxanthone. Deeply colored salts **2a**·(BF₄)₂^[6] [λ_{max} (lg ϵ) 480 (sh, 3.73), 450 (3.84), 378 (4.71), 260 nm (4.86) in MeCN] and **2b**·(BF₄)₂^[6] [531 (sh, 3.78), 498 (3.88), 387 (4.47), 282 (5.09)] were obtained in 94 and 91 % yield, respectively, by treating these diols with HBF₄ in (EtCO)₂O.

According to the voltammetric analyses the dicationic dye undergoes facile and reversible 2e reduction (**2a**²⁺: $E^{\text{red}} = +0.29$ V (2e); **2b**²⁺: $E^{\text{red}} = +0.26$, $E_2^{\text{red}} = +0.05$ V versus the saturated calomel electrode (SCE)),^[8] which indicates that the diradical **2**^{••} is a long-lived species in an argon atmosphere. In contrast, only irreversible reduction waves were observed in both cases when the same solutions were saturated with O₂ (Figure 1). On the basis of a detailed examination of the cyclic voltammograms, it seems likely that the cation radical intermediate **2**^{•+} is readily trapped by O₂ to form (**2-O**)^{•+}, which is further reduced to (**2-O**)^{••} at a slightly more negative potential than **2**²⁺ (ECE process). The resultant diradical (**2-O**)^{••} undergoes ring closure to the entropically disfavored nine-membered ring with unexpectedly high efficiency. Thus, peroxides **1a**^[6] [λ_{max} (lg ϵ) 291 (3.92), 284 nm (sh, 3.90) in MeCN] and **1b**^[6] [285 (sh, 4.02), 267 (4.28)] were isolated as colorless crystals in 80 and 71 % yield, respectively, when the **2**·(BF₄)₂ salts were reduced with Zn powder in aerated THF.^[9]

X-ray analyses^[10] on the novel heterocycles **1** (Figure 2) have revealed the extended conformation of the peroxide moiety (torsion angle for the C–O–O–C unit: 160.8(3)° for **1a**; 154.6(2)° for **1b**), and their O–O bond lengths (1.507(4) Å for **1a**; 1.502(3) Å for **1b**) are two of the longest values ever reported^[11]. Yet, the electrochemical oxidation does not break this weak O–O bond, but instead breaks the adjacent C–O bonds.^[3] It is also interesting to note that deoxygenation and

[*] Prof. Dr. T. Suzuki, J. Nishida, Dr. M. Ohkita, Prof. Dr. T. Tsuji
Division of Chemistry
Graduate School of Science, Hokkaido University
Sapporo 060-0810 (Japan)
Fax: (+81)11-746-2557
E-mail: tak@sci.hokudai.ac.jp

[**] This work was supported by the Ministry of Education, Science, and Culture, Japan (No. 10146101). Financial support by the Izumi Science and Technology Foundation and by a Research Grant from the Iwatani Naoji Foundation (T.S.) are gratefully acknowledged. J.N. thanks JSPS Research Fellowships for Young Scientists. We thank Prof. Tamotsu Inabe (Hokkaido University) for the use of facilities to analyze the X-ray structures. Technical support by Mr. Kenji Watanabe and Dr. Eri Fukushi (GC-MS & NMR Laboratory, Faculty of Agriculture, Hokkaido University) is gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.




Figure 1. Cyclic voltammogram of dication **2b**²⁺ a) under Ar and b) under O₂ (in CH₂Cl₂ containing 0.1 mol dm⁻³ nBu₄NBF₄, Pt electrode, scan rate 500 mV s⁻¹). The first reduction wave at +0.21 V in (b) is irreversible even when the scanning was reversed at +0.15 V just after the 1e reduction of **2b**²⁺ (dotted line). The voltammogram of **2a**²⁺ measured under O₂ also shows two well separated, irreversible peaks at +0.15 and -0.06 V. Full details of the voltammograms of **1** and **2**²⁺ are given in the Supporting Information.




Figure 2. Molecular geometry of peroxide **1a** determined by X-ray analysis with the selected bond lengths. Filled ellipsoids correspond to O atoms and dotted ones indicate C atoms comprising the nine-membered ring. X-ray analysis showed that **1b** adopts a similar molecular structure to **1a** although the deformation of the thioxanthene rings into a butterfly shape is much more pronounced.

not oxygenation is induced upon oxidation. Thus, the voltammograms of **1** show the irreversible oxidation waves (**1a**: +2.08 V; **1b** +1.89 V versus SCE in CH₂Cl₂), and their corresponding reduction peaks appear in the far cathodic region, which were assigned to those of the deoxygenated dications **2**²⁺. Furthermore, dication salts **2a**-(BF₄)₂ and **2b**-(BF₄)₂ were isolated in 85 and 65 % yield, respectively, when the peroxides **1** were treated with two equivalents of NOBF₄ in CH₂Cl₂. When 1.00 mmol of **1a** was treated with 2.06 equivalents of (2,4-Br₂C₆H₃)₃N⁺SbCl₆⁻ in 1,2-dichloroethane **2a**-(SbCl₆)₂ was obtained in 99 % yield. At the same time, 21.5 mL of gas evolved. This amount corresponds to 96 % of

the calculated volume of O₂ that would be released from **1a**.^[12]

These results indicate that **1** and **2**²⁺ constitute a novel redox system that undergoes reversible extrusion and trapping of O₂. The large changes in the UV/Vis spectrum upon electrolysis show the electrochromic nature^[13] of their interconversion (Figure 3). The several isosbestic points observed




Figure 3. a) Changes in the UV/Vis spectrum of **2a**²⁺ (3.5 mL; 2.07×10^{-5} mol dm⁻³ in MeCN containing 0.05 mol dm⁻³ nBu₄NBF₄) upon constant-current electrochemical reduction (40 μA) at 2 min intervals. b) Changes in the UV/Vis spectrum of **1b** (3.5 mL; 2.24×10^{-5} mol dm⁻³ in MeCN containing 0.05 mol dm⁻³ nBu₄NBF₄) upon constant-current electrochemical oxidation (70 μA) at 6 min intervals.

are indicative of the clean and quantitative transformation between **1** and **2**²⁺ although elucidation of the detailed reaction mechanism awaits further experiments.^[14]

Received: November 15, 1999 [Z14270]

Revised: January 17, 2000

- [1] A. Robert, B. Meunier, *Chem. Soc. Rev.* **1998**, 27, 273; C. W. Jefford, S. Kohmoto, D. Jaggi, G. Timári, J.-C. Rossier, M. Rudaz, O. Barbuzzi, D. Gérard, U. Burger, P. Kamalaprija, J. Mareda, G. Bernardinelli, I. Manzanares, C. J. Canfield, S. L. Fleck, B. L. Robinson, W. Peters, *Helv. Chim. Acta* **1995**, 78, 647; G. H. Posner, J. N. Cumming, P. Ploypradith, C. H. Oh, *J. Am. Chem. Soc.* **1995**, 117, 5885.
- [2] S. Antonello, M. Musumeci, D. D. M. Wayner, F. Maran, *J. Am. Chem. Soc.* **1997**, 119, 9541; R. L. Donkers, J. Tse, M. S. Workentin, *Chem. Commun.* **1999**, 135.
- [3] Y. Takahashi, K. Wakamatsu, S. Morishima, T. Miyashi, *J. Chem. Soc. Perkin Trans. 2* **1993**, 243; Y. Takahashi, S. Morishima, K. Wakamatsu, T. Suzuki, T. Miyashi, *J. Chem. Soc. Chem. Commun.* **1994**, 13.
- [4] Y. Ushigoe, Y. Torao, A. Masuyama, M. Nojima, *J. Org. Chem.* **1997**, 62, 4949; Y. Takahashi, M. Ando, T. Miyashi, *Tetrahedron Lett.* **1995**, 36, 1889; W. Fenical, *J. Am. Chem. Soc.* **1974**, 96, 5580.

- [5] T. Suzuki, J. Nishida, T. Tsuji, *Angew. Chem.* **1997**, *109*, 1387; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1329.
- [6] All new compounds gave satisfactory spectral data and analytical values. Selected ^1H NMR (400 MHz, TMS) data are as follows: **1a** (CDCl_3 , 24°C): δ = 7.47 (brd, J = 7.8 Hz, 4H), 7.40 (dd, J = 7.8, 1.5 Hz, 2H), 7.21–7.32 (m, 6H), 7.12 (m, 4H), 6.97 (brdd, J = 7, 7 Hz, 2H), 6.87–6.94 (m, 4H), 6.61 (brd, J = 7.8 Hz, 2H); **2a** (BF_4) $_2$ (CD_3CN , 20°C): δ = 8.48 (ddd, J = 6.8, 6.8, 1.5 Hz, 4H), 8.31 (brd, J = 8.3 Hz, 4H), 7.75 (ddd, J = 6.8, 6.8, 1.5 Hz, 2H), 7.73 (dd, J = 8.3, 1.5 Hz, 4H), 7.58 (ddd, J = 6.8, 6.8, 1.0 Hz, 4H), 7.47 (ddd, J = 7.8, 7.8, 1.0 Hz, 2H), 7.37 (dd, J = 7.8, 1.8 Hz, 2H), 7.32 (brd, J = 7.8 Hz, 2H).
- [7] H. Gilman, W. J. Trepka, *J. Org. Chem.* **1961**, *26*, 5202.
- [8] Comparisons of the coefficients obtained by linear combination of atomic orbitals (LCAO) in singly occupied molecular orbitals (SOMO) between xanthyl and thioxanthyl indicate that the considerable spin exists on S in thioxanthyl whereas much larger spin density is expected at C9 in xanthyl (K. Okada, T. Imakura, M. Oda, A. Kajiwaru, M. Kamachi, M. Yamaguchi, *J. Am. Chem. Soc.* **1997**, *119*, 5740). This fact may be related to the reason for the different number of electrons involved in E_1^{red} of **2** $^{2+}$. One possible explanation might be the rapid equilibrium of **2a** $^{2+}$ with some closed-shell species, which facilitates the disproportionation of **2a** $^{2+}$ to make the E_2^{red} wave ambiguous. Such a process is less likely in **2b** $^{2+}$, in which the spin density is delocalized over thioxanthyl moieties.
- [9] When the dication salt **2a** (BF_4) $_2$ was reduced by SmI_2 in THF followed by quenching with air, peroxide **1a** was obtained in 85% yield. This result indicates that (**2-O**) $^{2+}$ is also generated by 2e reduction of **2a** $^{2+}$ followed by C–O bond making (EEC process).
- [10] Crystal structure analyses: **1a**: $\text{C}_{38}\text{H}_{22}\text{O}_5$, M_r = 560.60, monoclinic, $P2_1/c$, a = 10.211(3), b = 16.311(5), c = 16.931(3) Å, β = 103.88(2) $^\circ$, V = 2737(1) Å 3 , Z = 4, ρ_{calc} = 1.360 g cm $^{-3}$, R_w = 0.069. **1b**: 0.5 AcOEt (-100°C): $\text{C}_{40}\text{H}_{28}\text{O}_4\text{S}_2$, M_r = 636.78, monoclinic, $C2/c$, a = 31.597(1), b = 10.2379(3), c = 22.9976(7) Å, β = 123.615(1) $^\circ$, V = 6195.3(3) Å 3 , Z = 8, ρ_{calc} = 1.365 g cm $^{-3}$, R_w = 0.055. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication nos. CCDC-136844 and -136845. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [11] The O–O bond length in 1,4-diphenyl-2,3-dioxabicyclo[2.2.1]heptane is 1.501(2) Å (D. J. Coughlin, R. S. Brown, R. G. Salomon, *J. Am. Chem. Soc.* **1979**, *101*, 1533) and that in tetrabenzopentacene endoperoxide is 1.496(5) Å (A. Izuoka, T. Murase, M. Tsukada, Y. Ito, T. Sugawara, A. Uchida, N. Sato, H. Inokuchi, *Tetrahedron Lett.* **1997**, *38*, 245). Long O–O distances are more common for small ring peroxides, such as dioxiranes (1.503(5) Å for dimesityldioxirane: W. Sander, K. Schroeder, S. Muthusamy, A. Kirschfeld, W. Kappert, R. Boese, E. Kraka, C. Sosa, D. Cremer, *J. Am. Chem. Soc.* **1997**, *119*, 7265) or dioxetanes (1.500 and 1.55 Å in 1,6-diphenyl-2,5,7,8-tetraoxobicyclo[4.2.0]octane derivatives: W. Adam, E. Schmidt, E.-M. Peters, K. Peters, H. G. von Schnering, *Angew. Chem.* **1983**, *95*, 566; *Angew. Chem. Int. Ed. Engl.* **1983**, *22*, 546). The last one is surprisingly long for an O–O bond although the error in the determination was not given.
- [12] When the similar oxidation was conducted under the presence of tetraphenylcyclopentadienone (10 equiv) no detectable amount of *cis*-dibenzoylstilbene was formed, which suggests that the major reaction path does not produce $^1\text{O}_2$.
- [13] P. M. S. Monk, R. J. Mortimer, D. R. Rosseinsky, *Electrochromism: Fundamentals and Applications*, VCH, Weinheim, **1995**, pp. 124–182; S. Hünig, M. Kemmer, H. Wenner, I. F. Perepichka, P. Bäuerle, A. Emge, G. Gescheid, *Chem. Eur. J.* **1999**, *5*, 1969.
- [14] Reductive oxygenation would also occur by the combination of **2** $^{2+}$ with $\text{O}_2^{\cdot-}$, which is generated by 1e reduction of O_2 . This process seems, however, less likely when it is considered that O_2 is a weaker electron acceptor ($E^{\text{red}} < -0.8$ V versus SCE in CH_2Cl_2) than **2** $^{2+}$.


A New Asymmetric Carbon–Carbon Bond Forming Reaction: Four-Component Stereoselective Synthesis of (Z)-4,6-Dihydroxy-3-methylalk-2-enyl Methyl Sulfones**

Vera Narkevitch, Kurt Schenk, and Pierre Vogel*

Dedicated to Professor Horst Prinzbach
on the occasion of his 68th birthday

In the presence of a Lewis acid, 1-alkoxy- or 1-silyloxy-1,3-dienes can be combined with enoxysilanes and sulfur dioxide to generate (Z)-6-oxo-4-oxyalk-2-ene sulfonates, which react with methyl iodide (S-alkylation) to afford the corresponding methyl sulfones.^[1, 2] We report here an asymmetric version of this new carbon–carbon bond forming reaction that can be used to construct polyketide fragments stereoselectively.

Enantiomerically pure (>99% ee) diene (+)-**2** was obtained by reaction of **1** with sodium (–)-(S)-1-(2,4,6-triisopropylphenyl)ethoxide^[3] followed by Wittig methylenation (Scheme 1).^[4] In the presence of $\text{Yb}(\text{OTf})_3$ ($\text{Tf} = \text{F}_3\text{CSO}_2$) and



Scheme 1. Asymmetric synthesis of 4,6-anti-(Z)-4,6-dihydroxy-3-methylalk-2-enyl methyl sulfones; see text for details.

an excess of SO_2 , (+)-**2** reacted with enoxysilane **3** to give a trimethylsilyl sulfinate, which was desilylated with Bu_4NF and treated with MeI to afford a 25:1 mixture of sulfone (–)-**4** and its diastereomer (79% yield, recovery of 20% of (+)-**2**).^[5]

[*] Prof. P. Vogel, V. Narkevitch
Section de chimie de l'Université de Lausanne, BCH
1015 Lausanne-Dorigny (Switzerland)
Fax: (+41) 21-692-39-75
E-mail: pierre.vogel@ico.unil.ch

K. Schenk
Institut de Cristallographie, BSP
Université de Lausanne, Lausanne-Dorigny (Switzerland)

[**] This work was supported by the Swiss National Science Foundation.