

- Soc.* **1994**, 116, 11014–11019; c) J. A. Ayllon, C. Gervaux, S. Sabo-Etienne, B. Chaudret, B. Hessen, *Organometallics* **1997**, 16, 2000–2002; d) S. Aime, R. Gobetto, E. Valls, *Organometallics* **1997**, 16, 5140–5141; e) H. S. Chu, C. P. Lau, K. Y. Wong, W. T. Wong, *Organometallics* **1998**, 17, 2768–2777; f) A. Cabalero, F. A. Jalón, B. R. Manzano, *Chem. Commun.* **1998**, 1879–1880.
- [3] a) J. P. Campbell, J. W. Hwang, V. G. Young, R. B. Von Dreele, C. J. Cramer, W. L. Gladfelter, *J. Am. Chem. Soc.* **1998**, 120, 521–531; b) K. Abdur-Rashid, D. G. Gusev, A. L. Lough, R. H. Morris, *Organometallics* **2000**, 19, 834–843.
- [4] a) R. Custelcean, J. E. Jackson, *J. Am. Chem. Soc.* **1998**, 120, 12935–12941; b) R. Custelcean, J. E. Jackson, *Angew. Chem.* **1999**, 111, 1748–1751; *Angew. Chem. Int. Ed.* **1999**, 38, 1661–1663; c) R. Custelcean, J. E. Jackson, *J. Am. Chem. Soc.* **2000**, 122, 5251–5257.
- [5] Spectroscopic data for **1**: ^1H NMR (300.1 MHz, $\text{C}_6\text{D}_5\text{N}$, 25 °C): δ = 6.03 (br. s, 4H; OH), 3.85 (t, J = 4.5 Hz, 8H; OCH_2), 2.49, 2.41 (br., 24H; NCH_2), 1.51 (q, J = 81.3 Hz, 4H; BH_4); ^{13}C NMR (75.4 MHz, CD_3CN , 25 °C): δ = 51.17 (NCH_2), 56.45 (NCH_2), 58.96 (OCH_2); ^{11}B NMR (96.23 MHz, CD_3CN , 25 °C, $\text{B}(\text{OCH}_3)_3$): δ = –54.21 (quintet, J = 81.4 Hz; BH_4); ^{11}B MAS NMR (128.33 MHz, 4013 Hz, 25 °C, $\text{B}(\text{OH})_3$): δ = –45.4 ($\Delta\nu_{1/2}$ = 2198 Hz); ^{23}Na MAS NMR (105.81 MHz, 6028 Hz, 25 °C, NaCl): δ = –12.3 ($\Delta\nu_{1/2}$ = 1440 Hz); IR (KBr): $\tilde{\nu}$ = 2298, 2225 cm^{-1} (B–H); elemental analysis calcd for $\text{C}_{16}\text{H}_{40}\text{N}_4\text{O}_4\text{BNa}$ (%): C 49.74, H 10.36, N 14.51; found: C 49.37, H 10.87, N 14.80; m.p. 180–181 °C.
- [6] A Siemens SMART CCD diffractometer with MoK_α radiation (λ = 0.71073) was used for the X-ray single-crystal analyses in this study. The structures were solved by direct methods and refined by full-matrix least-squares on F^2 using the SHELXTL package (Structure Analysis Program 5.1, Bruker AXS, Inc., Madison, WI, **1997**). Absorption corrections were applied using SADABS. All non-hydrogen atoms were refined anisotropically. Hydrogen atoms were located from the difference Fourier maps and refined isotropically. For the borohydride ions in **1**, however, only the hydridic hydrogen atoms involved in dihydrogen bonding could be found; the remaining H atoms from BH_4^- were calculated and placed in idealized positions. The observed O–H and B–H distances appear typically shorter than the ideal values of 0.96 and 1.21 Å, and were not corrected. Crystal data for **1**: crystal dimensions: $0.23 \times 0.18 \times 0.13$ mm; T = 173 K; orthorhombic, I_{222} ; a = 8.8240(3), b = 14.1344(6), c = 17.3533(7) Å, V = 2164.34(15) Å³, Z = 4; ρ_{calcd} = 1.186 g cm^{-3} ; $2\theta_{\text{max}}$ = 56.4°; 12 854 reflections collected, 2608 unique; 200 parameters; R_1 = 0.0487, wR_2 = 0.1221 for $I > 2\sigma(I)$; residual electron density: 0.488 e Å^{-3} . Crystal data for **3**: crystal dimensions: $0.47 \times 0.39 \times 0.29$ mm; T = 173 K; monoclinic, $P2_1/n$; a = 8.7696(1), b = 13.0571(1), c = 17.7724(2) Å, β = 96.36°, V = 2022.50(4) Å³, Z = 4; ρ_{calcd} = 1.301 g cm^{-3} ; $2\theta_{\text{max}}$ = 56.5°; 20 248 reflections collected, 4840 unique; 462 parameters; R_1 = 0.0352, wR_2 = 0.0921 for $I > 2\sigma(I)$; residual electron density: 0.261 e Å^{-3} . Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication nos. CCDC-143751 and CCDC-143752. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [7] a) S. Buoën, J. Dale, J. Krane, *J. Chem. Soc. Chem. Commun.* **1982**, 1172–1174; b) S. L. Whitbread, S. Politis, A. K. W. Stephens, J. B. Lucas, R. S. Dhilon, S. F. Lincoln, K. P. Wainwright, *J. Chem. Soc. Dalton Trans.* **1996**, 1379–1384.
- [8] ^{11}B MAS NMR (128.33 MHz, 4038 Hz, 25 °C, $\text{B}(\text{OH})_3$): δ = –8.0 ($\Delta\nu_{1/2}$ = 2462 Hz); ^{23}Na MAS NMR (105.81 MHz, 6001 Hz, 25 °C, NaCl): δ = –15.6 ($\Delta\nu_{1/2}$ = 3421 Hz).
- [9] a) T. Suzuki, T. Fukushima, Y. Yamashita, T. Miyashi, *J. Am. Chem. Soc.* **1994**, 116, 2793–2803; b) K. Tashiro, T. Kamae, M. Kobayashi, A. Matsumoto, K. Yokoi, S. Aoki, *Macromolecules* **1999**, 32, 2449–2454.
- [10] ^{11}B MAS NMR (128.33 MHz, 6000 Hz, 25 °C, $\text{B}(\text{OH})_3$): δ = –6.9 ($\Delta\nu_{1/2}$ = 1745 Hz); ^{23}Na MAS NMR (105.81 MHz, 6000 Hz, 25 °C, NaCl): δ = –8.9 ($\Delta\nu_{1/2}$ = 2704 Hz).
- [11] For this comparison, the following unit cell parameters derived from powder X-ray diffraction data at room temperature were used for **1**: a = 8.894(7), b = 14.214(13), c = 17.509(17) Å, V = 2214(3) Å³.
- [12] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, R. E. Stratmann, J. C. Burant, S. Dapprich, A. D. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Thomas, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. G. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle, J. A. Pople, Gaussian Inc., Pittsburgh PA, **1998**.
- [13] A. D. Becke, *J. Chem. Phys.* **1993**, 98, 5648.
- [14] ^{11}B MAS NMR (128.33 MHz, 6060 Hz, 25 °C, $\text{B}(\text{OH})_3$): δ = –7.5 ($\Delta\nu_{1/2}$ = 1880 Hz); ^{23}Na MAS NMR (105.81 MHz, 6058 Hz, 25 °C, NaCl): δ = –9.0 ($\Delta\nu_{1/2}$ = 2476 Hz).
- [15] A topochemical dihydrogen to covalent bonding transformation has been previously reported to occur in the solid-state conversion of cyclotrigallazane to nanocrystalline gallium nitride. However, initial loss of H_2 at 150 °C resulted in an amorphous GaN phase, and subsequent annealing at 600 °C was required to convert it into the final crystalline product. See: J.-W. Hwang, J. P. Campbell, J. Kozubowski, S. A. Hanson, J. F. Evans, W. L. Gladfelter, *Chem. Mater.* **1995**, 7, 517–525.

NMR and Theoretical Study of Acid Sites Formed by Adsorption of SO_3 onto Oxide Surfaces**

Jinhua Zhang, John B. Nicholas,* and James F. Haw*

The structure and function of acid sites formed by the treatment of oxide surfaces with sulfur compounds is a current problem in heterogeneous catalysis.^[1] While many studies have considered hydrogen sulfate surface complexes as the locus of solid acidity in such materials, other acidic surface sites have also been proposed, including those formed from adsorbed SO_3 . Haase and Sauer recently reported a periodic ab initio theoretical study of sulfuric acid adsorbed on low-index surfaces of ZrO_2 .^[2] One of the stable structures they found on $\text{ZrO}_2(001)$ was a tridentate-adsorbed SO_3 species. This conclusion was consistent with an experimental study of Babou and co-workers, who on the basis of infrared studies

[*] Prof. J. F. Haw

Loker Hydrocarbon Research Institute
and Department of Chemistry
University of Southern California
Los Angeles, CA 90089-1661 (USA)
Fax: (+1) 213-740-6679
E-mail: haw@methy.usc.edu

Dr. J. B. Nicholas

Environmental Molecular Sciences Laboratory
Pacific Northwest National Laboratory
Richland, WA 99352 (USA)
Fax: (+1) 509-375-6631
E-mail: john.nicholas@pnl.gov

J. Zhang

Department of Chemistry
Texas A&M University
Austin, TX (USA)

[**] This work was supported by the US NSF.

concluded that adsorbed SO_3 formed on sulfated zirconia prepared by using sulfuric acid as the sulfur source.^[3] Yamaguchi and co-workers found that generation of strong acid sites on Fe_2O_3 was independent of the sulfur source including sulfate, SO_2 , and SO_3 .^[4]

We treated various oxide materials with SO_3 and studied them using a combination of solid-state NMR spectroscopy and theoretical methods.^[5] Here we report results obtained for a material prepared from silica gel ($\text{SO}_3/\text{silica}$) and one prepared from zirconia ($\text{SO}_3/\text{zirconia}$). Figures 1a and 1b show the ^1H MAS NMR spectra of activated silica and $\text{SO}_3/\text{silica}$ following evacuation at 473 K, respectively. Activated




Figure 1. 359.7 MHz ^1H MAS NMR spectra and 50.1 MHz ^{13}C MAS NMR spectra characterizing activated silica and $\text{SO}_3/\text{silica}$: a) ^1H MAS spectrum of activated silica; b) ^1H MAS spectrum of $\text{SO}_3/\text{silica}$; c) ^{13}C MAS spectrum of acetone-2- ^{13}C on activated silica; d) ^{13}C MAS spectrum of acetone-2- ^{13}C on $\text{SO}_3/\text{silica}$. Asterisks (*) denote spinning side bands. All spectra were acquired at 298 K.

silica displays isolated silanol groups at $\delta = 1.9$, but $\text{SO}_3/\text{silica}$ has a new resonance signal at $\delta = 10.6$. The ^{13}C isotropic chemical shift of adsorbed acetone is a useful spectroscopic probe of relative acid strength.^[6] Figure 1c shows the ^{13}C MAS NMR spectrum of acetone-2- ^{13}C on activated silica; the ^{13}C shift at $\delta = 210$ indicates hydrogen bonding but negligible acidity. Figure 1d shows the ^{13}C MAS NMR spectrum of acetone-2- ^{13}C on $\text{SO}_3/\text{silica}$. The peak due to unreacted acetone is at $\delta = 217$; this shift indicates acidity, but of slightly lower strength than zeolite HZSM-5, which induces an acetone shift at $\delta = 223$. Figure 1d also shows small peaks at $\delta = 211$ and 189, due to mesityl oxide, which forms from the acid-catalyzed reaction of acetone.^[6]

We probed the nature of the acid sites formed by SO_3 treatment by measuring the ^{15}N MAS NMR spectra of ^{15}N -pyridine on both materials.^[7] The ^{15}N spectrum of pyridine on silica (Figure 2a) shows only hydrogen-bonded pyridine ($\delta = -88$) and a small amount of “free” pyridine ($\delta = -65$). Figure 2b shows that much of the pyridine on $\text{SO}_3/\text{silica}$ is protonated by Brønsted sites ($\delta = -174$). Figure 2c is the ^{15}N




Figure 2. 36.5 MHz ^{15}N CP/MAS NMR spectra of ^{15}N -pyridine on the four oxide materials (0.3 mmol g^{-1} on silica materials and 0.2 mmol g^{-1} on zirconia materials). a) Silica; b) $\text{SO}_3/\text{silica}$; c) zirconia; d) $\text{SO}_3/\text{zirconia}$. All spectra were acquired at 123 K. Asterisks (*) denotes spinning side bands.

spectrum of pyridine on zirconia; the pyridine is complexed to Lewis sites on this sample and resonates at $\delta = -101$. The thermal stability of $\text{SO}_3/\text{zirconia}$ is remarkable; much of the SO_3 initially adsorbed remains on the surface after evacuation to 773 K. Figure 2d shows that this material contains Brønsted ($\delta = -180$) as well as Lewis ($\delta = -106$) sites. The latter are presumably the adsorbed SO_3 species described by Hasse and Sauer.

We applied theoretical methods to model the Brønsted sites formed by treatment of silica with SO_3 . We used cluster models based on of β -cristabolite to approximate the catalysts and density functional theory to optimize the geometries of the reaction products. Initial calculations used a cluster of stoichiometry $\text{Si}_4\text{O}_{13}\text{H}_{10}$, fixing the terminal OH groups in crystallographic positions^[8] and optimizing the inner nine atoms of the cluster and the adsorbed SO_3 . Final energies were calculated by using a cluster to which an additional shell of atoms was added ($\text{Si}_{13}\text{O}_{13}\text{H}_{28}$). We considered both the interaction of SO_3 with the silanol lone pairs in a dative fashion, and as an insertion complex which creates an ester of hydrogen sulfate bound to the silica surface (Figure 3). The reaction of SO_3 with the surface silanol to form a dative complex is exothermic by $5.1 \text{ kcal mol}^{-1}$. Formation of the hydrogen sulfate ester is further exothermic by $5.0 \text{ kcal mol}^{-1}$. The theoretical deprotonation enthalpy of the ester is $294.4 \text{ kcal mol}^{-1}$, whereas it is $290.2 \text{ kcal mol}^{-1}$ for the dative complex. For comparison, the deprotonation enthalpy of the unactivated silanol group is $326.3 \text{ kcal mol}^{-1}$, in good agreement with the experimental value of $332.2 \pm 6.0 \text{ kcal mol}^{-1}$.^[9] The ester is predicted to be a much stronger acid than the unactivated surface silanols. In fact, the calculated acidity of the ester is comparable to that of a similar model of zeolite H-ZSM5 at the same level of theory ($294.1 \text{ kcal mol}^{-1}$). These




Figure 3. Optimized B3LYP/DZVP2 geometry of the hydrogen sulfate ester formed by reaction of SO_3 on β -cristabolite.

theoretical values compare to experimental deprotonation enthalpies for ZSM-5 of 291–300 kcal mol⁻¹.^[10]

In summary, NMR experiments confirm that adsorption of SO_3 onto oxide surfaces introduces acid sites. On SO_3/ZrO_2 , both Lewis and Brønsted sites are present, and theoretical modeling, in agreement with NMR studies, supports the formation of Brønsted sites on $\text{SO}_3/\text{silica}$.

Experimental Section

Electronic grade silicon(IV) oxide (Puratrem, 99.999 %) was obtained from Strem. This was evacuated at 873 K, cooled to 298 K, and then contacted with SO_3 (0.8 mmol g⁻¹). $\text{SO}_3/\text{silica}$ samples were evacuated at 473 K prior to investigation. Zirconia was prepared by the hydrolysis of zirconyl chloride octahydrate (98 %, Aldrich), followed by washing and drying the gel. All zirconia samples were first calcined at 773 K for two hours in flowing air. SO_3 (1.43 mmol g⁻¹) was adsorbed at 298 K and all samples of $\text{SO}_3/\text{zirconia}$ were further evacuated at 773 K before study. ¹H and ¹⁵N MAS NMR spectra were measured at 9.4 Tesla on a Chemagnetics CMX-360 spectrometer and ¹³C MAS NMR spectra were measured at 4.7 Tesla on a home-built spectrometer.

Theoretical calculations were performed by using Gaussian 94^[11] and the B3LYP^[12] exchange-correlation functional and the DZVP2^[13] basis set. The terminal O and H atoms were held fixed in crystallographic positions, with the OH bond lengths set to 0.9675 Å. Frequency calculations on $\text{H}_3\text{SiOHSO}_3$ and $\text{H}_3\text{SiOSO}_3\text{H}$ were used to estimate the zero-point and thermal energy contributions for the determination of the deprotonation enthalpies of the larger models.

Received: September 27, 1999
Revised: May 29, 2000 [Z14070]

- [1] a) X. Song, A. Sayari, *Catal. Rev. Sci. Eng.* **1996**, *38*, 229–412; b) A. Corma, *Chem. Rev.* **1995**, *95*, 559–614; c) V. Adeeva, J. W. Dehaan, J. Janchen, G. D. Lei, V. Schunemann, L. J. M. van Devan, W. M. H. Sachtler, R. A. van Santen, *J. Catal.* **1995**, *151*, 364–372; d) T. Reimer, D. Spielbauer, M. Hunger, G. A. H. Mekheimer, H. Knözinger, *Chem. Comm.* **1994**, 1181–1182.
- [2] F. Haase, J. Sauer, *J. Am. Chem. Soc.* **1998**, *120*, 13 503–13 512.
- [3] F. Babou, G. Coudurier, J. C. Vedrine, *J. Catal.* **1995**, *152*, 341–349.
- [4] T. Yamaguchi, T. Jin, K. Tanabe, *J. Phys. Chem.* **1986**, *90*, 3148–3152.
- [5] a) J. F. Haw, J. B. Nicholas, T. Xu, L. W. Beck, D. B. Ferguson, *Acc. Chem. Res.* **1996**, *29*, 259–267; b) T. Xu, N. Kob, R. S. Drago, J. B. Nicholas, J. F. Haw, *J. Am. Chem. Soc.* **1997**, *119*, 12 231–12 239.
- [6] T. Xu, E. J. Munson, J. F. Haw, *J. Am. Chem. Soc.* **1994**, *116*, 1962–1972.
- [7] J. F. Haw, I.-S. Chuang, B. L. Hawkins, G. E. Maciel, *J. Am. Chem. Soc.* **1983**, *105*, 7206–7207.
- [8] R. W. G. Wyckoff, *Crystal Structures Vol. IV*, Wiley, New York, **1982**, p. 318.
- [9] E. A. Paukshtis, E. N. Yurchenko, *Usp. Khim.* **1983**, *52*, 426.
- [10] H. V. Brand, L. A. Curtiss, L. E. Iton, *J. Phys. Chem.* **1993**, *97*, 12 773–12 782.
- [11] Gaussian 94, Revision B.2: M. J. Frisch, G. W. Trucks, H. B. Schlegel, P. M. W. Gill, B. G. Johnson, M. A. Robb, J. R. Cheeseman, T. Keith, G. A. Petersson, J. A. Montgomery, K. Raghavachari, M. A. Al-Laham, V. G. Zakrzewski, J. V. Ortiz, J. B. Foresman, J. Cioslowski, B. B. Stefanov, A. Nanayakkara, M. Challacombe, C. Y. Peng, P. Y. Ayala, W. Chen, M. W. Wong, J. L. Andres, E. S. Replogle, R. Gomperts, R. L. Martin, D. J. Fox, J. S. Binkley, D. J. Defrees, J. Baker, J. P. Stewart, M. Head-Gordon, C. Gonzalez, J. A. Pople, Gaussian Inc., Pittsburgh, PA, **1994**.
- [12] A. D. Becke, *J. Chem. Phys.* **1993**, *98*, 5648–5652.
- [13] N. Godbout, D. R. Salahub, J. Andzelm, E. Wimmer, *Can. J. Chem.* **1992**, *70*, 560–571.

Syntheses, Structures, and Magnetic Properties of Two Gadolinium(III)–Copper(II) Coordination Polymers by a Hydrothermal Reaction**

Yucang Liang, Rong Cao,* Weiping Su,
Maohun Hong,* and Wenjian Zhang

Studies on the syntheses, structures, and properties of lanthanide–transition metal complexes are currently of great interest because they are good models to investigate the nature of the magnetic exchange interactions between 3d and 4f metal ions in magnetic materials that contain rare earth metals.^[1–5] Many such studies focused on discrete complexes, which were synthesized from conventional, self-assembly

[*] R. Cao, Prof. M. Hong, Y. Liang, Dr. W. Su, W. Zhang
State Key Laboratory of Structural Chemistry
Fujian Institute of Research on the Structure of Matter
Chinese Academy of Sciences
Fuzhou, Fujian 350002 (P. R. China)
Fax: (+86) 591-371-4946
E-mail: rcao@ms.fjirsm.ac.cn, hmc@ms.fjirsm.ac.cn

[**] This work is supported by the NNSF of China, the NSF of Fujian Province, and the Key Project from the CAS.

Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.