

purified by flash chromatography on silica gel (Et₂O/hexanes 50/50 → 100/0) to give the desired product (205 mg, 0.871 mmol, 88% yield) as a pale yellow oil. R_f = 0.27 (Et₂O/hexanes 90/10); IR (neat): $\tilde{\nu}$ = 3506.1 cm⁻¹ (br.), 3115.9, 2921.6, 2710.9, 1692.9, 1648.3, 1605.4, 1512.0, 1404.1, 1345.1, 1237.9, 1106.8, 1063.1, 1019.7, 850.8; ¹H NMR (400 MHz, CDCl₃): δ = 9.43 (s, 1H; H-1), AB signals (4H, δ = 8.25, 7.58, J_{AB} = 8.7 Hz; aromatic H), 6.56 (tq, 1H, J = 7.2, 1.2 Hz; H-3), 5.07 (dd, 1H, J = 6.2 Hz; H-5), 2.89–2.77 (m, 2H; H-4), 2.20 (br. s, 1H; OH), 1.71 (s, 3H; CH₃); ¹³C NMR (100 MHz, CDCl₃): δ = 195.17, 150.94, 148.99, 147.25, 141.35, 126.46, 123.73, 71.90, 38.37, 9.31; MS: m/z (%) = 218.22 (3.5) [M^+ – OH], 152.15 (57.5), 84.13 (100.0); HR-MS: calcd for C₁₂H₁₄NO₄ = 236.0923, found = 236.0921; molecular formula: C₁₂H₁₃NO₄ (235.23 g mol⁻¹).

Received: June 5, 2000 [Z15226]

Total Synthesis of (+)-Eurylene and (+)-14-Deacetylene**

Yoshiki Morimoto,* Koji Muragaki, Toshiyuki Iwai, Yukio Morishita, and Takamasa Kinoshita

Recently biologically active and structurally novel triterpene polyethers, natural products which are thought to be biogenetically derived from squalene, have been isolated from both marine and terrestrial plants. Among the polyethers are cytotoxic eurylene (**1**) and 14-deacetylene (**2**; see Scheme 1), isolated from the wood of *Eurycoma longifolia* by Itokawa et al.^[1] The stereostructures and conformations of **1** and **2** have been elucidated by X-ray crystallographic analysis and spectroscopic methods.^[2] The mechanism of action for the cytotoxic activities of these natural polyethers, however, remains to be clarified, because these molecules are available only in restricted amounts from natural sources. Therefore, the development of an efficient synthesis was desired for these polyethers. The novel structures, the cytotoxic activities, and the conformation–activity relationships^[2] of **1** and **2** have attracted the attention of many synthetic organic chemists; however, the total synthesis of **2** with potent cytotoxic activity has never been accomplished.^[3] Here we report the efficient and stereoselective total synthesis of (+)-eurylene (**1**) and (+)-14-deacetyl eurylene (**2**), featuring monool- and diol-differentiated chemoselective oxidative cyclizations promoted by rhenium(VII) and chromium(VI) oxo species, respectively.

Our retrosynthetic analysis of **1** and **2** is depicted in Scheme 1. The key events for the total synthesis are the stereoselective construction of the *trans* and *cis* tetrahydrofuran (THF) rings and the differentiation of the 14-hydroxy group. To solve this problem, we focused on the hydroxy-directed *syn* oxidative cyclization of acyclic bishomoallylic alcohols promoted by rhenium(VII)^[4] and chromium(VI) oxides.^[5] Thus, the *trans* THF ring will be constructed by applying our Re^{VII} protocol^[6] to the bishomoallylic monool moiety in triol **3**, while the *cis* THF ring is constructed by the Cr^{VI}-induced *cis*-selective cyclization of the bishomoallylic vicinal diol. The pseudo-*meso* triol **3** will, in turn, be derived


- [1] a) J. Colonge, G. Descottes, Y. Bahurel, *Bull. Soc. Chim. Fr.* **1965**, 619; b) P. Miginiac, C. Prévost, *Bull. Soc. Chim. Fr.* **1966**, 704; c) R. J. Anderson, *J. Am. Chem. Soc.* **1970**, 92, 4978; d) R. W. Herr, C. R. Johnson, *J. Am. Chem. Soc.* **1970**, 92, 4979; e) M. Tamura, G. Suzukamo, *Tetrahedron Lett.* **1981**, 22, 577; f) S. K. Taylor, *Tetrahedron* **2000**, 56, 1149, and references therein.
- [2] a) For the Lewis acid catalyzed rearrangement of oxiranes, see: B. J. Rickborn in *Comprehensive Organic Synthesis*, Vol. 3 (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, pp. 733–775, and references therein; b) P. Wipf, W. Xu, *J. Org. Chem.* **1993**, 58, 825; a stoichiometric amount of the organozirconocene was needed.
- [3] a) W. E. Bachmann, J. P. Horwitz, R. J. Warzynski, *J. Am. Chem. Soc.* **1953**, 75, 3268; b) the rearrangement under various conditions was reported: D. Serramedan, F. Marc, M. Perey, C. Filliatre, P. Chabardès, B. Delmond, *Tetrahedron Lett.* **1992**, 33, 4457; c) F. Marc, B. Soulet, D. Serramedan, B. Delmond, *Tetrahedron* **1994**, 50, 3381; d) C. Chapuis, R. Brauchli, *Helv. Chim. Acta.* **1993**, 76, 2070.
- [4] For an alternative to the use of β,γ -unsaturated aldehydes, see: G. Hughes, M. Lautens, C. Wen, *Org. Lett.* **2000**, 2, 107.
- [5] R. A. Batey, A. N. Thadani, D. V. Smill, *Tetrahedron Lett.* **1999**, 40, 4289.
- [6] All new products have been fully characterized by ¹H and ¹³C NMR spectroscopy, IR spectroscopy, mass spectrometry, and high-resolution mass spectrometry.
- [7] J. M. Coxon, C.-E. Lim, *Aus. J. Chem.* **1977**, 30, 1137.
- [8] Compounds **7** and **14** were already synthesized by another strategy: S. Saito, M. Shiozawa, M. Ito, H. Yamamoto, *J. Am. Chem. Soc.* **1998**, 120, 813.
- [9] Those conditions favor the formation of the dimerization product.
- [10] M. Lautens, S. Raepel, S. G. Ouellet, full paper in preparation.
- [11] M. T. Crimmins, S. J. Kirincich, A. J. Wells, A. L. Choy, *Syn. Commun.* **1998**, 28, 3675, and references therein.
- [12] Nucleophiles such as allyltributyltin and tetraallyltin also add to afford the expected bishomoallylic alcohols.
- [13] I. Paterson, J. D. Smith, R. A. Ward, *Tetrahedron* **1995**, 51, 9413.
- [14] a) T. Mukaiyama, K. Banno, K. Narasaka, *J. Am. Chem. Soc.* **1974**, 96, 7503; b) T. Mukaiyama, A. Ishida, *Chem. Lett.* **1975**, 319; T. Mukaiyama, A. Ishida, *Chem. Lett.* **1975**, 1201; c) A. Ishida, T. Mukaiyama, *Chem. Lett.* **1977**, 467; d) T. Mukaiyama, *Tetrahedron* **1999**, 55, 8609; for a review on the vinylogous aldol reaction, see: G. Casiraghi, F. Zanardi, G. Appendino, G. Rassu, *Chem. Rev.* **2000**, 100, 1929.

[*] Prof. Dr. Y. Morimoto, K. Muragaki, T. Iwai, Y. Morishita, Prof. Dr. T. Kinoshita
Department of Chemistry, Graduate School of Science
Osaka City University
Sumiyoshiku, Osaka 558-8585 (Japan)
Fax: (+81) 6-6605-2522
E-mail: morimoto@sci.osaka-cu.ac.jp

[**] We thank K. Ujihara (Sumitomo Chemical Co.) for generously supplying an authentic sample and a copy of the ¹H NMR spectrum of synthetic eurylene. We are grateful to Prof. H. Iio and Dr. Y. Usuki (Osaka City University) for the HPLC purification. T.I. thanks the Japan Society for the Promotion of Science for providing a JSPS Research Fellowship for Young Scientists. This research was supported by the Naito Foundation, the Fujisawa Pharmaceutical Award in Synthetic Organic Chemistry, Japan, the SUNBOR GRANT from the Suntory Institute for Bioorganic Research, and a Grant-in Aid for Encouragement of JSPS Research Fellows from the Japan Society for the Promotion of Science.




Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.



Scheme 1. Retrosynthetic analysis of **1** and **2**. Pv = pivaloyl; MPM = 4-methoxyphenylmethyl.

from the appropriately protected diepoxide **4** by extending the side chains with two neryl units in a bidirectional manner. It was envisaged that the diepoxide **4** can be prepared by Sharpless asymmetric epoxidation of the readily available diol **5** and subsequent regioselective ring-opening reactions.


The stereoselective preparation of the diepoxide **4** began with monoprotection of the known diol **5**^[7] as a *tert*-butyldimethylsilyl (TBS) ether ($\rightarrow$ **6**; Scheme 2). Sharpless asymmetric epoxidation^[8] of the allylic alcohol **6**^[9] using (–)-diethyl D-tartrate (D-(–)-DET) afforded the epoxy alcohol **7** in high optical purity.^[10] The pivalate group was introduced regioselectively into **7** by a titanium-assisted epoxide-opening



Scheme 2. a) TBSCl, imidazole, CH_2Cl_2 , RT, 1 h, 55%; b) TBHP, $\text{Ti}(\text{O}i\text{Pr})_4$, D-(–)-DET, MS 4 Å, CH_2Cl_2 , -20°C , 2 h, 83% (98% ee); c) $\text{Ti}(\text{O}i\text{Pr})_4$, PvOH, benzene, 0°C , 2 h, 83%; d) 2,2-dimethoxypropane, camphorsulfonic acid, CH_2Cl_2 , 0°C , 2 h, 88%; e) Bu_4NF , THF, RT, 3 h, 99%; f) TBHP, $\text{Ti}(\text{O}i\text{Pr})_4$, L-(+)-DET, MS 4 Å, CH_2Cl_2 , -20°C , 3 h, 89%; g) $\text{Ti}(\text{OMPM})_4$, MPMOH, benzene, 60°C , 12 h; h) $\text{AcOH}/\text{H}_2\text{O}$ (4/1), RT, 5 h; i) MsCl , Py, CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{RT}$, 5 h; j) K_2CO_3 , MeOH, RT, 1 h, 41% (4 steps). Ms = methanesulfonyl, Py = pyridine, TBHP = *tert*-butyl hydroperoxide.

reaction^[11] to yield 1,2-diol **8** as a single diastereomer; subsequent acetone protection and desilylation furnished the allylic alcohol **10** in good overall yield. The asymmetric epoxidation of **10** using L-(+)-DET provided the epoxy alcohol **11**, which was subjected to a $\text{Ti}(\text{OMPM})_4$ -mediated epoxide-opening reaction^[11, 12] with *p*-anis alcohol to produce the desired 1,2-diol **12** and a 1,3-diol derivative in a ratio of approximately 3:1. Deprotection of the acetonide in 1,2-diol **12**, mesylation of both primary hydroxy groups in the resultant tetraol, and subsequent basic treatment of the dimesylate finally gave the requisite diepoxide **4** in 41% yield (based on four steps from **11**).

The bidirectional chain extension proceeded smoothly by alkylation of **13**, the lithio derivative of neryl phenyl sulfide,^[6b] with the diepoxide **4** in the presence of *N,N,N',N'*-tetramethylethylenediamine (TMEDA), yielding the bisulfide **14** as a mixture of diastereomers (Scheme 3).^[13] Compound **14** was



Scheme 3. a) Compound **13**, TMEDA, THF, -78°C , 30 min, then 0°C , 2 h; b) Na, THF/*i*PrOH (2/1), reflux, 15 h, 86% (2 steps); c) Ac_2O , Py, RT, 12 h, 91%; d) DDQ, MS 4 Å, CH_2Cl_2 , 0°C , 2 h, 68%; e) $\text{AcOH}/\text{H}_2\text{O}$ (4/1), RT, 16 h, 89%; f) $[(\text{CF}_3\text{CO}_2)\text{ReO}_3 \cdot 2\text{CH}_3\text{CN}]$, TFAA, $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{CN}$ (9/1), -40°C , 1.5 h, 84%; g) PCC, CH_2Cl_2 , RT, 30 min, 47%; h) Ac_2O , Py, RT, 40 h, 80%.

desulfurized under Bouvault–Blanc conditions^[6b, 14] to provide the triol **15**, and selective acetylation of the *sec*-alcohol afforded the diol **16** in good overall yield. The MPM protecting group in **16** was converted into the 4-methoxybenzylidene acetal through interaction of the neighboring hydroxy groups with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ).^[15] Subsequent acidic hydrolysis furnished the key substrate **3** required for the oxidative cyclizations.

The optimal conditions were determined, and the treatment of triol **3** with eight equivalents of the oxorhenium(VII) complex $[(\text{CF}_3\text{CO}_2)\text{ReO}_3 \cdot 2\text{CH}_3\text{CN}]$ ^[16] and ten equivalents of trifluoroacetic anhydride (TFAA) in a mixed solvent system ($\text{CH}_2\text{Cl}_2/\text{CH}_3\text{CN}$ 9/1)^[6] at -40°C for 1.5 h diastereoselectively

gave the expected *trans* THF product **17** in 84% yield; the right-hand part of the molecule was intact. On the other hand, when **17** was treated with a stoichiometric amount of the oxochromium(vi) complex pyridinium chlorochromate (PCC),^[17] in CH₂Cl₂ at room temperature for 30 min,^[5] the bishomoallylic vicinal diol moiety was oxidatively cyclized, probably via a chelated dialkoxychromium intermediate,^[5a, 18] with complete *cis* diastereoselectivity to produce (+)-14-deacetylerylene (**2**).^[19] The spectral characteristics (¹H and ¹³C NMR, IR, MS, and HRMS) of synthetic **2** were identical to those reported for the natural product.^[1b] Finally, selective acetylation of the 14-hydroxy group in **2** afforded another objective, (+)-eurylene (**1**),^[19] whose spectral data were also consistent with the ¹H NMR spectrum of an authentic sample^[3a] and with the data reported for the natural product.^[1]

In conclusion, we have accomplished the first total synthesis of (+)-14-deacetylerylene (**2**), featuring chemoselective THF ring formations stereocontrolled by *syn*-oxidative cyclizations of the bishomoallylic monool and diol induced by oxorhenium(vii) and -chromium(vi) species, respectively. The reason for the intriguing relationship^[2] between the conformations and cytotoxicities of **1** and **2** is under investigation.

Received: May 18, 2000

Revised: August 21, 2000 [Z15142]

- [1] a) H. Itokawa, E. Kishi, H. Morita, K. Takeya, Y. Iitaka, *Tetrahedron Lett.* **1991**, 32, 1803–1804; b) H. Morita, E. Kishi, K. Takeya, H. Itokawa, Y. Iitaka, *Phytochemistry* **1993**, 34, 765–771.
- [2] Itokawa et al. have also proposed a very interesting relationship between the conformations and cytotoxicities of **1** and **2**: Compound **2**, which adopts the folded conformation, exhibits prominent cytotoxic activity on KB cells (IC₅₀ = 0.52 µg mL⁻¹), whereas **1**, possessing an extended conformation, does not (IC₅₀ > 100 µg mL⁻¹).
- [3] For the previous total synthesis of eurylene (**1**), see: a) K. Ujihara, H. Shirahama, *Tetrahedron Lett.* **1996**, 37, 2039–2042; for synthetic approaches toward **1**, see: b) M. K. Gurjar, U. K. Saha, *Tetrahedron Lett.* **1993**, 34, 1833–1836; c) S. Yoshio, H. Minami, H. Mori, Y. Oshita, T. Tabata, Y. Fukuyama, M. Kodama, H. Takayanagi, Y. Morinaka, Y. Fukazawa, *Tennen Yuki Kagobutsu Toronkai Koen Yoshishu* **1993**, 35, 471–478 [*Chem. Abstr.* **1994**, 121, 256071v]; d) G. A. Molander, S. Swallow, *J. Org. Chem.* **1994**, 59, 7148–7151.
- [4] For reviews on rhenium(vii) chemistry, see a) W. A. Herrmann, F. E. Kühn, *Acc. Chem. Res.* **1997**, 30, 169–180; b) C. C. Romão, F. E. Kühn, W. A. Herrmann, *Chem. Rev.* **1997**, 97, 3197–3246. For Re^{VII}-induced oxidative cyclizations of 5-hydroxyalkenes and their application to natural product syntheses, see c) R. M. Kennedy, S. Tang, *Tetrahedron Lett.* **1992**, 33, 3729–3732; d) S. C. Sinha, A. Sinha-Bagchi, E. Keinan, *J. Am. Chem. Soc.* **1995**, 117, 1447–1448; e) S. C. Sinha, A. Sinha-Bagchi, A. Yazbak, E. Keinan, *Tetrahedron Lett.* **1995**, 36, 9257–9260; f) E. Keinan, A. Sinha, A. Yazbak, S. C. Sinha, S. C. Sinha, *Pure Appl. Chem.* **1997**, 69, 423–430; g) T. B. Towne, F. E. McDonald, *J. Am. Chem. Soc.* **1997**, 119, 6022–6028; h) S. C. Sinha, A. Sinha, S. C. Sinha, E. Keinan, *J. Am. Chem. Soc.* **1997**, 119, 12014–12015; i) S. C. Sinha, A. Sinha, S. C. Sinha, E. Keinan, *J. Am. Chem. Soc.* **1998**, 120, 4017–4018; j) S. C. Sinha, E. Keinan, S. C. Sinha, *J. Am. Chem. Soc.* **1998**, 120, 9076–9077.
- [5] For Cr^{VI}-induced oxidative cyclizations of 5,6-dihydroxyalkenes and their application to natural product syntheses, see: a) D. M. Walba, G. S. Stoudt, *Tetrahedron Lett.* **1982**, 23, 727–730; b) D. M. Walba, G. S. Stoudt, *J. Org. Chem.* **1983**, 48, 5404–5406; c) E. J. Corey, D.-C. Ha, *Tetrahedron Lett.* **1988**, 29, 3171–3174; d) F. E. McDonald, C. C. Schultz, *Tetrahedron* **1997**, 53, 16435–16448; e) F. E. McDonald, T. B. Towne, C. C. Schultz, *Pure Appl. Chem.* **1998**, 70, 355–358.
- [6] a) Y. Morimoto, T. Iwai, *J. Am. Chem. Soc.* **1998**, 120, 1633–1634; b) Y. Morimoto, T. Iwai, T. Kinoshita, *J. Am. Chem. Soc.* **1999**, 121, 6792–6797.
- [7] T. Lindel, B. Franck, *Tetrahedron Lett.* **1995**, 36, 9465–9468.
- [8] Y. Gao, R. M. Hanson, J. M. Klunder, S. Y. Ko, H. Masamune, K. B. Sharpless, *J. Am. Chem. Soc.* **1987**, 109, 5765–5780.
- [9] All new compounds reported here were satisfactorily characterized based on their ¹H and ¹³C NMR, IR, MS, and HRMS spectra (see the Supporting Information).
- [10] The optical purity of **7** was determined to be 98% *ee* by derivatization of **7** to a Mosher ester with (R)-MTPACl (MTPA = *α*-methoxy-*α*-(trifluoromethyl)phenylacetic acid) and integration of the signals in the ¹H NMR spectrum (J. A. Dale, D. L. Dull, H. S. Mosher, *J. Org. Chem.* **1969**, 34, 2543–2549).
- [11] M. Caron, K. B. Sharpless, *J. Org. Chem.* **1985**, 50, 1557–1560.
- [12] M. Oikawa, T. Ueno, H. Oikawa, A. Ichihara, *J. Org. Chem.* **1995**, 60, 5048–5068.
- [13] During this alkylation, the pivaloyl group was removed by an excess of **13**.
- [14] M. Hashimoto, H. Harigaya, M. Yanagiya, H. Shirahama, *J. Org. Chem.* **1991**, 56, 2299–2311.
- [15] Y. Oikawa, T. Yoshioka, O. Yonemitsu, *Tetrahedron Lett.* **1982**, 23, 885–888 and 889–892.
- [16] a) W. A. Herrmann, W. R. Thiel, F. E. Kühn, R. W. Fischer, M. Kleine, E. Herdtweck, W. Scherer, J. Mink, *Inorg. Chem.* **1993**, 32, 5188–5194; b) P. Edwards, G. Wilkinson, *J. Chem. Soc. Dalton Trans.* **1984**, 2695–2702.
- [17] E. J. Corey, J. W. Suggs, *Tetrahedron Lett.* **1975**, 2467–2470.
- [18] It is postulated that, in this case, the formation of the chelated dialkoxychromium intermediate is favored over that of the monoalkoxychromium counterpart of the left-hand moiety due to use of one equivalent of PCC. The material balances other than **2** in this reaction were the oxidatively cleaved products of the vicinal diol moiety, the corresponding aldehyde (33%), and nerylaceton (40%).
- [19] **1**: M.p. 145–147 °C, [α]_D²⁵ = +4.86 (*c* = 0.275 in CHCl₃); ref. [1] m.p. 146–148 °C, [α]_D = +4.32 (*c* = 0.14 in CHCl₃). **2**: M.p. 64.3–67.4 °C, [α]_D²⁵ = +6.02 (*c* = 0.600 in CHCl₃); ref. [1b] m.p. 63–65 °C, [α]_D = +6 (*c* = 1.03 in CHCl₃).

Cuprophilicity: Spectroscopic and Structural Evidence for Cu–Cu Bonding Interactions in Luminescent Dinuclear Copper(II) Complexes with Bridging Diphosphane Ligands*

Chi-Ming Che,* Zhong Mao, Vincent M. Miskowski, Man-Chung Tse, Chi-Keung Chan, Kung-Kai Cheung, David Lee Phillips,* and King-Hung Leung

Copper(I)–copper(I) bonding interactions have widely been invoked to be a driving force for the self-assembly of copper(I) aggregates and to play an important role with regard to the photoluminescence of polynuclear copper(I) complexes con-

[*] Prof. Dr. C.-M. Che, Dr. D. L. Phillips, Z. Mao, Dr. V. M. Miskowski, Dr. M.-C. Tse, Dr. C.-K. Chan, Dr. K.-K. Cheung, Dr. K.-H. Leung
Department of Chemistry, The University of Hong Kong
Pokfulam Road, Hong Kong SAR (P.R. China)
Fax: (+852) 2857-1586
E-mail: cmche@hku.hk
phillips@hku.hk

[**] We are grateful for financial support from The University of Hong Kong, the Hong Kong University Foundation, and the Research Grants Council of Hong Kong SAR, P.R. China [HKU 7298/99P and HKU 974/94P].