

(202.3 MHz, CDCl₃): δ = 116.96; ¹H NMR (500 MHz, CDCl₃): δ = 8.48 (br, 1H), 7.54 (d, 1H, $J_{4,5}$ = 7.5 Hz), 7.51 (d, 1H, $J_{6,7}$ = 7.5 Hz), 7.45 (s, 1H), 7.19–7.35 (m, 7H), 6.12 (m, 1H), 4.90 (m, 1H), 4.36 (m, 1H), 3.45–4.00 (m, 8H), 3.50 (m, 3H), 3.26 (m, 1H), 3.13 (m, 1H), 2.96 (m, 1H), 2.23 (m, 1H), 2.01–2.06 (m, 1H), 1.82–1.95 (m, 7H), 0.91 (s, 9H), 0.08 (2s); HRMS (FAB): calcd for C₄₀H₅₃N₅O₇PSi: 774.3452; found: 774.3455.

15: DBU (37 μ L, 0.25 mmol) was added to a stirred solution of **12** (38.6 mg, 0.05 mmol) and 3'-O-TBDMS-thymidine (35.6 mg, 0.1 mmol) in dry THF (2 mL) at room temperature. After 20 min, the ³¹P NMR spectrum of the reaction mixture showed a single signal at around δ = 140. The mixture was then loaded onto a short column to filter off DBU. After evaporation of the solvent, the residue was dissolved in dry CH₂Cl₂, and Beaucage's reagent (10 mg, 0.05 mmol) was added at room temperature. After stirring for 5 min, ³¹P NMR spectroscopy revealed a single peak at around δ = 68. The solvent was removed, the residue was dissolved in THF, and TBAF was added. After 2 h, the solvent was removed in vacuo. CH₂Cl₂ was added to the residue, and the extract was washed with water and brine and dried over magnesium sulfate. Purification by column chromatography afforded **15** as a white solid (28 mg, 61 %).

16 and 17: Phosphorothioate **15** (28 mg, 0.026 mmol) was dissolved in concentrated ammonia/ethanol (3/1, 5 mL). The mixture was heated at 55 °C for 16 h. The ³¹P NMR signal shifted quantitatively from δ = 68 to 56. The reaction mixture was extracted with chloroform several times. The combined organic extracts were washed with brine and dried over magnesium sulfate. The solvent was removed to provide **17** (9.2 mg, 91 %) as a light yellow solid. The aqueous crude product was purified by flash chromatography to give **16** (15 mg, 89 %).


17: ¹H NMR (500 MHz, CDCl₃): δ = 7.41 (d, 1H, J = 8.0 Hz), 7.23–7.35 (m, 6H), 7.13 (m, 1H), 7.02 (m, 1H), 4.71 (br, 1H), 4.04–4.14 (m, 2H), 3.94 (m, 2H), 3.68–3.86 (m, 2H), 3.11–3.23 (m, 4H), 2.97 (m, 1H), 2.82 (m, 1H), 1.55–1.82 (m, 5H); HRMS (FAB): calcd for C₂₄H₂₉N₄O: 389.2341 [(M+1)⁺]; found: 389.2343.

Received: July 17, 2000 [Z15468]

Early Transition Metal α -Diazoalkane Complexes**

Victorio Cadierno, Maria Zablocka, Bruno Donnadieu, Alain Igau, and Jean-Pierre Majoral*

The reaction of diazoalkanes with transition metals results either in coordination through the nitrogen atoms or in formation of a carbon–metal bond. Many N-bonded substituted transition metal diazoalkanes have been studied,^[1] but only four series of C-bonded late transition metal compounds have been isolated.^[2] α -Metalated diazoalkanes L_nM–C(N₂)R containing an early transition metal have not yet been reported. Treatment of zirconocene Zr^{IV} complexes with diazoalkanes results in insertion into Zr–C,^[3] Zr–H,^[3c] Zr–P,^[4] and zirconium–metal bonds^[5] to give the corresponding hydrazone ligands **I** [Eq. (1)].



We have shown that 1-aza-zirconacyclopentene complexes **1a** and **1b**^[6] (see Scheme 1) contain a strongly electrophilic metal center (mostly because of the inductive electron-withdrawing properties of the σ -bonded nitrogen atom) and a rather nucleophilic imido group that does not share its lone-pair electron density with the adjacent metal center because of the strain in the metallacycle.^[7] Indeed phosphanes **1a** and **1b** activate C–H bonds in relatively acidic carbonic acids, such as acetylenes and methylene compounds.^[8] Although electrophilic, metal-mediated, aliphatic C–H bond cleavage is precedented,^[9] its application to the functionalization of complex substrates is very rare. The present report deals with a novel electrophilic sp²-C–H bond activation and its application for the preparation of α -diazomethylzirconium complexes, which are the first C-metalated diazoalkanes L_nMC(N₂)R with an early transition metal.

Treatment of **1a** and **1b** with ethyl diazoacetate (**2**) in THF at room temperature gave the light yellow, thermally stable solids **4a** and **4b** (Scheme 1) in excellent yields (>88 %). The ³¹P NMR spectra of **4a** and **4b** display a singlet at δ = 49.4 and 58.5, respectively. The total disappearance of the sp²-CH


[*] Dr. J.-P. Majoral, Dr. V. Cadierno, Dr. B. Donnadieu, Dr. A. Igau
Laboratoire de Chimie de Coordination, CNRS
205 route de Narbonne, 31077 Toulouse Cedex (France)
Fax: (+33) 05-61-55-30-03
E-mail: majoral@lcc.toul.lcc-toulouse.fr
Dr. M. Zablocka
Polish Academy of Sciences
Sienkiewicza 112, 90-363 Lodz (Poland)

[**] This work was supported by the CNRS, France and KBN (Poland, Grant No 3T09A03619). V.C. thanks the Spanish Ministerio de Educacion y Cultura for postdoctoral fellowship.



Supporting information for this article is available on the WWW under <http://www.wiley-vch.de/home/angewandte/> or from the author.

- [1] E. Uhlmann, A. Peyman, *Chem. Rev.* **1990**, 90, 543.
- [2] See webpage at <http://www.isip.com>.
- [3] W. J. Stec, A. Grajkowski, A. Kobylanska, B. Karwowski, M. Koziolkiewicz, K. Misiura, A. Okruszek, A. Wilk, P. Guga, M. Boczkowska, *J. Am. Chem. Soc.* **1995**, 117, 12019.
- [4] A. Wilk, A. Grajkowski, L. R. Phillips, S. L. Beaucage, *J. Am. Chem. Soc.* **2000**, 122, 2149.
- [5] a) Z. Xin, G. Just, *Tetrahedron Lett.* **1996**, 37, 969; b) Y. Jin, G. Biancotto, G. Just, *Tetrahedron Lett.* **1996**, 37, 973; c) E. Marsault, G. Just, *Tetrahedron Lett.* **1996**, 37, 977; d) E. Marsault, G. Just, *Tetrahedron* **1997**, 53, 16945; e) J. Wang, G. Just, *Tetrahedron Lett.* **1997**, 38, 705; f) J. Wang, G. Just, *Tetrahedron Lett.* **1997**, 38, 3797; g) J. Wang, G. Just, A. P. Guzaev, M. Manoharan, *J. Org. Chem.* **1999**, 24, 2595; h) Y. Jin, G. Just, *J. Org. Chem.* **1998**, 63, 3647; i) Y. Lu, G. Just, *Tetrahedron* **2000**, 56, 4355.
- [6] a) F. Ungemach, M. DiPierro, R. Weber, J. M. Cook, *J. Org. Chem.* **1981**, 46, 164; b) E. D. Cox, L. K. Hamaker, J. Li, P. Yu, K. M. Czerwinski, L. Deng, D. W. Bennett, J. M. Cook, *J. Org. Chem.* **1997**, 62, 44.
- [7] a) R. P. Iyer, D. Yu, T. Devlin, N.-H. Ho, S. Agrawal, *J. Org. Chem.* **1995**, 60, 5388; b) R. P. Iyer, D. Yu, N.-H. Ho, W. Tan, S. Agrawal, *Tetrahedron: Asymmetry* **1995**, 6, 1051.
- [8] R. P. Iyer, W. Egan, J. B. Regan, S. L. Beaucage, *J. Am. Chem. Soc.* **1990**, 112, 1253.
- [9] M. J. Damha, P. A. Giannaris, S. U. Zabarylo, *Nucleic Acids Res.* **1990**, 18, 3813.
- [10] a) W. J. Stec, G. Zon, B. Uznanski, *J. Chromatogr.* **1985**, 326, 263; b) H. Kanehara, T. Wada, M. Mizuguchi, K. Makino, *Nucleosides Nucleotides* **1996**, 15, 1169.



Scheme 1. Synthesis of α -diazomethyl zirconium complexes **4a**, **b** and **5a**, **b**.

chemical shift of the starting reagent **2** was monitored by ^1H NMR spectroscopy, which showed the presence of a broad, low-field signal for the NH proton at $\delta = 9.55$ and 9.74 in the final products **4a** and **4b**, respectively. In addition to the chemical shifts corresponding to the aza-zirconocene metallacycle skeleton (the resonances for the Cp groups appear as one singlet), the ^1H and ^{13}C NMR spectra exhibited signals in the region expected for the CO_2Et ester group bonded to the diazo functionality. The most striking data of the C-coordinated $\text{C}(\text{N}_2)$ diazo group in **4a** and **4b** are the ^{13}C NMR chemical shifts at $\delta = 48.1$ and 47.0 ,^[2, 10] respectively, and the intense band at 2029 cm^{-1} in the IR spectra for $\text{C}=\text{N}=\text{N}$, which is shifted to lower frequency in comparison with that of the starting diazoalkanes.^[2] All these data are fully in accordance with the structure proposed for **4a** and **4b**.


An identical experimental procedure was conducted with trimethylsilyldiazomethane (**3**) and **1a**, **b** to prepare the corresponding C- α -metalated diazoalkane complexes **5a** and **5b**, which were isolated in good yield (Scheme 1). The ^1H and ^{13}C NMR spectra exhibited the expected chemical shifts for the SiMe_3 fragment and showed signals at $\delta = 30.5$ (**5a**) and 36.7 (**5b**) for the $\text{Zr}-\text{C}(\text{N}_2)$ carbon atom.^[2, 10] The IR spectra displayed a characteristic band at 2069 (**5a**) and 2049 cm^{-1} (**5b**) for the diazo $\text{C}=\text{N}=\text{N}$ group. Thus spectroscopic and analytical data^[11] are consistent with the formulation of **4a**, **b** and **5a**, **b** as the first α -C-zirconated diazoalkanes.

Interestingly when **1a** and **1b** were allowed to react with the “nonterminal” diazo compound **6**, the pyrazoles **8a** and **8b** were obtained in good yield (Scheme 2). These new compounds were unambiguously characterized by mass spectrometry as well as by IR and NMR spectroscopy (see the Experimental Section). Formation of the metalated heterocycles **8a** and **8b** could be rationalized by O–H bond activation of the enol tautomer of substrate **6**, leading to the zirconocene diazo intermediates **7a** and **7b**, which rearrange by an intramolecular [3+2] cycloaddition to give the isolated products **8a** and **8b**.


In a first approach, the reactivity of the C- α -metalated diazoalkane complex **4a** was tested by addition of phenyl isocyanate (**9**) to afford the 1,2,3-triazolato zirconocene enolate complex **13** (Scheme 3). As evidenced from ^{31}P , ^1H , and ^{13}C NMR spectroscopies, the structure of the aza-zirconocene metallacycle skeleton remained unchanged. In particular, the ^1H NMR signal of the NH group appeared as expected at $\delta = 9.59$ and the ^{13}C NMR shift for $\text{P}-\text{C}=\text{N}$ was observed at $\delta = 201.3$ with $J_{\text{C,P}} = 38.5\text{ Hz}$ (**4a**: $\delta(^{13}\text{C}) = 203.9$,

$J_{\text{C,P}} = 39.3\text{ Hz}$). No IR absorption corresponding to a diazo group was identified in **13**. In the ^{13}C NMR spectrum the signals corresponding to the triazole skeleton at $\delta = 119$ and 162 are in the range usually observed for sp^2 carbon atoms bonded to ester and ether groups, respectively.

An identical reaction was conducted with naphthyl isocyanate (**10**) and **4a** (Scheme 3). The isolated triazole **14**^[11]



Scheme 2. Reactions of the enolizable diazo compound **6** with **1a**, **b**: **a**: R = NiPr_2 , **b**: R = NCy_2 .



Scheme 3. Cycloaddition reactions of the α -diazomethyl zirconium complex **4a**. **9**, **11**, **13**: R = Ph; **10**, **12**, **14**: R = α -Naph; **15**, **17**: R = CO_2Et , R' = Et; **16**, **18**: R = H, R' = Me.

exhibited the same spectroscopic data as **13**. However, to characterize unambiguously the cycloaddition products **13** and **14**, a single-crystal X-ray analysis of **14** was undertaken (Figure 1, see the Experimental Section). The features at the




Figure 1. ZORTEP drawing of the molecular structure of **14**. Selected bond lengths [Å] and angles [°]: Zr–O1 2.220(4), Zr–N1 2.258(4), Zr–C7 2.390(4), N1–C1 1.281(5), C1–P 1.861(4); O1–Zr–N1 70.89(12), N1–Zr–C7 67.71(13), Zr–O1–C8 146.1(2), Zr–N1–C1 126.1(3), Zr–C7–C2 116.3(3).



$\text{Cp}_2\text{Zr}^{\text{IV}}\text{-N}$ σ bonds ($\approx 2.20 \text{ \AA}$) and $\text{Cp}_2\text{Zr}^{\text{IV}}\text{-N}$ π -type interactions ($\approx 2.40 \text{ \AA}$), are the truly outstanding structural features in **14**.^[14] Therefore this product could formally be represented as the iminium–zirconate complex **II**.^[15]



Diazo compounds are classical precursors of carbenes. It is well known that transition metal complexes catalyze the decomposition of diazoacetates in the presence of acetylenes.^[18] Addition of **4a** to the acetylene dicarboxylate **15** resulted in loss of nitrogen and formation of the stable C-metalated cyclopropene compound **17**,^[11] which was isolated in 96 % yield (Scheme 3). The intense $\nu(\text{C}(\text{N}_2))$ absorption at 2029 cm^{-1} for **4a** totally disappeared, while characteristic bands for **17** at 1566 cm^{-1} ($\text{C}=\text{N}$), 1725 cm^{-1} (CO), and 3335 cm^{-1} (NH) were observed. The ^1H and ^{13}C chemical shifts of the aza-zirconocene metallacycle skeleton in **17** appeared in the expected range (^1H NMR: $\delta(\text{NH})=10.64$; ^{13}C NMR: $\delta(\text{P}-\text{C}=\text{N})=202.0$, $J_{\text{C,P}}=38.5\text{ Hz}$). The cyclopropene ring in **17** exhibited, beside the chemical shift of the corresponding CO_2Et ester groups, two singlets at $\delta=118.7\text{ ppm}$ ($\text{sp}^2\text{-C}$) and 146.1 ppm ($\text{Zr}-\text{C}$).

Organometallocene Group 4 complexes are useful intermediates in organic synthesis since various procedures for the cleavage of carbon–zirconium bonds with electrophiles have been reported.^[20] Addition of chlorodiphenylphosphane (**19**)

4700 © WILEY-VCH Verlag GmbH, D-69451 Weinheim

Experimental Section

4a, b: Ethyl diazoacetate (**2**; 0.048 mL, 0.465 mmol for **4a**; 0.063 mL, 0.599 mmol for **4b**) was added at room temperature to a solution of **1a** (0.258 g, 0.465 mmol) or **1b** (0.428 g, 0.599 mmol) in toluene (5 mL). The mixture was stirred at room temperature for 10 min (30 min for **4b**) and then evaporated to dryness. The resulting solid residue was extracted with pentane (20 mL) and filtered. The volatile components were removed from the solution to give the product.

4b: Yellow solid, 0.437 g (88 % yield). IR (KBr): $\tilde{\nu}$ = 1571 $\text{v}(\text{C}=\text{N})$, 1661 $\text{v}(\text{C}=\text{O})$, 2029 $\text{v}(\text{C}=\text{N}=\text{N})$, 3355 cm^{-1} $\text{v}(\text{N}-\text{H})$; $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 58.5 (s); ^1H NMR (C_6D_6): δ = 0.83–1.89 (m, 40H; CH_2), 1.21 (t, J_{HH} = 7.1 Hz, 3H; OCH_2CH_3), 3.13 (m, 4H; NCH), 4.23 (q, J_{HH} = 7.1 Hz, 2H; OCH_2), 5.85 (s, 10H; CH_3P), 7.08–7.20 (m, 2H; CH_arom), 7.52 (d, J_{HH} = 8.0 Hz, 1H; CH_arom), 8.50 (m, 1H; CH_arom), 9.74 (d, J_{HP} = 5.5 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 15.4 (s; OCH_2CH_3), 26.3, 27.1 (s; CH_2), 35.5 (d, J_{CP} = 6.1 Hz; CH_2), 36.1 (d, J_{CP} = 4.2 Hz; CH_2), 47.0 (s; $\text{C}=\text{N}=\text{N}$), 58.8 (d, J_{CP} = 9.8 Hz; NCH), 59.9 (s; OCH_2), 109.6 (s; CH_3P), 123.4, 129.3, 142.1 (s; CH_arom), 130.1 (d, J_{CP} = 19.3 Hz; CH_arom), 146.7 (d, J_{CP} = 28.2 Hz; ZrCC), 174.5 (s; $\text{C}=\text{O}$), 196.9 (d, J_{CP} = 6.5 Hz; ZrC), 204.0 (d, J_{CP} = 43.6 Hz; $\text{PC}=\text{N}$); elemental analysis calcd for $\text{C}_{43}\text{H}_{64}\text{N}_5\text{O}_2\text{PZr}$ (829.23) (%): C 65.18, H 7.77, N 8.44; found: C 65.12, H 7.81, N 8.42.

5a,b: (Trimethylsilyl)diazomethane (**3**, 2.0 M in hexanes; 0.302 mL, 0.604 mmol for **5a**; 0.489 mL, 0.979 mmol for **5b**) was added at room temperature to a solution of **1a** (0.335 g, 0.604 mmol) or **1b** (0.700 g, 0.979 mmol) in toluene (8 mL). The mixture was heated under reflux for 20 min and then evaporated to dryness. The resulting solid residue was extracted with pentane (20 mL) and filtered. The volatile components were removed from the solution to give the product.

5a: Yellow solid, 0.287 g (71 % yield). IR (KBr): $\tilde{\nu}$ = 1571 ν (C=N), 2069 ν (C=N=N), 3325 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 57.2 (s); ^1H NMR (C_6D_6): δ = 0.35 (s, 9H; SiMe₃), 1.11 (d, $J_{\text{H,H}}$ = 6.6 Hz, 12H; CH₃), 1.18 (d, $J_{\text{H,H}}$ = 6.6 Hz, 12H; CH₃), 3.25 (m, 4H; NCH), 5.90 (s, 10H; CH_{CP}), 6.98–7.40 (m, 3H; CH_{arom}), 8.18 (d, $J_{\text{H,H}}$ = 7.9 Hz, 1H; CH_{arom}), 9.62 (d, $J_{\text{H,P}}$ = 5.6 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 2.5 (s; SiMe₃), 24.5 (d, $J_{\text{C,P}}$ = 7.1 Hz; CH₃), 24.6 (d, $J_{\text{C,P}}$ = 6.2 Hz, CH₃), 30.5 (s, C=N=N), 49.6 (d, $J_{\text{C,P}}$ = 10.9 Hz, NCH), 110.2 (s; CH_{CP}), 122.7, 129.5, 141.2 (s; CH_{arom}), 129.7 (d, $J_{\text{C,P}}$ = 12.7 Hz; CH_{arom}), 146.9 (d, $J_{\text{C,P}}$ = 23.9 Hz; ZrCC), 198.1 (d, $J_{\text{C,P}}$ = 6.4 Hz; ZrC), 202.2 (d, $J_{\text{C,P}}$ = 37.4 Hz; PC=N); elemental analysis calcd for C₃₃H₅₂N₃PSiZr (669.08) (%): C 59.24, H 7.83, N 10.46; found: C 59.19, H 7.79, N 10.51.

5b: Yellow solid, 0.600 g (74 % yield); IR (KBr): $\tilde{\nu}$ = 1571 ν (C=N), 2049 ν (C=N=N), 3315 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 63.7 (s); ^1H NMR (C_6D_6): δ = 0.35 (s, 9H; SiMe₃), 0.83–2.05 (m, 40H; CH₂), 2.91 (m, 4H; NCH), 5.93 (s, 10H; CH_{CP}), 6.98–7.40 (m, 3H; CH_{arom}), 8.28 (d, $J_{\text{H,H}}$ = 7.6 Hz, 1H; CH_{arom}), 9.70 (d, $J_{\text{H,P}}$ = 5.7 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 2.5 (s; SiMe₃), 26.3, 27.5 (s; CH₃), 35.6 (d, $J_{\text{C,P}}$ = 5.6 Hz; CH₂), 35.9 (br; CH₂), 36.7 (s; C=N=N), 59.7 (br; NCH), 110.2 (s; CH_{CP}), 122.7, 129.4, 141.1 (s; CH_{arom}), 129.8 (d, $J_{\text{C,P}}$ = 15.6 Hz; CH_{arom}), 147.1 (d, $J_{\text{C,P}}$ = 23.8 Hz; ZrCC), 198.2 (d, $J_{\text{C,P}}$ = 6.2 Hz; ZrC), 201.9 (d, $J_{\text{C,P}}$ = 39.2 Hz; PC=N); elemental analysis calcd for C₄₅H₆₈N₃PSiZr (829.34) (%): C 65.17, H 8.26, N 8.44; found: C 65.21, H 8.29, N 8.50.

8a,b: Ethyl diazoacetate (**6**; 0.053 mL, 0.388 mmol for **8a**; 0.069 mL, 0.500 mmol for **8b**) was added at room temperature to a solution of **1a** (0.215 g, 0.388 mmol) or **1b** (0.357 g, 0.500 mmol) in toluene (5 mL). The mixture was stirred at room temperature for 1 h and then evaporated to dryness. The resulting solid residue was extracted with pentane (15 mL) and filtered. The volatile components were removed from the solution to give the product.

8a: Yellow solid, 0.242 g (88 % yield). MS (FAB/MNBA): m/z : 709 [M^+]; IR (KBr): $\tilde{\nu}$ = 1551 ν (C=N), 1591 ν (C=N), 1715 ν (C=O), 3204 ν (N-H), 3355 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 51.6 (s); ^1H NMR (C_6D_6): δ = 0.87 (t, $J_{\text{H,H}}$ = 7.1 Hz, 3H; OCH₂CH₃), 1.11 (d, $J_{\text{H,H}}$ = 6.7 Hz, 12H; CH₃), 1.21 (d, $J_{\text{H,H}}$ = 6.4 Hz, 12H; CH₃), 3.46 (m, 4H; NCH), 4.16 (q, $J_{\text{H,H}}$ = 7.1 Hz, 2H; CH₂), 5.96 (s, 10H; CH_{CP}), 7.04–7.41 (m, 3H; CH_{arom}), 7.46 (s, 1H; =CH), 8.34 (m, 1H; CH_{arom}), 9.82 (d, $J_{\text{H,P}}$ = 5.3 Hz, 1H; C=NH), 11.29 (br, 1H; =NNH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 15.2 (s; OCH₂CH₃), 24.5 (d, $J_{\text{C,P}}$ = 6.0 Hz; CH₃), 24.8 (d, $J_{\text{C,P}}$ = 5.0 Hz; CH₃), 49.3 (d, $J_{\text{C,P}}$ = 11.2 Hz; NCH), 59.6 (s; CH₂), 111.2 (s; CH_{CP}), 122.9, 129.5, 141.3 (s; CH_{arom}), 129.8 (d, $J_{\text{C,P}}$ = 17.4 Hz; CH_{arom}), 139.5 (s; =CH), 147.8 (d, $J_{\text{C,P}}$ = 26.1 Hz; ZrCC), 148.9 (s; C=N), 158.5 (s; =C-O), 160.5 (s; C=O), 198.6 (s; ZrC), 201.3 (d, $J_{\text{C,P}}$ = 38.6 Hz; PC=N); elemental analysis calcd for C₃₅H₅₀N₃O₃PZr (711.00) (%): C 59.12, H 7.08, N 9.85; found: C 59.07, H 7.12, N 9.81.

8b: Orange solid, 0.309 g (71 % yield). IR (KBr): $\tilde{\nu}$ = 1536 ν (C=N), 1561 ν (C=N), 1725 ν (C=O), 3296 ν (N-H), 3325 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 64.0 (s); ^1H NMR (C_6D_6): δ = 0.77–1.94 (m, 43H; CH₂, CH₃), 2.97 (m, 4H; NCH), 4.25 (q, $J_{\text{H,H}}$ = 6.8 Hz, 2H; OCH₂), 6.04 (s, 10H; CH_{CP}), 6.99–7.24 (m, 2H; CH_{arom}), 7.41 (d, $J_{\text{H,H}}$ = 6.8 Hz, 1H; CH_{arom}), 7.47 (s, 1H; =CH), 8.47 (d, $J_{\text{H,H}}$ = 7.1 Hz, 1H; CH_{arom}), 9.88 (d, $J_{\text{H,P}}$ = 5.2 Hz, 1H; C=NH), 10.54 (br, 1H; =NNH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 15.2 (s; CH₃), 26.3 (s; CH₂), 27.4 (d, $J_{\text{C,P}}$ = 6.3 Hz; CH₂), 35.6 (br; CH₂), 59.5 (br; NCH), 59.7 (s; OCH₂), 111.3 (s; CH_{CP}), 122.7, 129.5, 141.1 (s; CH_{arom}), 130.2 (d, $J_{\text{C,P}}$ = 13.1 Hz; CH_{arom}), 139.4 (s; =CH), 147.4 (d, $J_{\text{C,P}}$ = 24.1 Hz; ZrCC), 147.6 (s; C=N), 157.9 (s; =C-O), 160.6 (s; C=O), 199.0 (d, $J_{\text{C,P}}$ = 5.0 Hz; ZrC), 201.6 (d, $J_{\text{C,P}}$ = 41.3 Hz; PC=N); elemental analysis calcd for C₄₇H₆₆N₃O₃PZr (871.26) (%): C 64.79, H 7.63, N 8.03; found: C 64.73, H 7.68, N 8.07.

17 and **18:** Diethyl acetylenedicarboxylate (**15**; 0.049 mL, 0.311 mmol for **17**) or methyl propiolate (**16**; 0.071 mL, 0.803 mmol for **18**) was added at room temperature to a solution of **4a** (0.208 g, 0.311 mmol for **17**; 0.537 g, 0.803 mmol for **18**) in toluene (5 mL). The mixture was stirred at room

temperature for 1 h and then evaporated to dryness. The resulting solid residue was extracted with pentane (15 mL) and filtered. The volatile components were removed from the solution to give the product.

17: Yellow solid, 0.242 g (96 % yield). IR (KBr): $\tilde{\nu}$ = 1566 ν (C=N), 1725 ν (C=O), 3335 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 40.9 (s); ^1H NMR (C_6D_6): δ = 0.94–1.18 (m, CH₃, 27H; OCH₂CH₃), 1.35 (d, $J_{\text{H,H}}$ = 6.5 Hz, 6H; CH₃), 3.69 (m, 4H; NCH), 4.22 (q, $J_{\text{H,H}}$ = 7.2 Hz, 2H; CH₂), 4.27 (q, $J_{\text{H,H}}$ = 7.2 Hz, 4H; CH₂), 5.86 (s, 10H; CH_{CP}), 7.02–7.56 (m, 3H; CH_{arom}), 8.49 (m, 1H; CH_{arom}), 10.64 (d, $J_{\text{H,P}}$ = 4.8 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 14.7, 14.9 (s; OCH₂CH₃), 24.5 (d, $J_{\text{C,P}}$ = 6.3 Hz; CH₃), 24.8 (d, $J_{\text{C,P}}$ = 6.6 Hz; CH₃), 49.3 (d, $J_{\text{C,P}}$ = 12.3 Hz; NCH), 60.7, 61.4 (s; CH₂), 111.4 (s; CH_{CP}), 118.7 (s; =C), 123.5, 129.6, 141.6 (s; CH_{arom}), 130.4 (d, $J_{\text{C,P}}$ = 24.4 Hz; CH_{arom}), 146.1 (s; ZrCCO₂Et), 147.3 (d, $J_{\text{C,P}}$ = 30.1 Hz; ZrCC), 164.7, 164.8 (s; C=O), 197.8 (d, $J_{\text{C,P}}$ = 6.7 Hz; ZrC), 202.0 (d, $J_{\text{C,P}}$ = 40.2 Hz; PC=N); elemental analysis calcd for C₄₁H₅₈O₆N₃PZr (811.10) (%): C 60.71, H 7.20, N 5.18; found: C 60.76, H 7.18, N 5.21.

18: Yellow solid, 0.456 g (78 % yield). IR (KBr): $\tilde{\nu}$ = 1561 ν (C=N), 1710 ν (C=O), 3305 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 38.3 (s); ^1H NMR (C_6D_6): δ = 0.97 (d, $J_{\text{H,H}}$ = 6.8 Hz, 6H; CH₃), 1.02 (d, $J_{\text{H,H}}$ = 6.5 Hz, 6H; CH₃), 1.10 (m, CH₃, 9H; OCH₂CH₃), 1.37 (d, $J_{\text{H,H}}$ = 6.5 Hz, 6H; CH₃), 3.57 (s, 3H; OCH₃), 3.82 (m, 4H; NCH), 4.15 (q, $J_{\text{H,H}}$ = 7.1 Hz, 2H; CH₂), 5.88 (s, 10H; CH_{CP}), 6.99–7.53 (m, 3H; CH_{arom}), 7.79 (s; =CH), 8.53 (m, 1H; CH_{arom}), 11.22 (d, $J_{\text{H,P}}$ = 5.0 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 14.9 (s; OCH₂CH₃), 24.7 (d, $J_{\text{C,P}}$ = 6.4 Hz; CH₃), 24.9 (d, $J_{\text{C,P}}$ = 6.6 Hz; CH₃), 49.2 (d, $J_{\text{C,P}}$ = 12.7 Hz; NCH), 51.3 (s; OCH₃), 60.3 (s; CH₂), 111.6 (s; CH_{CP}), 117.3 (s; =CH), 123.2, 129.9, 140.9 (s; CH_{arom}), 130.4 (d, $J_{\text{C,P}}$ = 26.7 Hz; CH_{arom}), 144.7, 145.3 (s; ZrCCO₂Et, =C), 147.5 (d, $J_{\text{C,P}}$ = 32.7 Hz; ZrCC), 163.5, 163.9 (s; C=O), 198.9 (d, $J_{\text{C,P}}$ = 7.9 Hz; ZrC), 201.5 (d, $J_{\text{C,P}}$ = 41.2 Hz; PC=N); elemental analysis calcd for C₃₇H₅₂O₆N₃PZr (725.01) (%): C 61.29, H 7.22, N 5.79; found: C 61.25, H 7.16, N 5.83.

21, 23: Chlorophosphane **20** (0.056 mL, 0.311 mmol) was added at room temperature to a solution of **5a** (0.208 g, 0.311 mmol) in toluene (5 mL). The mixture was stirred at room temperature for 1 h and then evaporated to dryness. The resulting solid residue was extracted with pentane (20 mL) and filtered. The volatile components were removed from the solution to give **21** and **23** as the unique products of the reaction according to ^{31}P NMR spectroscopy. Compound **23** was identified in the reaction mixture by comparison with an authentic sample.^[22] Independently, addition of HCl to **4a** allowed the preparation and isolation of **21** as a yellow solid: IR (KBr): $\tilde{\nu}$ = 1576 ν (C=N), 3305 cm^{-1} ν (N-H); $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6): δ = 58.2 (s); ^1H NMR (C_6D_6): δ = 1.11 (d, $J_{\text{H,H}}$ = 6.6 Hz, 12H; CH₃), 1.21 (d, $J_{\text{H,H}}$ = 6.6 Hz, 12H; CH₃), 3.27 (d sept, $J_{\text{H,H}}$ = 6.6 Hz, $J_{\text{H,P}}$ = 12.1 Hz, 4H; NCH), 5.90 (s, 10H; CH_{CP}), 7.20 (m, 2H; CH_{arom}), 7.88 (m, 4H; CH_{arom}), 8.25 (m, 1H; CH_{arom}), 10.12 (d, $J_{\text{H,P}}$ = 5.0 Hz, 1H; NH); $^{13}\text{C}\{^1\text{H}\}$ NMR (C_6D_6): δ = 24.4 (d, $J_{\text{C,P}}$ = 6.8 Hz; CH₃), 25.1 (d, $J_{\text{C,P}}$ = 5.0 Hz; CH₃), 49.6 (d, $J_{\text{C,P}}$ = 11.2 Hz; NCH), 111.3 (s; CH_{CP}), 122.8, 129.8, 141.2 (s; CH_{arom}), 130.1 (d, $J_{\text{C,P}}$ = 14.3 Hz; CH_{arom}), 146.1 (d, $J_{\text{C,P}}$ = 24.1 Hz; ZrCC), 198.6 (d, $J_{\text{C,P}}$ = 6.3 Hz; ZrC), 202.5 (d, $J_{\text{C,P}}$ = 40.6 Hz; PC=N).

X-ray crystal structure analysis of **14** (C₄₄H₄₉N₆O₃PZr·1.5C₆H₆): M_r = 979.72, monoclinic, space group $P2_1/n$, T = 160 K, a = 13.811(6), b = 18.356(5), c = 20.031(8) Å, β = 98.26(5)°, V = 5025(3) Å³, Z = 4, μ = 0.296 mm⁻¹. Of 27371 reflections collected, 6500 were independent; R_{int} = 0.0970; final R indices: $R1$ = 0.0402, $wR2$ = 0.0889 ($I > 2\sigma(I)$), $R1$ = 0.0729, $wR2$ = 0.0992 (all data). Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-149738. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).

Received: July 12, 2000 [Z15438]

- [1] For N-bonded diazoalkane transition metal compounds, see for example: G. L. Hillhouse, B. L. Haymore, *J. Am. Chem. Soc.* **1982**, *104*, 1537; L. K. Bell, W. A. Hermann, G. W. Kriechbaum, H. Pfisterer, M. L. Ziegler, *J. Organomet. Chem.* **1982**, *240*, 381; C. Woodcock, R. Eisenberg, *Organometallics* **1985**, *4*, 4; M. Cowie, S. J. Loeb, I. R. McKeer, *Organometallics* **1986**, *5*, 854; J. Wolf, L. Brandt, Y. Fries, H. Werner, *Angew. Chem.* **1990**, *102*, 584; *Angew. Chem. Int. Ed. Engl.*

- 1990, 29, 510; Y. Mizobe, Y. Ishii, M. Hidai, *Coord. Chem. Rev.* **1995**, 139, 281.
- [2] For C-bonded diazoalkane late transition metal compounds, see the following: Pd complexes: S. I. Murahashi, Y. Kitani, T. Hosokawa, K. Miki, N. Kasai, *J. Chem. Soc. Chem. Commun.* **1979**, 450; Os complexes: M. A. Gallo, T. C. Jones, C. E. F. Rickard, W. R. Roper, *J. Chem. Soc. Chem. Commun.* **1979**, 450; Rh complexes: E. Deydier, M.-J. Menu, M. Dartiguenave, Y. Dartiguenave, M. Simard, A. L. Beauchamp, J. C. Brewer, H. B. Gray, *Organometallics* **1996**, 15, 1166; M.-J. Menu, J. Desrosiers, M. Dartiguenave, Y. Dartiguenave, G. Bertrand, *Organometallics* **1987**, 6, 1822; Ni complexes: M.-J. Menu, H. König, M. Dartiguenave, Y. Dartiguenave, H. F. Klein, *J. Am. Chem. Soc.* **1990**, 112, 5351.
- [3] a) T. Luker, R. J. Whitby, M. Webster, *J. Organomet. Chem.* **1995**, 492, 53; b) G. A. Vaughan, G. L. Hillhouse, A. L. Rheingold, *J. Am. Chem. Soc.* **1990**, 112, 7994; c) S. Gamboratta, C. Floriani, A. Chiesi-Villa, C. Guastini, *Inorg. Chem.* **1983**, 22, 2029.
- [4] E. Hey, U. Müller, *Z. Naturforsch. B* **1989**, 44, 1538; E. Hey, M. F. Lappert, J. L. Atwood, S. G. Bott, *Polyhedron* **1988**, 7, 2083; E. Hey, F. Weller, *Chem. Ber.* **1988**, 1207.
- [5] T. A. Hanna, A. M. Baranger, R. G. Bergman, *Angew. Chem.* **1996**, 108, 693; *Angew. Chem. Int. Ed. Engl.* **1996**, 35, 653.
- [6] V. Cadierno, B. Donnadieu, A. Igau, J.-P. Majoral, *Eur. J. Inorg. Chem.* **2000**, 417.
- [7] U. Blaschke, G. Erker, M. Nissinen, E. Wegelius, R. Fröhlich, *Organometallics* **1999**, 18, 1224, and references therein.
- [8] V. Cadierno, M. Zablocka, B. Donnadieu, A. Igau, J.-P. Majoral, A. Skowronska, *Chem. Eur. J.*, in press.
- [9] J. A. Davies, P. L. Watson, J. F. Liebman, A. Greeberg, *Selective Hydrocarbon Activation*, VCH, New York, **1990**; *Activation and Functionalization of Alkanes* (Ed.: C. L. Hill), Wiley, New York, **1989**; A. E. Shilov, *Activation of Saturated Hydrocarbons by Transition Metal Complexes*, Reidel, Hingham, MA, **1984**; W. D. Jones, F. J. Feher, *Acc. Chem. Res.* **1989**, 2, 91; A. D. Ryabov, *Chem. Rev.* **1990**, 90, 403; A. Sen, *Acc. Chem. Res.* **1988**, 21, 421; R. H. Crabtree, *Chem. Rev.* **1985**, 85, 245.
- [10] K. Horchler von Locquenien, R. Réau, G. Bertrand, *J. Chem. Soc. Chem. Commun.* **1991**, 1192.
- [11] Elemental analyses gave satisfactory results, see the Experimental Section.
- [12] D. J. Cardin, M. F. Lappert, C. L. Raston, *Chemistry of Organo-Zirconium and -Hafnium Compounds*, Wiley, New York, **1986**, and references therein.
- [13] L. Pauling, *The Nature of the Chemical Bond*, 3rd ed., Cornell University Press, Ithaca, NY, **1960**, p. 260.
- [14] For Zr–N σ/π bonding, see for example C. E. Holloway, I. M. Walker, M. Melnik, *J. Organomet. Chem.* **1987**, 321, 143; R. Fandos, M. Lanfranchi, A. Otero, M. A. Pellinghelli, M. J. Ruiz, P. Terreros, *Organometallics* **1996**, 15, 4725; P. J. Walsh, M. J. Carney, R. G. Bergman, *J. Am. Chem. Soc.* **1984**, 106, 2210.
- [15] For other five-coordinate zirconate complexes (containing Cp₂Zr) bonded to at least one nitrogen atom, see for example V. Cadierno, M. Zablocka, B. Donnadieu, A. Igau, J.-P. Majoral, A. Skowronska, *Chem. Eur. J.* **2000**, 6, 345; V. Cadierno, M. Zablocka, B. Donnadieu, A. Igau, J.-P. Majoral, *Organometallics* **1999**, 18, 1882; T. Luker, R. J. Whitby, M. Webster, *J. Organomet. Chem.* **1995**, 492, 53.
- [16] M. Regitz, *Diazoalkanes*, Thieme, Stuttgart, **1977**; D. S. Wulfman, G. Linstrumelle, C. F. Cooper, in *The Chemistry of Diazonium and Diazo Groups*, Vol. 2 (Ed.: S. Patai), Chap. 18, Wiley, New York, **1978**.
- [17] See for example D. M. Roddick, J. E. Bercaw, *Chem. Ber.* **1989**, 122, 1579; R. Choukroun, B. Douziech, F. Soleil, *J. Chem. Soc. Chem. Commun.* **1995**, 2017.
- [18] G. Maas, *Top. Curr. Chem.* **1987**, 137, 75.
- [19] The NMR spectroscopic assignment have been unambiguously confirmed by 2D HMQC NMR experiments: S. Braun, H. O. Kalinowski, S. Berger, *100 and more Basic NMR Experiments*, VCH, Weinheim, **1996**.
- [20] E. Negishi, J.-L. Montchamp in *Metalloenes, Synthesis, Reactivity, Applications*, Vol. 1 (Eds.: A. Togni, R. L. Halterman), Wiley-VCH, Weinheim, **1998**, p. 241; E. Negishi, *Tetrahedron* **1995**, 51, 4255; reviews: U. Annby, S. Karlsson, *Acta Chem. Scand.* **1993**, 47, 425; J. A. Labinger in *Comprehensive Organic Synthesis*, Vol. 3 (Eds.: B. M.

Trost, I. Fleming, G. Pattenden), Pergamon, Oxford, **1991**, p. 667; E. Negishi, *Comprehensive Organic Synthesis*, Vol. 5 (Eds.: B. M. Trost, I. Fleming, L. A. Paquette), Pergamon, Oxford, **1991**, p. 1163; G. Erker, M. Aulbach, M. Mena, R. Pfaff, F. Sosna, *Chem. Scr.* **1989**, 29, 451.

[21] H. Staudinger, J. Meyer, *Helv. Chim. Acta* **1919**, 2, 61; G. Singh, H. Zimmer, *Organomet. Chem. Rev.* **1967**, 279.

[22] IR and NMR spectroscopies were in full agreement with the reported data: A. Baceiredo, G. Bertrand, G. Sicard, *J. Am. Chem. Soc.* **1985**, 107, 4781.

Hybrid Coordination Polymers—Metal Oxide Compounds with Chiral Structures**

Li-Min Zheng, Tabatha Whitfield, Xiqu Wang, and Allan J. Jacobson*

Many investigations of the design and synthesis of metal–organic compounds with porous structures have been motivated by potential applications in molecule recognition, separation, and catalysis.^[1–11] By using rigid, multifunctional ligands, two- or three-dimensional microporous coordination polymers have been obtained with desired pore sizes and shapes.^[3–11] Other metal–organic frameworks have been reported that, though they may have other valuable properties, are not microporous because of lattice interpenetration,^[12] or are unstable on removal or exchange of their guest molecules or ions. Very few examples of either framework or microporous chiral metal–organic structures have been reported.^[13–17]

One strategy to enhance the stability with respect to removal of the guest species is to attach the metal–organic components to a thermally more stable inorganic backbone. Several examples of compounds of this type containing both ligated metal cation and metal oxide components are known.^[18–20] These extended lattices are formed in different ways, for example, by connecting a metal oxide cluster with a coordination complex, a metal oxide cluster anion with a coordination polymer, a metal oxide lattice with an isolated coordination complex, and metal oxide lattice with a coordination polymer. All of the inorganic components in the known examples are molybdenum or vanadium oxides.^[18–20]

[*] Prof. A. J. Jacobson, Dr. L.-M. Zheng,^[+] T. Whitfield, Dr. X. Wang
Department of Chemistry
University of Houston
Houston, TX 77204-5641 (USA)
Fax: (+1) 713-743-2787
E-mail: ajacob@uh.edu

[+] Permanent address:
Coordination Chemistry Institute, Department of Chemistry
Nanjing University, Nanjing 210093 (P. R. China)

[**] We thank the National Science Foundation (DMR-9805881) and the Robert A. Welch Foundation for support. This work made use of MRSEC/TCSUH Shared Experimental Facilities supported by the National Science Foundation under Award Number DMR-9632667 and the Texas Center for Superconductivity at the University of Houston.