

7.57 (d, $J = 7$ Hz, 1H), 7.48 (d, $J = 7$ Hz, 1H), 7.40 (t, $J = 7$ Hz, 1H), 5.00 (d, $J = 3$ Hz, 6H), 3.83 (d, $J = 9$ Hz, 6H), 3.61 (dd, $J' = 10$, $J'' = 3$ Hz, 6H), 3.52 (s, 3H), 3.49 (t, $J = 10$ Hz, 6H), 3.34 (d, $J = 10$ Hz, 6H), 3.31 (s, 18H), 3.26 (s, 18H), 3.24 (s, 18H), 3.19 (t, $J = 10$ Hz, 6H), 3.01 (dd, $J' = 10$, $J'' = 3$ Hz, 6H); ^{13}C NMR (125 MHz, D_2O): $\delta = 169.1, 165.4, 164.0, 150.7, 150.5, 147.7, 143.8, 141.8, 141.5, 138.5, 133.9, 133.1, 131.7, 129.94, 129.93, 129.7, 128.9, 128.1, 124.9, 123.7, 121.3, 118.5, 117.4, 100.0, 82.4, 81.4, 81.1, 71.4, 71.1, 62.5, 59.0, 57.6, 39.1$; UV/Vis (H_2O): $\lambda_{\text{max}} = 364$ nm, $\lg \epsilon = 4.3$; ES-MS (negative): m/z : 964.2 [$\text{M}^{3-} + \text{Na}^+$].

Received: November 17, 2000 [Z16126]

- [1] D. J. Cram, M. E. Tanner, R. Thomas, *Angew. Chem.* **1991**, *103*, 1048–1051; *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 1024–1027; R. Warmuth, *Angew. Chem.* **1997**, *109*, 1406–1409; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1347–1350; R. Warmuth, M. A. Marvel, *Angew. Chem.* **2000**, *112*, 1168–1171; *Angew. Chem. Int. Ed.* **2000**, *39*, 1117–1119.
- [2] a) S. Anderson, T. D. W. Claridge, H. L. Anderson, *Angew. Chem.* **1997**, *109*, 1367–1370; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1310–1313; b) S. Anderson, W. Clegg, H. L. Anderson, *Chem. Commun.* **1998**, 2379–2378; c) M. R. Craig, T. D. W. Claridge, M. G. Hutchings, H. L. Anderson, *Chem. Commun.* **1999**, 1537–1538; d) J. E. H. Buston, J. R. Young, H. L. Anderson, *Chem. Commun.* **2000**, 905–906.
- [3] S. Anderson, R. T. Aplin, T. D. W. Claridge, T. Goodson, A. C. Maciel, G. Rumbles, J. F. Ryan, H. L. Anderson, *J. Chem. Soc. Perkin Trans. 1* **1998**, 2383–2397; P. N. Taylor, M. J. O'Connell, L. A. McNeill, M. J. Hall, R. T. Aplin, H. L. Anderson, *Angew. Chem.* **2000**, *112*, 3598–3602; *Angew. Chem. Int. Ed.* **2000**, *39*, 3456–3460.
- [4] For recent work on rotaxanes, see: *Molecular Catenanes, Rotaxanes and Knots* (Eds.: J.-P. Sauvage, C. Dietrich-Buchecker), Wiley-VCH, Weinheim, **1999**; S. A. Nepogodiev, J. F. Stoddart, *Chem. Rev.* **1998**, *98*, 1959–1976.
- [5] Other rotaxanes containing azo-benzene units have been reported: H. Murakami, A. Kawabuchi, K. Kotoo, M. Kunitake, N. Nakashima, *J. Am. Chem. Soc.* **1997**, *119*, 7605–7606; C. Kauffmann, W. M. Müller, F. Vögtle, S. Weinman, S. Abramson, B. Fuchs, *Synthesis* **1999**, 849–853; M. Asakawa, P. R. Ashton, V. Balzani, C. L. Brown, A. Credi, O. A. Matthews, S. P. Newton, F. M. Raymo, A. N. Shipway, N. Spence, A. Quick, J. F. Stoddart, A. J. P. White, D. J. Williams, *Chem. Eur. J.* **1999**, *5*, 860–875; T. Fujimoto, Y. Sakata, T. Kaneda, *Chem. Commun.* **2000**, 2143–2144.
- [6] H. Zollinger, *Color Chemistry*, VCH, Weinheim, **1991**.
- [7] M. Kunitake, K. Kotoo, O. Manabe, T. Muramatsu, N. Nakashima, *Chem. Lett.* **1993**, 1033–1036.
- [8] K. Harata, *J. Chem. Soc. Perkin Trans. 2* **1990**, 799–804; K. Harata, K. Uekama, M. Otagiri, F. Hirayama, *Bull. Chem. Soc. Jpn.* **1982**, *55*, 3904–3910.
- [9] UV/Vis titrations in water at 298 K showed that the complex **2**·TM- α -CD has a stability constant of $(5 \pm 1) \times 10^5 \text{ M}^{-1}$ compared to only $(1.1 \pm 0.1) \times 10^4 \text{ M}^{-1}$ for **2**· α -CD. It is difficult to obtain an accurate value of K_1 for **2**·TM- α -CD because a weak complex **2**·2 TM- α -CD is also formed, $K_2 \approx 100 \text{ M}^{-1}$. For other work on the stability of azo dye CD complexes, see: N. Yoshida, *J. Chem. Soc. Perkin Trans. 2* **1995**, 2249–2256.
- [10] The reason for exclusive formation of one isomer of **1** has yet to be elucidated but it is probably related to the phenomenal stability of the complex **2**·TM- α -CD.
- [11] A. Püntener, J. Fennen, C. T. Page, *J. Soc. Leather Technol. Chem.* **1996**, *80*, 1–5; L. F. Fieser, *Organic Synthesis, Vol. II*, Wiley, New York, **1943**, p. 35.
- [12] F. Ferreo, *J. Soc. Dyers Colour.* **2000**, *116*, 148–153.
- [13] Light-fastness was measured in conformance with International Standard ISO 105-B02 in a Xenotest 150 fadeometer, using a xenon arc lamp to mimic daylight. The degree of fading was assessed against ISO blue wool standards, irradiated simultaneously. Cotton samples dyed with rotaxane **1** had a light-fastness rating of 5, whereas those dyes with free dye **3** had a rating of 3–4.

2-Pyridyldimethylsilyl as a Removable Hydrophilic Group in Aqueous Diels–Alder Reactions**

Kenichiro Itami, Toshiki Nokami, and Jun-ichi Yoshida*

During the last two decades water has been rediscovered as a most intriguing medium for organic synthesis.^[1] However, the execution of organic reactions in water is not straightforward since many organic compounds, reagents, and catalysts are insoluble or unstable in it. Therefore, many of the reported aqueous organic reactions have had to rely on the use of organic co-solvents at the expense of the many inherent advantages and unique properties of water, such as hydrophobic effects, hydrogen-bonding interactions, and high polarity.^[1]

During the course of our study on the development of multifunctional phase tags for solution-phase synthesis it was found that organic molecules bearing the 2-pyridyldimethylsilyl (2-PyMe₂Si) group are miscible with water when HCl is added.^[2,3] Molecular aggregation seems to be involved in their dissolution so as to minimize the energetically unfavorable contact between water molecules and nonpolar carbon chains. In addition, we have also established that the 2-PyMe₂Si group can be easily removed from organic molecules by oxidation with H₂O₂ to afford the corresponding alcohols.^[4]

We envisioned that if the 2-PyMe₂Si group works as a removable hydrophilic group and induces molecular aggregation in water, organic reactions in aqueous molecular aggregates should be possible when the reaction site is located on a hydrophobic side chain of the 2-PyMe₂Si-substituted molecule (Figure 1).^[5] The advantage of using the 2-PyMe₂Si




Figure 1. New strategy for aqueous organic reactions using a removable hydrophilic group.


[*] Prof. J.-i. Yoshida, Dr. K. Itami, T. Nokami
Department of Synthetic Chemistry and Biological Chemistry
Graduate School of Engineering, Kyoto University
Yoshida, Kyoto 606-8501 (Japan)
Fax: (+81) 75-753-5911
E-mail: yoshida@sbchem.kyoto-u.ac.jp

[**] This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture, Japan, and in part by the Mitsubishi Foundation. We thank Professor Kazunari Akiyoshi (Kyoto University) for assistance with dynamic light-scattering experiments and fruitful discussions. Professor Susumu Yoshikawa and Professor Masahiro Kinoshita (Kyoto University) are gratefully acknowledged for critical discussions.




Supporting information for this article is available on the WWW under <http://www.angewandte.com> or from the author.

group is apparent, as it can be readily removed from the molecules when required either by desilylation or oxidation. We chose an intermolecular Diels–Alder reaction of the 2-PyMe₂Si-substituted 1,3-dienes **1–3** to evaluate our strategy of using a removable hydrophilic group in aqueous organic reactions.^[1, 6–9]



First, the solution behavior of the 2-PyMe₂Si-substituted 1,3-dienes **1–3** in water was examined by dynamic light-scattering experiments on aqueous solutions (containing 1.0 equiv of HCl) of these dienes. These experiments revealed the presence of molecular aggregates of **1–3** with average hydrodynamic radii of 197, 138, and 307 nm, respectively. However, the determination of the exact structure of the molecular aggregates must await further investigations. The large radii observed can probably be attributed to nonordered structures of molecular aggregates in aqueous solution. The detection of molecular aggregates led us to explore aqueous reactions of **1–3**, irrespective of the structures of their molecular aggregates.

The Diels–Alder reactions of **1** and **2** with *p*-benzoquinone occurred at room temperature in water with simultaneous desilylation and aromatization to afford, respectively, naphthoquinones **4** and **5** quantitatively [Eq. (1)]. In these cases, an additional chemical operation to remove the 2-PyMe₂Si group was unnecessary.




Several control experiments were conducted to assess the effect of the 2-PyMe₂Si group (Table 1). Decreasing the amount of HCl added slowed down the reaction enormously and only a trace amount of **4** was formed (entries 2 and 3). We contend that at least one equivalent of HCl is required for the 2-PyMe₂Si-substituted diene to form molecular aggregates and that the reaction probably occurs in the interior of aggregates where there are enhanced hydrophobic interactions,^[10] and not in the aqueous bulk phase.^[6] The use of H₂SO₄ also gave rise to a rate enhancement, but not as

Table 1. Aqueous Diels–Alder reactions of **1** with *p*-benzoquinone.^[a]


Entry	Solvent	Additive (equiv)	Time [h]	Yield of 4 [%] ^[b]
1	H ₂ O	HCl (1.0) ^[c]	38	quant
2	H ₂ O	HCl (0.2)	47	2
3	H ₂ O	–	89	4
4	H ₂ O	H ₂ SO ₄ (1.0)	38	73
5	Et ₂ O/toluene	HCl (1.0)	38	22
6	Et ₂ O/toluene	–	38	6

[a] All reactions were performed at room temperature using **1** (1.0 equiv, 1.0 M concentration) and *p*-benzoquinone (3.0 equiv). [b] Yields of isolated product. [c] The addition of 2.0 equivalents of HCl did not affect the rate and yield.

dramatic as with HCl (entry 4). The reaction carried out in organic solvent resulted in a much lower yield of the product (entries 5 and 6), which indicates the rate acceleration occurs in water. In contrast, the reaction with the PhMe₂Si-substituted diene **6** gave no Diels–Alder adduct at all in water [Eq. (2)].




Next, we examined the aqueous reaction of 1,3-pentadiene with *p*-benzoquinone in the presence of protonated 2-pyridyldimethyloctylsilane (**7**) as a surfactant. The hard-to-dissolve 1,3-pentadiene dissolved in water when **7** and HCl were added. Thereafter, *p*-benzoquinone was added to the solution. However, in sharp contrast with the reaction of **1** or **2**, no cycloadduct was observed at room temperature [Eq. (3)]. It is apparent that the use of a reacting surfactant has an advantage in terms of reactivity over the combined use of an external surfactant.^[6b, 11]



The aqueous Diels–Alder reaction of **3** with *p*-benzoquinone was complete within one hour to give the cycloadduct **8** (Scheme 1).^[12] In this case a catalytic amount of HCl (0.2 equiv) was enough to promote the reaction. Interestingly, the formation of molecular aggregates (average hydrodynamic radii: 65 nm) was observed even at the end of the reaction. Moreover, the rate acceleration in water was again observed (12% yield of **8** in Et₂O/toluene).

The transformation of **8** was examined to test the chemical stability and removability of the 2-PyMe₂Si group (Scheme 1). The reduction of **8** afforded the substituted hydroquinone **9** in 73% yield. The protection of the phenolic hydroxyl groups with benzyl groups afforded **10** in 59% yield (78% by NMR spectroscopy). Importantly, the 2-PyMe₂Si group was unaf-



Scheme 1. Aqueous Diels–Alder reaction of **3** and removal of the 2-PyMe₂Si group. Bn = benzyl.

ected throughout these transformations, which indicates the reasonable chemical stability of this group. Finally, the oxidation of **10** with H₂O₂ afforded **11** in 89% yield without affecting the C–C double bond or eroding the stereochemistry at the allylic carbon atoms.

In summary, we have demonstrated the “proof-of-principle” of our strategy for aqueous organic reactions by utilizing the 2-PyMe₂Si group as a removable hydrophilic group. Importantly, the strategy described herein should not be limited to aqueous Diels–Alder reactions but could in principle be applied to other aqueous organic reactions as well.

Received: September 27, 2000 [Z15862]

- [1] a) *Organic Synthesis in Water* (Ed.: P. A. Grieco), Blackie Academic & Professional, London, **1998**; b) C.-J. Li, T.-H. Chen, *Organic Reactions in Aqueous Media*, Wiley, New York, **1997**.
- [2] J. Yoshida, K. Itami, K. Mitsudo, S. Suga, *Tetrahedron Lett.* **1999**, *40*, 3403.
- [3] a) K. Itami, K. Mitsudo, J. Yoshida, *Tetrahedron Lett.* **1999**, *40*, 5533; b) K. Itami, K. Mitsudo, J. Yoshida, *Tetrahedron Lett.* **1999**, *40*, 5537; c) K. Itami, T. Nokami, J. Yoshida, *Org. Lett.* **2000**, *2*, 1299.
- [4] K. Itami, K. Mitsudo, J. Yoshida, *J. Org. Chem.* **1999**, *64*, 8709.
- [5] For reviews on micellar organic reactions, see a) S. Tascioglu, *Tetrahedron* **1996**, *52*, 11 113; b) J. B. F. N. Engberts, *Pure Appl. Chem.* **1992**, *64*, 1653.
- [6] a) D. C. Rideout, R. Breslow, *J. Am. Chem. Soc.* **1980**, *102*, 7816; b) R. Breslow, U. Maitra, *Tetrahedron Lett.* **1984**, *25*, 1239.
- [7] a) P. A. Grieco, P. Garner, Z. He, *Tetrahedron Lett.* **1983**, *24*, 1897; b) P. A. Grieco, K. Yoshida, P. Garner, *J. Org. Chem.* **1983**, *48*, 3137; c) P. A. Grieco, P. Galatsis, R. F. Spohn, *Tetrahedron* **1986**, *42*, 2847.

- [8] a) A. Lubineau, Y. Queneau, *J. Org. Chem.* **1987**, *52*, 1001; b) D. A. Jaeger, H. Shinozaki, P. A. Goodson, *J. Org. Chem.* **1991**, *56*, 2482; c) W. Blokzijl, M. J. Blandamer, J. B. F. N. Engberts, *J. Am. Chem. Soc.* **1991**, *113*, 4241.
- [9] For recent examples, see a) Z. Zhu, J. H. Espenson, *J. Am. Chem. Soc.* **1997**, *119*, 3507; b) S. Otto, J. B. F. N. Engberts, *J. Am. Chem. Soc.* **1999**, *121*, 6798; c) K. Manabe, Y. Mori, S. Kobayashi, *Tetrahedron* **1999**, *55*, 11 203; d) D. A. Jaeger, D. Su, A. Zafar, B. Pikhova, S. B. Hall, *J. Am. Chem. Soc.* **2000**, *122*, 2749.
- [10] a) R. Breslow, *Acc. Chem. Res.* **1991**, *24*, 159; b) W. Blokzijl, J. B. F. N. Engberts, *Angew. Chem.* **1993**, *105*, 1610; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1545.
- [11] a) R. Braun, F. Schuster, J. Sauer, *Tetrahedron Lett.* **1986**, *27*, 1285; b) V. K. Singh, B. N. S. Raju, P. T. Deota, *Synth. Commun.* **1988**, *18*, 567; c) N. K. Sangwan, H.-J. Schneider, *J. Chem. Soc. Perkin Trans. 2* **1989**, 1223; d) I. Hunt, C. D. Johnson, *J. Chem. Soc. Perkin Trans. 2* **1991**, 1051; e) G. K. van der Wel, J. W. Wijnen, J. B. F. N. Engberts, *J. Org. Chem.* **1996**, *61*, 9001; f) J. W. Wijnen, J. B. F. N. Engberts, *J. Org. Chem.* **1997**, *62*, 2039; g) K. Chiba, M. Jinno, A. Nozaki, M. Tada, *Chem. Commun.* **1997**, 1403; h) M. J. Diego-Castro, H. C. Hailes, *Tetrahedron Lett.* **1998**, *39*, 2211; i) S. Otto, J. B. F. N. Engberts, J. C. T. Kwak, *J. Am. Chem. Soc.* **1998**, *120*, 9517.
- [12] The cycloadduct formed initially was easily air-oxidized to **8** when subjected to chromatography on silica gel.

The Combination of Spontaneous Resolution and Asymmetric Catalysis: A Model for the Generation of Optical Activity from a Fully Racemic System**

Olivier Tissot, Maryse Gouygou,* Frédéric Dallemer, Jean-Claude Daran, and Gilbert G. A. Balavoine*

Dedicated to Professor Henri Kagan on the occasion of his 70th birthday

Considerable efforts have been devoted to the development of new chiral ligands owing to the growing importance of transition metal catalyzed asymmetric synthesis.^[1] Among these chiral ligands, diphosphanes have played a dominant role, in particular those possessing C₂ symmetry.^[2] We were interested in the possibilities offered by 1,1'-diphenyl-3,3',4,4'-tetramethyl-2,2'-biphosphole (**1**) (BIPHOS), first synthesized by Mathey et al. in 1986.^[3] This diphosphane combines the axial chirality generated by the biphenyl framework with the central chiralities of the phosphorus atoms. This implies the existence of six stereoisomers, corresponding to three pairs of enantiomers, which are in a fast equilibrium in

[*] Dr. M. Gouygou, Prof. G. G. A. Balavoine, Dr. O. Tissot, Dr. J.-C. Daran
Laboratoire de Chimie de Coordination du CNRS
205, Route de Narbonne, 31077 Toulouse Cedex (France)
Fax: (+33) 5-61-55-30-03
E-mail: gouygou@lcc-toulouse.fr, balavoine@lcc-toulouse.fr
Dr. F. Dallemer
Rhodia Recherches, Centre de Recherche de Lyon
85, Avenue des Freres Perret, BP 62, 69192 Saint-Fons Cedex (France)

[**] We thank the Centre National de la Recherche Scientifique and Rhodia for financial support, Prof. H. B. Kagan for helpful discussions, and H. Ait-Haddou and O. Hoarau for assistance in catalytic tests.