




Scheme 1. Aqueous Diels–Alder reaction of **3** and removal of the 2-PyMe₂Si group. Bn = benzyl.

ected throughout these transformations, which indicates the reasonable chemical stability of this group. Finally, the oxidation of **10** with H₂O₂ afforded **11** in 89% yield without affecting the C–C double bond or eroding the stereochemistry at the allylic carbon atoms.

In summary, we have demonstrated the “proof-of-principle” of our strategy for aqueous organic reactions by utilizing the 2-PyMe₂Si group as a removable hydrophilic group. Importantly, the strategy described herein should not be limited to aqueous Diels–Alder reactions but could in principle be applied to other aqueous organic reactions as well.

Received: September 27, 2000 [Z15862]

- [1] a) *Organic Synthesis in Water* (Ed.: P. A. Grieco), Blackie Academic & Professional, London, **1998**; b) C.-J. Li, T.-H. Chen, *Organic Reactions in Aqueous Media*, Wiley, New York, **1997**.
- [2] J. Yoshida, K. Itami, K. Mitsudo, S. Suga, *Tetrahedron Lett.* **1999**, *40*, 3403.
- [3] a) K. Itami, K. Mitsudo, J. Yoshida, *Tetrahedron Lett.* **1999**, *40*, 5533; b) K. Itami, K. Mitsudo, J. Yoshida, *Tetrahedron Lett.* **1999**, *40*, 5537; c) K. Itami, T. Nokami, J. Yoshida, *Org. Lett.* **2000**, *2*, 1299.
- [4] K. Itami, K. Mitsudo, J. Yoshida, *J. Org. Chem.* **1999**, *64*, 8709.
- [5] For reviews on micellar organic reactions, see a) S. Tascioglu, *Tetrahedron* **1996**, *52*, 11 113; b) J. B. F. N. Engberts, *Pure Appl. Chem.* **1992**, *64*, 1653.
- [6] a) D. C. Rideout, R. Breslow, *J. Am. Chem. Soc.* **1980**, *102*, 7816; b) R. Breslow, U. Maitra, *Tetrahedron Lett.* **1984**, *25*, 1239.
- [7] a) P. A. Grieco, P. Garner, Z. He, *Tetrahedron Lett.* **1983**, *24*, 1897; b) P. A. Grieco, K. Yoshida, P. Garner, *J. Org. Chem.* **1983**, *48*, 3137; c) P. A. Grieco, P. Galatsis, R. F. Spohn, *Tetrahedron* **1986**, *42*, 2847.

- [8] a) A. Lubineau, Y. Queneau, *J. Org. Chem.* **1987**, *52*, 1001; b) D. A. Jaeger, H. Shinozaki, P. A. Goodson, *J. Org. Chem.* **1991**, *56*, 2482; c) W. Blokzijl, M. J. Blandamer, J. B. F. N. Engberts, *J. Am. Chem. Soc.* **1991**, *113*, 4241.
- [9] For recent examples, see a) Z. Zhu, J. H. Espenson, *J. Am. Chem. Soc.* **1997**, *119*, 3507; b) S. Otto, J. B. F. N. Engberts, *J. Am. Chem. Soc.* **1999**, *121*, 6798; c) K. Manabe, Y. Mori, S. Kobayashi, *Tetrahedron* **1999**, *55*, 11 203; d) D. A. Jaeger, D. Su, A. Zafar, B. Pikhova, S. B. Hall, *J. Am. Chem. Soc.* **2000**, *122*, 2749.
- [10] a) R. Breslow, *Acc. Chem. Res.* **1991**, *24*, 159; b) W. Blokzijl, J. B. F. N. Engberts, *Angew. Chem.* **1993**, *105*, 1610; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1545.
- [11] a) R. Braun, F. Schuster, J. Sauer, *Tetrahedron Lett.* **1986**, *27*, 1285; b) V. K. Singh, B. N. S. Raju, P. T. Deota, *Synth. Commun.* **1988**, *18*, 567; c) N. K. Sangwan, H.-J. Schneider, *J. Chem. Soc. Perkin Trans. 2* **1989**, 1223; d) I. Hunt, C. D. Johnson, *J. Chem. Soc. Perkin Trans. 2* **1991**, 1051; e) G. K. van der Wel, J. W. Wijnen, J. B. F. N. Engberts, *J. Org. Chem.* **1996**, *61*, 9001; f) J. W. Wijnen, J. B. F. N. Engberts, *J. Org. Chem.* **1997**, *62*, 2039; g) K. Chiba, M. Jinno, A. Nozaki, M. Tada, *Chem. Commun.* **1997**, 1403; h) M. J. Diego-Castro, H. C. Hailes, *Tetrahedron Lett.* **1998**, *39*, 2211; i) S. Otto, J. B. F. N. Engberts, J. C. T. Kwak, *J. Am. Chem. Soc.* **1998**, *120*, 9517.
- [12] The cycloadduct formed initially was easily air-oxidized to **8** when subjected to chromatography on silica gel.

The Combination of Spontaneous Resolution and Asymmetric Catalysis: A Model for the Generation of Optical Activity from a Fully Racemic System**

Olivier Tissot, Maryse Gouygou,* Frédéric Dallemer, Jean-Claude Daran, and Gilbert G. A. Balavoine*

Dedicated to Professor Henri Kagan on the occasion of his 70th birthday

Considerable efforts have been devoted to the development of new chiral ligands owing to the growing importance of transition metal catalyzed asymmetric synthesis.^[1] Among these chiral ligands, diphosphanes have played a dominant role, in particular those possessing C₂ symmetry.^[2] We were interested in the possibilities offered by 1,1'-diphenyl-3,3',4,4'-tetramethyl-2,2'-biphosphole (**1**) (BIPHOS), first synthesized by Mathey et al. in 1986.^[3] This diphosphane combines the axial chirality generated by the biphenyl framework with the central chiralities of the phosphorus atoms. This implies the existence of six stereoisomers, corresponding to three pairs of enantiomers, which are in a fast equilibrium in

[*] Dr. M. Gouygou, Prof. G. G. A. Balavoine, Dr. O. Tissot, Dr. J.-C. Daran
Laboratoire de Chimie de Coordination du CNRS
205, Route de Narbonne, 31077 Toulouse Cedex (France)
Fax: (+33) 5-61-55-30-03
E-mail: gouygou@lcc-toulouse.fr, balavoine@lcc-toulouse.fr

Dr. F. Dallemer
Rhodia Recherches, Centre de Recherche de Lyon
85, Avenue des Freres Perret, BP 62, 69192 Saint-Fons Cedex (France)

[**] We thank the Centre National de la Recherche Scientifique and Rhodia for financial support, Prof. H. B. Kagan for helpful discussions, and H. Ait-Haddou and O. Hoarau for assistance in catalytic tests.

Table 1. Asymmetric allylic substitution of 1,3-diphenylprop-2-enyl acetate with the anion of dimethyl malonate, using 1 mol % of the palladium complex **2**.

Entry	Catalyst	Solvent	<i>T</i> [°C]	Conversion [% of 3] ^[a]	<i>ee</i> [% of 4] ^[b]
1	<i>R</i> [SS]-(-)- 2	CH ₂ Cl ₂	37	84	80 (<i>R</i>)
2	<i>R</i> [SS]-(-)- 2	THF	25	93	80 (<i>R</i>)
3	<i>S</i> [RR]-(+)- 2	THF	25	93	80 (<i>S</i>)

[a] Evaluated by ¹H NMR spectroscopy. [b] Determined by HPLC analysis using a chiral column (Pharmacir 7C).

as effective as CHIRAPHOS,^[10a] BPPF,^[10b] DPPBA,^[10c] and the BIPNOR^[10d]/Pd^{II} system in the catalytic asymmetric allylic substitution of 1,3-diphenylprop-2-enyl acetate with the anion of dimethyl malonate (Table 1). Our results provide an interesting example of the spontaneous generation of optical activity from a racemic mixture by crystallization of a molecule that is stereolabile in solution, followed by amplification in an enantioselective catalytic process. The crystallization of chiral ligands or chiral catalytic complexes as conglomerates has already been described,^[11] but to the best of our knowledge, the stereolabile BIPHOS ligand appears to be a special case. Because of the fast equilibrium between the two enantiomers in solution, the whole BIPHOS racemic mixture can be transformed into a single enantiomer,^[12] giving the enantiomerically pure catalytic complex quantitatively.

Experimental Section

1: The biphosphole was prepared as described in the literature.^[2b] To obtain single crystals, a very high purity of **1** is crucial. After purification by filtration through a short pad of silica gel (dichloromethane/pentane 70:30), the resulting oily residue was washed with pentane until a pale yellow solid was obtained. Subsequently, the slow evaporation of an undisturbed dichloromethane solution under argon gave single crystals of **1**.

2: A single crystal of BIPHOS **1** (0.050 mg, 0.13 mmol) was added to a stirred suspension of [PdCl₂(CH₃CN)₂] (0.033 mg, 0.13 mmol) in dichloromethane (5 mL) maintained at -78 °C. The mixture was stirred at -78 °C for 16 h, and then warmed to room temperature and filtered. The slow addition of pentane to the solution gave complex **2** as an orange microcrystalline solid (0.067 mg, 93 %), which was washed with pentane (2 × 2 mL). [α]_D²⁰ = +200 (*c* = 0.02 in CH₂Cl₂); ¹H NMR (CD₂Cl₂): δ = 1.87 (d, ⁴*J*_{PH} = 2.9 Hz, 6H; Me21, 121), 2.14 (d, ⁴*J*_{HH} = 1.5 Hz, 6H; Me31, 131), 6.63 (m, 2H; =CH-P), 7.40–7.87 (m, 10H; Ph); ³¹P NMR (CD₂Cl₂): δ = 33.4. Single crystals of *S*[RR]-(+)-**2** were obtained by the slow evaporation of a dichloromethane solution. Complex *R*[SS]-(-)-**2** was obtained in a similar way, starting from a single crystal of BIPHOS *R*[SS]-(-)-**1**.

General procedure for the allylic substitution: A solution of sodium dimethyl malonate resulting from the treatment of dimethyl malonate (0.148 mL, 1.29 mmol, 1.2 equiv) in THF (4 mL) with sodium hydride (80 % NaH, 1.07 mmol) was added to a solution of complex **2** (7 mg, 0.0123 mmol) and 1,3-diphenylprop-2-enylacetate (310 mg, 1.23 mmol) in THF (2 mL). After 24 h at room temperature, the resulting mixture was diluted with diethyl ether (5 mL) and quenched with a saturated aqueous NH₄Cl solution (5 mL). The aqueous phase was extracted with diethyl ether, the combined organic phases were dried over MgSO₄, filtered, and evaporated to give the crude product, which was purified by chromatography (SiO₂, ethyl acetate/pentane = 15:85). The enantiomeric excess was determined by chiral HPLC using a Pharmacir 7C column, hexane/isopropanol (90:10), 0.7 mL min⁻¹, *R_f* for the *S* isomer = 13.84 min, *R_f* for the *R* isomer = 15.15 min.

Received: October 27, 2000 [Z16001]

- [1] For a recent review, see: *Comprehensive Asymmetric Catalysis* (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, 1999.
- [2] a) T. P. Dang, H. B. Kagan, *J. Chem. Soc. Chem. Commun.* **1971**, 2287; b) J. K. Whitesell, *Chem. Rev.* **1989**, 89, 1581, and references therein.
- [3] a) F. Mercier, S. Holand, F. Mathey, *J. Organomet. Chem.* **1986**, 316, 271; b) E. Deschamps, F. Mathey *Bull. Soc. Chim. Fr.* **1992**, 129.
- [4] O. Tissot, M. Gouygou, J.-C. Daran, G. G. A. Balavoine, *Chem. Commun.* **1996**, 2287.
- [5] O. Tissot, Doctoral Thesis, University Paul Sabatier, Toulouse, 1999.
- [6] To define the absolute configuration, the axial chirality is given first and the central chiralities are given between square brackets.
- [7] M. Gouygou, O. Tissot, J.-C. Daran, G. G. A. Balavoine, *Organometallics* **1997**, 16, 1008.
- [8] Crystal data for *S*[RR]-(+)-**2** (C₂₄H₂₄P₂Cl₂Pd): *M_r* = 551, monoclinic, space group *C*₂, *a* = 18.485(3), *b* = 10.2561(2), *c* = 12.975(2) Å, β = 105.13(2)°, *V* = 2374.6(5) Å³, *Z* = 4, ρ_{calcd} = 1.543 g cm⁻³, μ = 11.396 cm⁻¹, MoK α radiation, *T* = 180 K, φ scan (Stoe IPDS diffractometer), $2\theta_{\text{max}}$ = 48.3°, reflections collected 9335, unique 3675 (*R*_{int} = 0.0525), observed 3424 (*I* > 2 σ (*I*)), parameters refined 264, *R*/*R_w* = 0.0289/0.0337, GOF = 1.044, $\Delta/\sigma_{\text{max}}$ = 0.025, [$\Delta\rho$]_{min} – 0.30, [$\Delta\rho$]_{max} 0.47, Flack's parameter = –0.04(2). Structure solution (SIR97) and refinement (CRYSTALS). Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-150425. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [9] B. M. Trost, D. L. Van Vranken, *Chem. Rev.* **1996**, 96, 395.
- [10] a) CHIRAPHOS = 1,2-dimethyl-1,2-ethanediylbis[(diphenyl)phosphine]; P. B. Mackenzie, J. Whelan, B. Bosnich, *J. Am. Chem. Soc.* **1985**, 107, 2046; b) BPPF = 1-[1-(dimethylamino)ethyl]-1',2'-bis(diphenylphosphanyl)ferrocene; T. Hayashi, A. Yamamoto, T. Hagihara, Y. Ito, *Tetrahedron Lett.* **1986**, 27, 191; c) DPPBA = 2-(diphenylphosphanyl)benzoic acid; B. M. Trost, R. C. Bunt, *J. Am. Chem. Soc.* **1994**, 116, 4089; d) BIPNOR = 2,2',3,3'-tetraphenyl-4,4',5,5'-tetramethyl-6,6'-bis-1-phosphanorborna-2,5-dienyl; F. Robin, F. Mercier, L. Ricard, F. Mathey, *Chem. Eur. J.* **1997**, 3, 1365.
- [11] D. Carmona, C. Cativela, S. Elipe, F. J. Lahoz, M. P. Lamata, M. P. Lopez-Ram de Viu, L. A. Oro, C. Vega, F. Viguri, *Chem. Commun.* **1997**, 2351.
- [12] The spontaneous resolution of a stereolabile racemic mixture has been described for binaphthyl; see: a) R. E. Pincock, R. R. Perkins, A. S. Ma, K. R. Wilson, *Science* **1971**, 1081; b) D. K. Kondepudi, J. Laudadio, K. Asakura, *J. Am. Chem. Soc.* **1999**, 121, 1448.