


Synthesis, Structure, and Properties of a Novel Heterooctametallic Complex Containing a Cyclic Ru₄Ni₄ Core**

Indrani Pal, Falguni Basuli, Thomas C. W. Mak, and Samaresh Bhattacharya*


There is considerable current interest in the chemistry of polynuclear complexes, largely because of their fascinating properties.^[1] Mononuclear transition metal complexes of multidentate ligands in which some of the donor sites are unoccupied often serve as efficient building blocks for the construction of polynuclear assemblies.^[2] In our attempts to generate new heteropolynuclear systems, we planned a stepwise synthetic strategy, and salicylaldehyde thiosemicarbazone (H₃saltsc, **1**) was chosen as the multidentate ligand for this study. This ligand has five potential donor sites: three N, one O, and one S atom. However, in its reaction with ruthenium and osmium, we observed an apparently unusual coordination mode in which the H₂saltsc ligand utilizes only one N and one S donor site and forms a four-membered chelate ring (**2**).^[3] Further studies showed that this type of



coordination mode is in fact quite normal not only for H₂saltsc but also for benzaldehyde thiosemicarbazones.^[4] The three unused donor sites in **2** and their relative dispositions in space suggested that complexes containing this moiety should be able to bind to a second metal ion as a tridentate N,N,O donor ligand, and such a possibility is explored in the present study.

The mononuclear mixed-ligand ruthenium(II) complex [Ru(bpy)₂(H₂saltsc)]⁺ (bpy = 2,2'-bipyridine) was prepared first. 2,2'-Bipyridine was chosen as the coligand because it is a recognized stabilizer of bivalent ruthenium, and complexes containing the Ru(bpy)₂ chromophore exhibit interesting luminescence properties,^[5] so that a polynuclear complex constructed from [Ru(bpy)₂(H₂saltsc)]⁺ building blocks may also be expected to display such behavior. [Ru(bpy)₂-

(H₂saltsc)]ClO₄ was obtained from the reaction of H₃saltsc with [Ru(bpy)₂(EtOH)₂]²⁺, formed in situ from [Ru(bpy)₂Cl₂] and AgNO₃ in warm ethanol in the presence of a base [Eq. (1)]. The crystal structure of [Ru(bpy)₂(H₂saltsc)]ClO₄ is



displayed in Figure 1.^[6] The H₂saltsc ligand is coordinated to ruthenium as a bidentate N,S donor in a four-membered chelate ring. The N₃S coordination sphere around ruthenium





Figure 1. ORTEP plot of the structure of [Ru(bpy)₂(H₂saltsc)]⁺. Selected bond lengths [Å] and angles [°]: Ru1-S1 2.430(5), Ru1-N6 2.068(12), S1-C28 1.70(2), C28-N7 1.33(2), C28-N6 1.41(2), N6-N5 1.41(2), N5-C27 1.24(2), C27-C26 1.47(3), C26-C21 1.40(4), C21-O1 1.28(4); N6-Ru1-S1 66.7(5), Ru1-S1-C28 82.6(7), N6-C28-S1 107.1(15), N7-C28-S1 125.6(17), C28-N6-N5 118.5(17), N6-N5-C27 119.3(19), N5-C27-C26 123(2), C27-C26-C21 121(3), O1-C21-C26 123(3).

is distorted from ideal octahedral geometry. The phenolic oxygen atom O1, the imine nitrogen atom N5, and the amine nitrogen atom N7 of the H₂saltsc ligand remained uncoordinated, and the Ru(H₂saltsc) fragment is almost planar. The [Ru(bpy)₂(H₂saltsc)]⁺ complex cation thus appears to be a suitable building block for the construction of polynuclear complexes, and to test this hypothesis, [Ru(bpy)₂(H₂saltsc)]ClO₄ was treated with one equivalent of Ni(ClO₄)₂ · 6H₂O in warm acetonitrile/ethanol. This reaction indeed afforded a polynuclear complex, namely [{Ru(bpy)₂(saltsc)}₄Ni₄](ClO₄)₄ [Eq. (2)].



Formation of the centrosymmetric, octametallic complex cation was confirmed by X-ray crystallography (Figure 2).^[7] The coordinated H₂saltsc ligand of [Ru(bpy)₂(H₂saltsc)]ClO₄ has lost two further protons, one from the phenolic OH and the other from the NH₂ group, and the resulting saltsc ligand exhibits N,N,O coordination to the nickel(II) ion. The fourth coordination site of the nickel ion is occupied by the sulfur

[*] Dr. S. Bhattacharya, I. Pal, F. Basuli
Department of Chemistry
Inorganic Chemistry Section
Jadavpur University
Kolkata 700032 (India)
Fax: (+91)33-473-4266
E-mail: samaresh_b@hotmail.com

Prof. Dr. T. C. W. Mak
Department of Chemistry
The Chinese University of Hong Kong
Shatin, New Territories (Hong Kong)

[**] This work was supported by a grant (Grant No. SP/S1/F33/98) from the Department of Science and Technology, New Delhi, India and Hong Kong Research Grants Council Earmarked Grant CUHK 4022/98P.




Figure 2. ORTEP plot of the structure of $[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4]^{4+}$. Selected bond lengths [Å] and angles [°]: Ru1-S1 2.450(2), Ru1-N6 2.0936, S1-C28 1.789(8), C28-N7 1.303(9), C28-N6 1.319(10), N6-N5 1.390(9), N5-C27 1.301(10), C27-C26 1.406(12), C26-C21 1.402(12), C21-O1 1.321(10), Ni1-O1 1.831(6), Ni1-N7 1.868(7), Ni1-N5 1.853(6), Ni1-S2 2.213(2); N6-Ru1-S1 66.55(19), Ru1-S1-C28 80.0(3), N6-C28-S1 106.8(5), N7-C28-S1 132.2(6), C28-N6-N5 110.4(6), N6-N5-C27 119.5(7), N5-C27-C26 125.0(8), C27-C26-C21 122.3(8), O1-C21-C26 123.2(7), Ni1-O1-C21 128.3(5), O1-Ni1-N5 94.3(3), N5-Ni1-N7 83.6(3), Ni1-N7-C28 111.2(5), O1-Ni1-S2 83.53(18).

atom of another $\text{Ru}(\text{bpy})_2(\text{saltsc})\text{Ni}$ fragment, and the bridging sulfur atoms result in the formation of the cyclic octametallic $[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4]^{4+}$ complex cation. It is noteworthy that in this complex all five available donor sites of the thiosemicarbazone ligand are involved in coordination, and this is unprecedented in the literature. The Ni_2OS coordination sphere around nickel is distorted from ideal square-planar geometry. While bond lengths within the NiN_2OS core are quite normal,^[8] those in the $\text{Ru}(\text{saltsc})$ fragment of $[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$ are slightly different to the corresponding bond lengths in the $\text{Ru}(\text{H}_2\text{saltsc})$ fragment of $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ owing to coordination to nickel. The significant elongation of the C28–S1 bond is attributable to the bridging nature of the sulfur atom.

Both the mononuclear and octametallic complexes are diamagnetic; this indicates the presence of Ru^{II} (low-spin d^6 , $S=0$) in both of them, and of Ni^{II} (square-planar d^8 , $S=0$) in the octametallic complex. The ^1H NMR, conductance, and analytical data of the two complexes support their formulations. The cyclic voltammograms of $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ and $[\text{Ru}\{\text{bpy}\}_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$ recorded in $(\text{Et}_4\text{N})\text{ClO}_4/\text{MeCN}$ are shown in Figure 3. For $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ two oxidation waves are observed on the positive side of the saturated calomel electrode (SCE), and four successive reduction waves on the negative side. The first oxidation is reversible ($E_{1/2} = 0.62$ V, $\Delta E_p = 60$ mV, $i_{pa} = i_{pc}$) and is assigned to $\text{Ru}^{\text{II}} \rightarrow \text{Ru}^{\text{III}}$ oxidation. The second oxidation is irreversible ($E_{pa} = 1.38$ V) and is probably ligand-centered. The reduction waves ($E_{pc} = -1.50$, -1.70 , -1.87 , and -2.39 V) are due to




Figure 3. Cyclic voltammograms of a) $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ and b) $[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$ in acetonitrile solution (0.1M $(\text{Et}_4\text{N})\text{ClO}_4$) at a scan rate of 50 mV s^{-1} .

reductions of the two coordinated bpy ligands. Each bpy ligand is known to undergo two one-electron reductions.^[9] Hence, four successive one-electron reductions are expected for $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]^+$, and all of them were observed experimentally. Cyclic voltammetric behavior of $[\text{Ru}\{\text{bpy}\}_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$ is qualitatively similar to that of $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$. However, both oxidations are shifted to more positive values. The first oxidation is now irreversible ($E_{1/2} = 0.75$ V, $i_{pa} > i_{pc}$), and the second remained irreversible ($E_{pa} = 1.56$ V). All four reductions of the two bpy ligands in the $\text{Ru}(\text{bpy})_2$ fragments were also observed ($E_{pc} = -1.55$, -1.81 , -2.06 , and -2.37 V) in this complex. The cyclic voltammetric results clearly indicate that there are no detectable metal–metal interactions in the octametallic complex.

Experimental Section

$[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$: AgNO_3 (65 mg, 0.38 mmol) was added to a solution of $[\text{Ru}(\text{bpy})_2\text{Cl}_2] \cdot 2\text{H}_2\text{O}$ (100 mg, 0.19 mmol) in ethanol (30 mL). The mixture was warmed and stirred for 30 min. The deposited AgCl was separated by filtration, and H_2saltsc (40 mg, 0.21 mmol) and NEt_3 (21 mg, 0.21 mmol) were added to the filtrate. The solution was heated to reflux for 2 h. It was then concentrated to ca. 10 mL, and a saturated aqueous solution of NaClO_4 (0.5 mL) was added. $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ separated as a dark precipitate, which was collected by filtration, washed with water, and dried in vacuo over P_4O_{10} . Recrystallization from acetonitrile/benzene gave 125 mg of $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ (92%) as a brownish red crystalline solid. Elemental analysis (%) calcd for $\text{C}_{28}\text{H}_{24}\text{N}_7\text{O}_5\text{SClRu}$: C 47.56, H 3.40, N 13.87; found: C 47.69, H 3.49, N 13.65.

$[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$: $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4$ (100 mg, 0.14 mmol) was dissolved in warm ethanol (30 mL), and NEt_3 (14 mg, 0.14 mmol) and a solution of $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (52 mg, 0.14 mmol) in acetonitrile (10 mL) were added to the stirred solution. The mixture was warmed and stirred for 5 h to afford 75 mg of $[\text{Ru}(\text{bpy})_2(\text{saltsc})_4\text{Ni}_4](\text{ClO}_4)_4$ (66%), which separated as a brown microcrystalline solid and was collected by filtration, washed with ethanol, and dried in air. Elemental analysis (%) calcd for $\text{C}_{28}\text{H}_{22}\text{N}_7\text{O}_5\text{SClRuNi} \cdot 2\text{H}_2\text{O}$: C 42.04, H 3.25, N 12.26; found: C 41.51, H 3.35, N 11.70.

Received: March 9, 2001
Revised: May 22, 2001 [Z16749]


- [1] a) V. Balzani, A. Juris, M. Venturi, S. Campagna, S. S. Wilcox, *Chem. Rev.* **1996**, 96, 759–833; b) F. C. Anson, C. Shi, B. Steiger, *Acc. Chem. Res.* **1997**, 30, 437–444; c) V. Balzani, S. Campagna, G. Denti, A. Juris, S. Serroni, M. Venturi, *Acc. Chem. Res.* **1998**, 31, 26–34; d) J. A. McCleverty, M. D. Ward, *Acc. Chem. Res.* **1998**, 31, 842–851; e) P. C. Ford, E. Cariati, J. Bourassa, *Chem. Rev.* **1999**, 99, 3625–3647; f) Y. Xiong, L. N. Ji, *Coord. Chem. Rev.* **1999**, 185/186, 711–733; g) H. E. Toma, K. Araki, *Coord. Chem. Rev.* **2000**, 196, 307–329; h) D. T. McQuade, A. E. Pullen, T. M. Swager, *Chem. Rev.* **2000**, 100, 2537–2574; i) A. D. Cutland, R. G. Makkani, J. W. Kampf, V. L. Pecoraro, *Angew. Chem.* **2000**, 112, 2801–2803; *Angew. Chem. Int. Ed.* **2000**, 39, 2689–2691; j) J. J. Bodwin, V. L. Pecoraro, *Inorg. Chem.* **2000**, 39, 3434–3435; k) T. Konno, Y. Chikamoto, K. I. Okamoto, T. Yamaguchi, T. Ito, M. Hirotsu, *Angew. Chem.* **2000**, 112, 4264–4267; *Angew. Chem. Int. Ed.* **2000**, 39, 4098–4101; l) M. M. Ali, F. M. MacDonnell, *J. Am. Chem. Soc.* **2000**, 122, 11527–11528; m) R. W. Saalfrank, S. Trummer, U. Reimann, M. M. Chowdhury, F. Hampel, O. Waldmann, *Angew. Chem.* **2000**, 112, 3634–3636; *Angew. Chem. Int. Ed.* **2000**, 39, 3492–3494; Erratum: R. W. Saalfrank, S. Trummer, U. Reimann, M. M. Chowdhury, F. Hampel, O. Waldmann, *Angew. Chem.* **2000**, 112, 3885; *Angew. Chem. Int. Ed.* **2000**, 39, 3736; n) E. Uller, B. Demleitner, I. Berndt, R. W. Saalfrank, *Struct. Bonding (Berlin)* **2000**, 96, 149–175.
- [2] a) M. D. Ward, *Inorg. Chem.* **1996**, 35, 1712–1714; b) F. Birkelbach, T. Weyhemüller, M. Lengen, M. Gardan, A. X. Trautwein, K. Wieghardt, P. Chaudhuri, *J. Chem. Soc. Dalton Trans.* **1997**, 4529–4537; c) W. Paw, R. Eisenberg, *Inorg. Chem.* **1997**, 36, 2287–2293; d) P. Majumdar, S. M. Peng, S. Goswami, *J. Chem. Soc. Dalton Trans.* **1998**, 1569–1574; e) G. A. Abakumov, V. K. Cherkasov, V. I. Nevodchikov, V. A. Kurapatov, B. C. Noll, C. G. Pierpont, *Inorg. Chem.* **1998**, 37, 6117–6119; f) A. K. Das, A. Rueda, L. R. Falvello, S. M. Peng, S. Bhattacharya, *Inorg. Chem.* **1999**, 38, 4365–4368; g) N. C. Fletcher, T. C. Robinson, A. Behrendt, J. C. Jeffery, Z. R. Reeves, M. D. Ward, *J. Chem. Soc. Dalton Trans.* **1999**, 2999–3006.
- [3] F. Basuli, S. M. Peng, S. Bhattacharya, *Inorg. Chem.* **1997**, 36, 5645–5647.
- [4] a) F. Basuli, M. Ruf, C. G. Pierpont, S. Bhattacharya, *Inorg. Chem.* **1998**, 37, 6113–6116; b) F. Basuli, S. M. Peng, S. Bhattacharya, *Inorg. Chem.* **2000**, 39, 1120–1127.
- [5] A. Juris, S. B. S. Campagna, V. Balzani, P. Belser, A. V. Zelewsky, *Coord. Chem. Rev.* **1988**, 84, 85–227.
- [6] a) X-ray structure analysis of $[\text{Ru}(\text{bpy})_2(\text{H}_2\text{saltsc})]\text{ClO}_4 \cdot 1/2 \text{C}_6\text{H}_6 \cdot 1/2 \text{CH}_2\text{Cl}_2$: $\text{C}_{31.50}\text{H}_{27}\text{Cl}_2\text{N}_7\text{O}_5\text{RuS}$, $M_r = 787.63$, monoclinic, space group $P2_1/c$, $a = 9.169(1)$, $b = 16.396(2)$, $c = 24.634(3)$ Å, $\beta = 99.075(8)^\circ$, $V = 3657.0(7)$ Å³, $Z = 4$, $\rho_{\text{calc}} = 1.431 \text{ Mg m}^{-3}$; crystal dimensions $0.50 \times 0.30 \times 0.02$ mm. Data ($2\theta_{\text{max}} = 51.1^\circ$) were collected at 293(2) K on a Rigaku RAXIS-IIIC imaging plate with graphite-monochromatized MoK_α radiation ($\lambda = 0.71073$ Å), $\mu = 0.597 \text{ mm}^{-1}$, transmission factors 0.755–1.148. The structure was solved by direct methods and refined by full-matrix least-squares methods on F^2 . Hydrogen atoms were included but not refined. $R1 = 0.1267$ for 3296 observed reflections ($I > 2\sigma(I)$) and 418 parameters; $wR2 = 0.3507$ for 4634 unique reflections. Programs used: SHELXS-97 (structure solution) and SHELXL-97 (structure refinement). b) Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-156075 and -156076. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [7] X-ray structure analysis of $[\{\text{Ru}(\text{bpy})_2(\text{saltsc})\}_4\text{Ni}_4](\text{ClO}_4)_4 \cdot 4 \text{C}_6\text{H}_6 \cdot 2 \text{MeCN}$: $\text{C}_{140}\text{H}_{118}\text{Cl}_4\text{N}_{30}\text{Ni}_4\text{O}_{20}\text{Ru}_4\text{S}_4$, $M_r = 3449.80$, monoclinic, space group $P2_1n$, $a = 18.386(3)$, $b = 12.268(2)$, $c = 36.398(6)$ Å, $\beta = 104.219(3)^\circ$, $V = 7959(2)$ Å³, $Z = 2$; crystal dimensions $0.60 \times 0.47 \times 0.28$ mm, intensities ($2\theta_{\text{max}} = 56.2^\circ$) were collected at 293(2) K on a Bruker SMART 1000 CCD diffractometer with MoK_α radiation ($\lambda = 0.71073$ Å), $\mu = 1.020 \text{ mm}^{-1}$; absorption corrections with the SADABS program yielded relative transmission factors of 0.8075–1. $R1 = 0.0795$ for 10301 observed reflections ($I > 2\sigma(I)$) and 932 parameters; $wR2 = 0.2672$ for 19288 reflections.^[6b]
- [8] a) S. Padhye, *Coord. Chem. Rev.* **1985**, 63, 127–160; b) M. Mathew, G. J. Palenik, G. R. Clark, *Inorg. Chem.* **1973**, 12, 446–451.
- [9] a) A. A. Vlcek, *Coord. Chem. Rev.* **1982**, 43, 39–62; b) J. L. Kahl, K. W. Hanck, K. DeArmond, *J. Phys. Chem.* **1978**, 82, 540–545.

Selective Photoproduction of O₂ from the Mn₄O₄ Cubane Core: A Structural and Functional Model for the Photosynthetic Water-Oxidizing Complex**

Masayuki Yagi, Kurt V. Wolf, Patrick J. Baesjou, Steven L. Bernasek, and G. Charles Dismukes*

Dedicated to Dr. Peter Gabriel

Both the announcement last year by the German Federal Government to eliminate the use of nuclear power generators by 2021^[1] and the recent shortage of electric power in California USA place renewed emphasis on development of socially acceptable energy sources such as solar^[2] and fuel cells.^[3] The latter cells rely on energy generated by the combination of H₂ (or hydrocarbons) with O₂. Both of these molecules are expected to be produced by solar-based water-splitting catalysts ($\text{H}_2\text{O} \rightarrow \frac{1}{2}\text{O}_2 + \text{H}_2$), hence, the efforts to understand Nature's photosynthetic process of O₂ generation by water oxidation in plants,^[4] and to functionally mimic the catalytic center,^[5,6] take on a pressing schedule. Key advances have occurred recently on both fronts with the first X-ray crystal structure at 3.8 Å resolution of the water-oxidizing complex (WOC) and its associated photochemical reaction center (photosystem II) from a cyanobacterium^[7] and the first report of intramolecular O₂ photoproduction from the bridging oxygen atoms of a manganese–oxo cluster of cubane-type geometry, L₆Mn₄O₄ (**1**; L = diphenylphosphinate; Ph₂PO₂[−])



Scheme 1. Reactions of L₆Mn₄O₄ cubane complexes **1** and **1'**: a) UV photochemical reaction in the gas phase and b) reductive dehydration reaction in solution. RXH = organoamine, phenol, etc. Bridging phosphinates omitted for clarity.

[*] Prof. Dr. G. C. Dismukes, Prof. Dr. M. Yagi, K. V. Wolf
Dr. P. J. Baesjou Prof. Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, NJ, 08544-1009, NJ (USA)
Fax: (+1) 609-258-1980
E-mail: dismukes@princeton.edu

[**] This research was supported by grants from the US National Institutes of Health (GM39932) and the National Science Foundation (CHE96-19190). Fellowship support was provided by the Japan Society for the Promotion of Science (M.Y.) and the Netherlands NOW Scholarship (P.B.).