


- [1] J. Simon, J. J. André, A. Skoulios, *New J. Chem.* **1986**, *10*, 295–311; J. Simon, F. Tournilhac, J. J. André, *New J. Chem.* **1987**, *11*, 383–399; J. Simon, P. Bassoul, S. Norvez, *New J. Chem.* **1989**, *13*, 13–31.
- [2] J.-M. Lehn, *Supramolecular Chemistry—Concepts and Perspectives*, VCH, Weinheim, **1995**; D. Philp, J. F. Stoddart, *Angew. Chem.* **1996**, *108*, 1242–1286; *Angew. Chem. Int. Ed.* **1996**, *35*, 1155–1196.
- [3] O. Kahn, C. J. Martínez, *Science* **1998**, *279*, 44.
- [4] P. L. Boudas, M. Gómez-Kaifer, L. Echegoyen, *Angew. Chem.* **1998**, *110*, 226–258; *Angew. Chem. Int. Ed.* **1998**, *37*, 216–247.
- [5] V. Balzani, F. Scandola, *Supramolecular Photochemistry*, Ellis Horwood, New York, **1991**.
- [6] See for example M. J. Zaworotko, *Chem. Commun.* **2001**, 1–9; R. Robson, *J. Chem. Soc. Dalton Trans.* **2000**, 3735–3744.
- [7] A. D. Burrows, C. W. Chan, M. M. Chowdhry, J. E. McGrady, D. M. P. Mingos, *Chem. Soc. Rev.* **1995**, *24*, 351–359; M. D. Ward, C. M. White, F. Barigelli, N. Armadori, G. Calogero, L. Flamigni, *Coord. Chem. Rev.* **1998**, *171*, 481–488.
- [8] L. J. Childs, N. W. Alcock, M. J. Hannon, *Angew. Chem.* **2001**, *40*, 1113–1115; *Angew. Chem. Int. Ed.* **2001**, *40*, 1079–1081; M. J. Hannon, C. L. Painting, W. Errington, *Chem. Commun.* **1997**, 1805–1806.
- [9] N. W. Alcock, P. R. Barker, J. M. Haider, M. J. Hannon, C. L. Painting, Z. Pikramenou, E. A. Plummer, K. Rissanen, P. Saarenketo, *J. Chem. Soc. Dalton Trans.* **2000**, 1447–1461.
- [10] J. D. Swalen, D. L. Allara, J. D. Andrade, E. A. Chandross, S. Garoff, J. Israelachvili, T. J. McCarthy, R. Murray, R. F. Pease, J. F. Rabolt, K. J. Wynne, H. Yu, *Langmuir* **1987**, *3*, 932–950.
- [11] For examples of thin films with metal ions, see T. E. Mallouk, J. A. Gavin, *Acc. Chem. Res.* **1998**, *31*, 209–217; M. Maskus, H. D. Abruna, *Acc. Chem. Res.* **1996**, *12*, 4455–4462; Y. Liang, R. H. Schmehl, *J. Chem. Soc. Chem. Commun.* **1995**, 1007–1008; D. G. Kurth, P. Lehmann, M. Schütte, *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 5704–5707; A. Hatzor, T. Moav, H. Cohen, S. Matlis, J. Libman, A. Vaskevich, A. Shanzer, I. Rubinstein, *J. Am. Chem. Soc.* **1998**, *120*, 13469–13477.
- [12] G. Decher, *Science* **1997**, *277*, 1232.
- [13] M. Lahav, R. Gabai, A. N. Shipway, I. Willner, *Chem. Commun.* **1999**, 1937–1938; M. Lahav, V. Heleg-Shabtai, J. Wasserman, E. Katz, I. Willner, H. Durr, Y.-Z. Hu, S. Bossmann, *J. Am. Chem. Soc.* **2000**, *122*, 11480–11487.
- [14] M. Schütte, D. G. Kurth, M. R. Linford, H. Cölfen, H. Möhwald, *Angew. Chem.* **1998**, *110*, 3058–3061; *Angew. Chem. Int. Ed.* **1998**, *37*, 2891–2894; D. G. Kurth, C. Schüler, F. Caruso, *Chem. Commun.* **1999**, 1579–1580.
- [15] T. Salditt, Q. An, A. Plech, J. Piesl, C. Eschbaumer, C. H. Weidl, U. S. Schubert, *Thin Solid Films* **1999**, *354*, 208–214; U. S. Schubert, C. Eschbaumer, Q. An, T. Salditt, *J. Inclusion Phenom. Macrocyclic Chem.* **1999**, *35*, 35–43.
- [16] A range of aryl tails, including the biphenyl tails previously described in ref. [9], facilitate incorporation in films; the discussion herein is restricted to the pyrenyl tails because of their useful spectroscopic tag.
- [17] A square-planar Pt^{II} complex of this Pyr-tpy ligand was recently reported: J. F. Michalec, S. A. Bejune, D. R. McMillin, *Inorg. Chem.* **2000**, *39*, 2708–2709.
- [18] Crystal data for C₆₂H₃₈N₆FeCl₂O₈·3(CH₃NO₂): *M_r* = 1304.87, triclinic, space group *P*1̄, *a* = 11.997(10), *b* = 12.800(10), *c* = 21.665(10) Å, *α* = 89.762(10)°, *β* = 75.457(10)°, *γ* = 65.724(10)°, *V* = 2916.79(4) Å³, *T* = 180(2) K, *λ* = 0.71073, *Z* = 2, *ρ*_{calcd} = 1.486 Mg m^{−3}, *F*(000) = 1344, *μ*(MoK_α) = 0.429 mm^{−1}; purple plate, crystal dimensions 0.3 × 0.2 × 0.19 mm; data were collected on a Siemens SMART three-circle system with CCD area detector. The crystal was held at 180(2) K with an Oxford Cryosystem Cryostream Cooler; *θ*_{max} = 29.16°. A total of 18515 reflections were measured, 13214 of which were unique (*R*_{int} = 0.0159). Absorption correction by Psi scans. Weighting scheme: *w* = 1/[*σ*²(*F*_o²) + (0.0383 *P*)²], where *P* = (*F*_o² + 2 *F*_c²)/3. GOF on *F*² was 1.507, *R*₁ = 0.0537 (for 10583 reflections with *I* > 2 *σ*(*I*)), *wR*₂ = 0.2091. Data/restraints/parameters: 13214/0/823. Largest difference Fourier peak and hole: 1.294 and −0.894 e Å^{−3}. Refinement used SHELXL97 (G. M. Sheldrick, Universität Göttingen, **1997**). Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-160346. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [19] *Spectroscopy and Relaxation of Molecular Liquids* (Eds.: D. Steele, J. Yarwood), Elsevier, Amsterdam, **1991**.
- [20] D. G. Kurth, D. Volkmer, M. Ruttorf, B. Richter, M. J. Koop, A. Müller, *Chem. Mater.* **2000**, *12*, 2829–2831.
- [21] The surface coverage *Γ* was calculated according to *Γ* = 3/4(*A_λ**N*)/*ε_λ*, where *ε_λ* is the molar extinction coefficient [M^{−1} cm^{−1}] corrected for in-plane anisotropy, *N* is Avogadro's constant, and *A_λ* is the absorbance. The calculation assumes that *ε* is approximately the same in the thin film as in solution.
- [22] D. G. Kurth, R. E. Osterhout, *Langmuir* **1999**, *15*, 4842–4846.
- [23] F. M. Winnik, *Chem. Rev.* **1993**, *93*, 587–614.
- [24] The π–π stacking in the free ligand was confirmed by crystallographic analysis, which will be reported in forthcoming publications.
- [25] While the multilayers are stable towards solvents (water, alcohol, benzene, etc.), exposure to ethylenediaminetetraacetic acid (EDTA) disassembles the films (pH 11, room temperature). Such externally induced disassembly offers novel opportunities to fabricate membranes for controlled-release applications (see for instance G. B. Sukhorukov, E. Donath, S. Moya, A. S. Susa, A. Voigt, J. Hartmann, H. Möhwald, *J. Microencapsulation* **2000**, *17*, 177–185).
- [26] The intensity is not in scale for the two spectra, which are not corrected for photomultiplier tube response.

Iodoaziridine Derivatives as Novel Azahomoallyl Radical Precursors for [3+2] Cycloaddition Reactions with Alkenes


Osamu Kitagawa, Yoichiro Yamada, Hiroki Fujiwara, and Takeo Taguchi*

The [3+2] cycloaddition of a homoallyl radical with an alkene is a powerful method for the one-step synthesis of a cyclopentanoid skeleton.^[1] If such a cycloaddition could be achieved with an azahomoallyl (allylaminy) radical, it would provide a new and useful methodology for the preparation of pyrrolidine derivatives from simple substrates. As far as we know, only one example of the generation of an azahomoallyl radical and its reaction with an alkene has been reported, by Newcomb et al. [Eq. (1)].^[2] This method using *N*-allyl *N*-hydroxypyridine-2-thione carbamates as an azahomoallyl



[*] Prof. Dr. T. Taguchi, Assis. Prof. Dr. O. Kitagawa, Y. Yamada, H. Fujiwara
Tokyo University of Pharmacy and Life Science
1432-1 Horinouchi, Hachioji, Tokyo 192-0392 (Japan)
Fax: (+81) 426-76-3257
E-mail: kitagawa@ps.toyaku.ac.jp
taguchi@ps.toyaku.ac.jp

radical precursor requires 100 equiv of alkene to give reasonable yields of products (52–59%). Moreover, a Brønsted acid for the formation of the reactive allylaminium radical^[3] and a thiol as a hydrogen donor for the resulting pyrrolidinylmethyl radical are required in these reactions. In addition, due to the limited number of examples, the scope and limitations of this reaction are unclear. Application to alkenes other than enol ethers and reactions with other azahomoallyl radicals such as cycloalkenyl- and prenylaminy radicals were not investigated. In the addition of a nitrogen-centered radical to an alkene, intramolecular cyclization with a 4-pentenylaminyl radicals has been extensively demonstrated,^[1f, 4] while the intermolecular version has so far been uncommon.^[5] This may be due to the lower reactivity of nitrogen-centered radicals relative to carbon-centered radicals^[1f, 4b, 6] and the lack of a suitable radical precursor. Here we report on radical, iodine-transfer [3+2] cycloaddition with various alkenes and *N*-tosyloidoaziridine derivatives **1** as novel azahomoallyl radical precursors [Eq. (2)].



We expected that: 1) In the presence of a suitable radical initiator, *N*-tosyloidoaziridine (**1a**) might efficiently produce *N*-tosylallylamidyl radical **1A** by regioselective cleavage of the C–N bond of an aziridinylmethyl radical;^[7] 2) the reaction of **1A** with an alkene would possibly give an iodoalkylated pyrrolidine derivative **2** by a radical iodine-transfer mechanism; 3) *N*-Tosylamidyl radical **1A** may be more reactive toward alkenes than an aminyl radical because of its more electronegative character;^[8] and 4) the mild reaction conditions for the generation of **1A** and its addition to alkenes would make a wide range of alkenes employable.^[3] Furthermore, **1a** and various other iodoaziridine derivatives **1b–d** can be easily prepared by iodoaziridination of *N*-allyltosylamide derivatives.^[9] Thus, the generation of various azahomoallyl radicals such as allyl-, prenyl-, and cycloalkenylamidyl radicals may be also possible.

The [3+2] cycloadditions of **1a–d** with alkenes (2 equiv) were investigated in the presence of several radical initiators. The reaction of **1** with electron-rich alkenes such as enol ethers proceeded smoothly (Table 1). For example, the reaction of iodomethylaziridine **1a** with butyl vinyl ether under conditions A (1 equiv Et₃B, O₂ in CH₂Cl₂) efficiently proceeded even with only 2 equiv of the alkene to give functionalized iodomethylpyrrolidine **2a** in good yield (64%; entry 1).^[10] The reaction under conditions B (0.1 equiv (*n*Bu₃Sn)₂, *hν*) gave a lower yield (48%; entry 2). Since the present reaction can be performed under nonaqueous and neutral conditions,^[3] application to acid-sensitive silyl enol ethers is also possible. Thus, hydroxypyrrolidine **2b** was obtained in good yield (conditions A, 66%; conditions B, 80%) after desilylation of the cycloaddition product with 2 %

Table 1. Radical [3+2] cycloaddition of various iodoaziridines **1** with enol ethers.

				Products 2 A: Et ₃ B (1 equiv), O ₂ / CH ₂ Cl ₂ B: (<i>n</i> Bu ₃ Sn) ₂ (0.3 equiv), <i>hν</i> / C ₆ H ₆		Yield [%] ^[b]		cis/trans	
Entry	Aziridine	R ^{2[a]}	Condi-tions	Products 2					
1	1a	<i>n</i> Bu	A		2a	64		1 ^[c]	
2	1a	<i>n</i> Bu	B		2a	48		1 ^[c]	
3	1a	TMS	A		2b	66		1.3 ^[d]	
4	1a	TMS	B		2b	80		1.5 ^[d]	
5 ^[e]	1b	TMS	A		2c	63 ^[f]		–	
6 ^[e]	1b	TMS	B		2c	37 ^[e]		–	
7	1c	TMS	A		2d	60		1/1.5 ^[d,h]	
8	1d	TMS	A		2e	62		1.4 ^[d,h]	


[a] TMS = trimethylsilyl. [b] Yield of isolated product. [c] The ratio was determined by ¹H NMR spectroscopy. [d] The ratio was determined on the basis of isolated products. [e] The product (tertiary iodide) was subsequently treated with DBU because of its instability. [f] A mixture of internal double bond product (internal-**2c**) and terminal double bond product (terminal-**2c**) was obtained in a ratio of 4:1. The diastereomer ratio of terminal-**2c** was 5:1, while the configurations were not determined. [g] Internal-**2c**/terminal-**2c** = 4.3. A diastereomer ratio of terminal-**2c** = 5:1 [h] The formation of two stereoisomers was observed. The ratio corresponds to α-OH/β-OH.

HCl (entries 3 and 4). In the reaction of **1a** with silyl enol ethers, conditions B gave better results than conditions A.

The radical [3+2] cycloaddition reaction with 2 equiv of silyl enol ether can also be applied to the iodoalkylated aziridine derivatives **1b–1d**, which produce prenyl- and cycloalkenylamidyl radicals (entries 5–8). Under conditions A, reaction of **1b** with silyl enol ether also proceeded efficiently to give **2c** in good yield (63%) after treatment of the cycloadduct with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU; entry 5), while under conditions B, a lower yield was obtained (37%, entry 6). The reactions of bicyclic iodoaziridines **1c** and **1d** gave functionalized bicyclic nitrogen-containing compounds **2d** and **2e** in 60 and 62% yield, respectively (entries 7 and 8).

The present reaction using iodoaziridine **1a** can also be applied to alkyl-substituted alkenes (2 equiv) such as methylenecyclohexane and allylsilane because of the higher reactivity of the *N*-tosylamidyl radical compared to a simple aminyl radical. In these reactions, gradual generation of an *N*-

tosylallylamidyl radical by portionwise addition of Et₃B (3 × 0.5 equiv) increased the yield.^[11] Thus, the [3+2] cycloaddition products **2f** and **2g** were obtained in 56 and 51 % yield, respectively [Eq. (3)]. When 1 equiv of Et₃B was added in one



portion, considerably lower yields of **28** and **33** %, respectively, were obtained, and *N*-allyltosylamide was formed as a side product. The reaction with the simple 1-alkene 1-hexene gave **2h** in low yield (34 %), even with portionwise addition of Et₃B. These results may indicate that the reactivity of the *N*-tosylamidyl radical towards the alkene decreases with decreasing electron richness of the alkene.

In all reactions shown in Table 1 and Equation (3), formation of regioisomers was not observed. Thus, both cleavage of the C–N bond in the *N*-tosylaziridinylmethyl radical and the attack of the resulting *N*-tosylamidyl radical on the terminal carbon atom of alkene should proceed with complete regioselectivity, while the diastereoselectivities of the reactions are generally low (*cis/trans* = 1–2.6).

In conclusion, we have developed the radical iodine-transfer [3+2] cycloaddition of various alkenes with iodoaziridine derivatives as novel precursors of azahomoallyl radicals. The present reaction should provide a new and efficient methodology for the synthesis of functionalized pyrrolidine derivatives.

Experimental Section

cis/trans-2b: Et₃B (0.5 mL, 1 M in hexane) was added to a solution of iodoaziridine **1a** (169 mg, 0.5 mmol) and trimethylsilyl vinyl ether (0.15 mL, 1 mmol) in CH₂Cl₂ (4 mL) under Ar. Dry air (20 mL) was subsequently introduced with a syringe. After stirring for 10 h at room temperature, 5 % HCl (4 mL) and MeOH (15 mL) were added, and the mixture was then stirred for 2 h at room temperature. After removal of MeOH by evaporation, the residue was extracted with Et₂O. The Et₂O extracts were washed with brine, dried over MgSO₄, and evaporated to dryness. Purification of the residue by column chromatography (hexane/AcOEt 1/1) gave a mixture of *cis-2b* and *trans-2b*. Further purification of the mixture by MPLC (hexane/AcOEt 2/1) gave *cis-2b* (71 mg, 37 %, less polar) and *trans-2b* (56 mg, 29 %, more polar). The configurations of *cis-2b* and *trans-2b* were determined by NOE experiments.

cis-2b: white solid, m.p. 130–131 °C; IR (KBr): $\tilde{\nu}$ = 3492, 1322, 1152 cm^{−1}; ¹H NMR (300 MHz, CDCl₃, 25 °C, TMS) δ = 7.72 (d, *J* = 8.0 Hz, 2H), 7.34 (d, *J* = 8.0 Hz, 2H), 4.31 (q, *J* = 3.8 Hz, 1H), 3.60 (dd, *J* = 7.9, 9.6 Hz, 1H), 3.51 (dd, *J* = 3.8, 11.6 Hz, 1H), 3.43 (dd, *J* = 1.0, 11.6 Hz, 1H), 3.17 (dd, *J* = 9.0, 9.8 Hz, 1H), 3.10 (dd, *J* = 6.8, 9.8 Hz, 1H), 3.01 (dd, *J* = 9.6, 10.6 Hz, 1H), 2.44 (s, 3H), 2.42 (m, 1H), 1.59 (d, *J* = 4.9 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃, 25 °C, TMS) δ = 143.8, 133.6, 129.8, 127.4, 71.2, 56.4, 50.9, 47.3, 21.5, 0.4; MS: *m/z*: 381 [*M*⁺]; HRMS calcd for C₁₂H₁₆INO₃S [*M*⁺]: 380.9896; found 380.9869.

trans-2b: white solid, m.p. 136–137 °C; IR (KBr): $\tilde{\nu}$ = 3501, 1338, 1160 cm^{−1}; ¹H NMR (300 MHz, CDCl₃, 25 °C, TMS): δ = 7.71 (d, *J* = 8.3 Hz, 2H), 7.34 (d, *J* = 8.3 Hz, 2H), 4.05 (m, 1H), 3.56 (dd, *J* = 5.9, 10.5 Hz, 1H), 3.51 (dd, *J* = 7.5, 10.5 Hz, 1H), 3.05–3.15 (m, 3H), 2.97 (dd,

J = 7.8, 10.5 Hz, 1H), 2.44 (s, 3H), 2.28 (m, 1H), 2.17 (brs, 1H); ¹³C NMR (75 MHz, CDCl₃, 25 °C, TMS): δ = 144.0, 132.8, 129.9, 127.5, 74.6, 54.3, 52.2, 48.7, 21.6, 4.9; MS: *m/z*: 381 [*M*⁺]; HRMS calcd for C₁₂H₁₆INO₃S [*M*⁺]: 380.9896; found: 380.9877.

Received: April 30, 2001 [Z17013]

- [1] Reactions with electron-deficient alkenes: a) D. L. J. Clive, A. G. Angoh, *J. Chem. Soc. Chem. Commun.* **1985**, 980–982; b) Z. Cekovik, R. Saicic, *Tetrahedron Lett.* **1986**, 27, 5893–5896; c) D. P. Curran, M. Chen, *J. Am. Chem. Soc.* **1987**, 109, 6558–6560; d) D. H. R. Barton, S. Z. Zard, E. daSilva, *J. Chem. Soc. Chem. Commun.* **1988**, 285–287; e) K. S. Feldman, A. L. Romanelli, R. E. Ruckle, Jr., R. F. Miller, *J. Am. Chem. Soc.* **1988**, 110, 3300–3302; f) D. P. Curran in *Comprehensive Organic Synthesis*, Vol. 4 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, **1991**, pp. 779–831, and references therein. Reactions with simple alkenes: g) K. Miura, K. Fugami, K. Ohshima, K. Utimoto, *Tetrahedron Lett.* **1988**, 29, 5135–5138; h) D. P. Curran, M. Chen, E. Spletzer, C. M. Seong, C. Chang, *J. Am. Chem. Soc.* **1989**, 111, 8872–8878; i) D. P. Curran, C. M. Seong, *Tetrahedron* **1992**, 48, 2157–2174; j) D. P. Curran, C. M. Seong, *Tetrahedron* **1992**, 48, 2175–2190; k) O. Kitagawa, H. Fujiwara, T. Taguchi, *Tetrahedron Lett.* **2001**, 42, 2165–2167.
- [2] M. Newcomb, M. U. Kumar, *Tetrahedron Lett.* **1990**, 31, 1675–1678.
- [3] It is well known that electrophilic aminium radical cations and metal-complexed aminyl radicals formed by the addition of Brønsted and Lewis acids are more reactive towards alkenes than simple aminyl radicals. a) R. S. Neals, N. L. Marcus, *J. Org. Chem.* **1967**, 32, 3273–3284; b) J. M. Surzur, L. Stella, P. Tordo, *Bull. Soc. Chim. Fr.* **1970**, 115–120; c) R. S. Neale, *Synthesis* **1971**, 1–15; d) J. M. Dickinson, J. A. Murphy, *J. Chem. Soc. Chem. Commun.* **1990**, 434–436; e) C. Ha, O. M. Musa, F. N. Martinez, M. Newcomb, *J. Org. Chem.* **1997**, 62, 2704–2710; f) M. Hemmerling, A. Sjöholm, P. Somfai, *Tetrahedron: Asymmetry* **1999**, 10, 4091–4094, and references therein.
- [4] Reviews: a) L. Stella, *Angew. Chem.* **1983**, 95, 368–381; *Angew. Chem. Int. Ed. Engl.* **1983**, 22, 337–350; b) J. L. Esker, M. Newcomb in *Advances in Heterocyclic Chemistry*, Vol. 58 (Ed.: A. R. Katritzky), Academic Press, San Diego, **1993**, pp. 1–45; c) A. G. Fallis, I. M. Brinza, *Tetrahedron* **1997**, 53, 17543–17593.
- [5] For a review: P. Mackiewicz, R. Furstoss, *Tetrahedron* **1978**, 34, 3241–3260.
- [6] a) M. Newcomb, M. T. Burchill, T. M. Deeb, *J. Am. Chem. Soc.* **1988**, 110, 6528–6535; b) O. M. Musa, J. H. Horner, H. Shahin, M. Newcomb, *J. Am. Chem. Soc.* **1996**, 118, 3862–3868.
- [7] Regioselective cleavage of aziridinylcarbinyl radicals and application to intramolecular radical cyclization: a) J. M. Dickinson, J. A. Murphy, *Tetrahedron* **1992**, 48, 1317–1326; b) N. De Kimpe, R. Jolie, D. De Smaele, *J. Chem. Soc. Chem. Commun.* **1994**, 1221–1222; c) D. De Smaele, P. Bogaert, N. De Kimpe, *Tetrahedron Lett.* **1998**, 39, 9797–9800; d) review: J. J. Li, *Tetrahedron* **2001**, 57, 1–24.
- [8] J. H. Horner, O. M. Musa, A. Bouvier, M. Newcomb, *J. Am. Chem. Soc.* **1998**, 120, 7738–7748.
- [9] O. Kitagawa, T. Suzuki, T. Taguchi, *J. Org. Chem.* **1998**, 63, 4842–4845.
- [10] This indicates that together with the addition of *N*-tosylamidyl radical to an enol ether, iodine transfer to the resulting pyrrolidinylmethyl radical from iodoaziridine **1a** should also proceed efficiently, that is, the formation of aziridinylmethyl radical may be the thermodynamically favorable process.
- [11] **Caution:** When Et₃B was added to the reaction mixture at a high concentration of O₂, an explosion occurred once. Hence, further addition of Et₃B must be performed under Ar or N₂ atmosphere.