

approximately the same level of bond fixation (23 %, 53 %, 48 %, 48 %, 49 %, shortest bond 1.358 Å, longest bond 1.456 Å). The interplanar angles between consecutive six- and four-membered rings starting from the terminus are 1.31, 2.98, 3.61, 4.67, 3.84, 5.47, 4.50, and 4.99°, a smoother progression than that of **3a**, likely due to the absence of crystal lattice forces.^[14] The closest contact is between H1 and C22f (2.99 Å), the terminal ring centroids are separated by 7.48 Å, with an interplanar angle of 33.2°. Taking the central six-membered ring as a reference, the inner helix climbs 5.07 Å and turns in plane by 415.7°.

Are heliphenes **3** and **4** sufficiently extended to render their enantiomers amenable to resolution? The answer is provided by variable-temperature NMR experiments carried out on derivatives **3b** and **4b**. Remarkably, decoalescence of the potentially diastereotopic methylene hydrogens of the substituent in **3b** occurred only on cooling, providing $\Delta G_{-45^\circ\text{C}}^\ddagger = 13.4 \pm 0.4 \text{ kcal mol}^{-1}$ (400 MHz, $\Delta\nu_{\text{max}} = 12.6 \text{ Hz}$, $[\text{D}_8]\text{toluene}$). This value is less than 1 kcal mol⁻¹ higher than that of the corresponding [7]heliphenes **2**.^[1] To our further surprise, [9]heliphenes **4b** did not show any change in the shape of the corresponding methylene singlet signal down to the (solubility) limiting temperature of -45°C (400 MHz, $[\text{D}_8]\text{toluene}$), suggesting a lower barrier to enantiomerization, less than around 12 kcal mol⁻¹. As proposed previously,^[1,15] it appears that steric activation of the helical ground state toward unwinding the helix is responsible for the observed dynamic behavior. Structural elements, such as additional stereocenters biasing the helix or steric blockers locking it, will have to be introduced to render the heliphenes frame configurationally robust. Finally, it is evident that extensions of the “stitching” procedures outlined in Schemes 1–3 should provide access to even higher members of the heliphenes series. Such is the focus of future research.

Received: April 24, 2002 [Z19166]

- [1] S. Han, A. D. Bond, R. L. Disch, D. Holmes, J. M. Schulman, S. J. Teat, K. P. C. Vollhardt, G. D. Whitener, *Angew. Chem.* **2002**, *114*, 3357–3361; *Angew. Chem. Int. Ed.* **2002**, *41*, 3223–3227
- [2] R. Boese, A. J. Matzger, D. L. Mohler, K. P. C. Vollhardt, *Angew. Chem.* **1995**, *107*, 1630; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1478.
- [3] Attempted direct Pd-catalyzed installation of the C₂ bridge with ethyne failed, although it is successful in related couplings, see: a) R. Boese, A. J. Matzger, K. P. C. Vollhardt, *J. Am. Chem. Soc.* **1997**, *119*, 2052; b) M. Iyoda, A. Vorasingha, Y. Kuwatani, M. Yoshida, *Tetrahedron Lett.* **1998**, *39*, 4701; c) K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* **1975**, 4467.
- [4] H.-D. Beckhaus, R. Faust, A. J. Matzger, D. L. Mohler, D. W. Rogers, C. Rüchardt, A. K. Sawhney, S. P. Verevkin, K. P. C. Vollhardt, S. Wolff, *J. Am. Chem. Soc.* **2000**, *122*, 7819.
- [5] C. Eickmeier, H. Junga, A. J. Matzger, F. Scherhag, M. Shim, K. P. C. Vollhardt, *Angew. Chem.* **1997**, *109*, 2194; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2103.
- [6] R. H. Schmidt-Radde, K. P. C. Vollhardt, *J. Am. Chem. Soc.* **1992**, *114*, 9713.
- [7] All new isolated compounds gave satisfactory analytical and/or spectral data (see Table 1).
- [8] R. A. Hites in *Chemical Analysis of Polycyclic Aromatic Compounds* (Ed.: T. Vo-Dinh), Wiley, New York, **1989**, p. 219.
- [9] D. L. Mohler, K. P. C. Vollhardt in *Advances in Strain in Organic Chemistry*, Vol. 5 (Ed.: B. Halton), JAI, London, **1996**, p. 121.
- [10] In analogy to: a) G. Wenz, M. A. Müller, M. Schmidt, G. Wegner, *Macromolecules* **1984**, *17*, 837; see also: b) R. E. Martin, F. Diederich,

Angew. Chem. **1999**, *111*, 1441; *Angew. Chem. Int. Ed.* **1999**, *38*, 1350; c) Y. Geerts, G. Klärner, K. Müllen in *Electronic Materials: The Oligomer Approach* (Eds.: K. Müllen, G. Wegner), Wiley-VCH, Weinheim, **1998**, p. 1; for a caveat in applying this simple correlation, see d) H. Meier, U. Stalmach, H. Kolshorn, *Acta Polymer.* **1997**, *48*, 379.

- [11] a) J. M. Schulman, R. L. Disch, *J. Phys. Chem. A* **1997**, *101*, 5596; b) J. M. Schulman, R. L. Disch, H. Jiao, P. von R. Schleyer, *J. Phys. Chem. A* **1998**, *102*, 8051.
- [12] At the B3LYP/6-31G* level; the maximum and (in parentheses) average deviations of the calculated from the experimental bond lengths are 0.023 (0.0037) Å.
- [13] X-ray structure analysis: **3a**·[CH₂Cl₂]_{0.5} (synchrotron radiation, Station 9.8, SRS Daresbury): crystal size 0.08 × 0.06 × 0.01 mm³, space group *P2₁/c* (no. 14), scan range 4.80 < 2θ < 47.7°, *a* = 8.2710(6), *b* = 23.1799(18), *c* = 16.2279(13) Å, *V* = 3090.5(4) Å³, *Z* = 4, $\rho_{\text{calcd}} = 1.374 \text{ g cm}^{-3}$, $\mu(\lambda = 0.6880 \text{ Å}) = 0.78 \text{ cm}^{-1}$, 15 619 total reflections, 5266 unique reflections (*R*_{int} = 0.0411) at -53°C, of which 4062 were taken as observed [*F*² > 2.00σ(*F*²)], conventional *R*₁ = 0.0878, *R*_w = 0.2369 (on all 5266 *F*² data). CCDC-183203 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44)1223-336-033; or deposit @ccdc.cam.ac.uk).
- [14] D. Holmes, S. Kumaraswamy, A. J. Matzger, K. P. C. Vollhardt, *Chem. Eur. J.* **1999**, *5*, 3399.
- [15] While we cannot rule out accidental isochronism of the methylene hydrogens, its selective occurrence in **4b**, but not **3b** or the corresponding derivative of **2** in ref. [1], would seem unlikely.

A New Pd⁰–Cu^I Bimetallic Catalyst for the Synthesis of Indoles from Isocyanates and Allyl Carbonates**

Shin Kamijo and Yoshinori Yamamoto*

In the research field of transition-metal-catalyzed organic synthesis, one of the modern trends is to use a bimetallic catalyst system to explore not only more efficient but also entirely new transformations. Representative examples of such bimetallic catalysis in palladium-catalyzed reactions^[1] are the following: 1) cat.Pd⁰/Cu^I in the Sonogashira coupling reaction, in which a catalytic amount of Cu salts activate terminal acetylenes to form copper–acetylide species;^[2] 2) cat Pd⁰/Cu^I in the Stille coupling reaction, in which organostannanes are transmetalated to give organocoppers which

[*] Prof. Dr. Y. Yamamoto, Dr. S. Kamijo
Department of Chemistry
Graduate School of Science, Tohoku University
Sendai 980-8578 (Japan)
Fax: (+81)22-217-6784
E-mail: yoshi@yamamoto1.chem.tohoku.ac.jp


[**] We thank faculties in the Instrumental Analysis Center, Department of Chemistry at Tohoku University, for measurement of mass spectra and elemental analysis. S.K. is also thankful for fellowship support from the Japan Society of Promotion of Science for Young Scientists.

 Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


accelerate the coupling;^[3] 3) cat.Pd⁰/excess Ag^I in the Heck reaction, in which Ag salts abstract the halide ion to produce a reactive cationic Pd intermediate;^[4] 4) cat.Pd⁰/Rh^I in the asymmetric Tsuji–Trost reaction, in which Rh activates pronucleophiles by coordinating to a cyano group.^[5] Accordingly, the above examples 1) and 2) are categorized into the Pd⁰–M group, while examples 3) and 4) belong to the Pd⁰–LA

Pd⁰–M: A starting substrate is converted into a more reactive organometallic reagent

1) Sonogashira coupling reaction




2) Stille coupling reaction




Pd⁰–LA: Lewis acid activates a starting substrate or intermediate

3) Heck reaction




4) Asymmetric Tsuji–Trost reaction



Scheme 1. Bimetallic catalyst systems in Pd-Catalyzed Reactions. Pd⁰–M: A starting material is converted into a more reactive organometallic complex. 1) Sonogashira coupling reaction; 2) Stille coupling reaction. Pd⁰–LA: LA acts as a Lewis acid to activate a starting substrate or an intermediate. 3) Heck reaction; 4) Asymmetric Tsuji–Trost reaction.

family (LA = Lewis acid, Scheme 1). We now report that the coupling of the (2-alkynyl)phenylisocyanate **1** and the allyl carbonate **2** proceeds using a palladium–copper bimetallic catalyst system to afford indoles **3**^[6–9] in high yields [Eq. (1)],



in which Cu^I ions most probably act as π -coordinative Lewis acids.^[10]

We first investigated several catalytic systems for the reaction of **1** with **2**. The results are summarized in Table 1. The reaction of 2-(1-pentynyl)phenyl-1-isocyanate (**1a**) and allyl methyl carbonate (**2a**) in the presence of a catalytic amount of [Pd(PPh₃)₄] gave the *N*-allylaniline derivative **4aa** in 83 % yield (entry 1). After intensive investigations, we

Table 1. Effect of the additive.^[a]

Entry	Additive	NMR yield (isolated) [%] ^[b]	
		3aa	4aa
1	none	0	89 (83)
2	CuCl	78 (73)	0
3	CuBr	70	trace
4	CuI	27	20
5	[(CuOTf) ₂ ·C ₆ H ₆] ^[c]	37	0
6	CuOAc	26	37
7	Cu ₂ O	0	69
8 ^[d]	CuCl ₂	0	0
9	ZnCl ₂	62	0
10	AgCl	0	70
11	PdCl ₂	0	74
12 ^[e]	LiCl	0	0
13	<i>n</i> Bu ₄ NCl	0	80

[a] **2a** (2 equiv) was added to a solution of **1a**, [Pd(PPh₃)₄] (5 mol %), and additive (20 mol %) in THF, and the mixture was heated at 100 °C for 1 h. [b] Yields were determined by ¹H NMR spectroscopy with *p*-xylene as an internal standard (yield is shown in parentheses). [c] 10 mol % additive was used. [d] The starting material **1a** was recovered in 29 % yield, as suggested by NMR spectroscopy. [e] The starting material **1a** was recovered in 69 % yield, as suggested by NMR studies.

found that Cu^I species effectively catalyze the cyclization to form the indole **3aa**. Of the additives we tested, CuCl showed the best catalytic activities and gave the desired indole **3aa** in 73 % yield (entry 2), while CuBr showed a comparable activity to CuCl (entry 3). Other Cu^I catalysts such as CuI, [(CuOTf)₂·C₆H₆], and CuOAc gave lower yields of **3aa** (entries 4–6) and Cu₂O displayed no catalytic activity for the cyclization (entry 7). However, Cu^{II} catalysts inhibited the formation of both **3aa** and **4aa** (entry 8). Among the other metal-based catalysts, ZnCl₂ was found to catalyze the cyclization to form **3aa** in 62 % yield and did not give **4aa** at all (entry 9). AgCl and PdCl₂ did not catalyze the cyclization and gave only **4aa** (entries 10 and 11). LiCl inhibited the formation of both **3aa** and **4aa** and resulted in the recovery of the starting material **1a** in 69 % yield (entry 12), while tetrabutylammonium chloride did not yield **3aa** (entry 13). We found after further optimization of the reaction conditions that the reaction proceeds efficiently in THF by using 1 mol % of [Pd(PPh₃)₄] and 4 mol % of CuCl as cocatalysts [Eq. (1)].

We investigated the scope and limitations of the indole synthesis based on the bimetallic catalyst system. We first treated the (2-alkynyl)phenylisocyanate **1a** [R¹ = Pr in Eq. (1)] with various allyl carbonates **2** (Table 2). The reaction of **1a** and allyl methyl carbonate (**2a**) was conducted in the presence of catalytic amounts of [Pd(PPh₃)₄] (1 mol %) and CuCl (4 mol %) in THF (1M) at 100 °C for 1 h. The desired 3-allyl-*N*-(methoxycarbonyl)-2-propylindole (**3aa**) was obtained in 81 % yield (entry 1). The reaction with bulky allyl carbonates **2b** and **2c** also proceeded smoothly to give the corresponding indoles **3ab** and **3ac** in 69 and 72 % yields, respectively (entries 2 and 3). Even allyl phenyl carbonate

Table 2. Indole synthesis from **1a** and various allyl carbonates.^[a]

Entry	2	R ²	3	Yield [%]
1	2a	Me	3aa	81
2	2b	<i>i</i> Pr	3ab	69
3	2c	<i>t</i> Bu	3ac	72
4	2d	Ph	3ad	86
5	2e	Bn	3ae	83

[a] **2** (1.2 equiv) was added to a solution of **1a**, [Pd(PPh₃)₄] (1 mol %), and CuCl (4 mol %) in THF, and the mixture was stirred at 100 °C for 1 h.

(**2d**) and allyl benzyl carbonate (**2e**) gave the corresponding indoles **3ad** and **3ae** in good yields (entries 4 and 5).

We next treated allyl methyl carbonate (**2a**) [R² = Me in Eq. (1)] with isocyanate **1** having various acetylenic terminal groups (Table 3). The isocyanates with propyl (**1a**) and cyclopentyl (**1b**) substituents afforded the corresponding


Table 3. Indole synthesis from **2a** and various isocyanates.^[a]

Entry	1	R ¹	Reaction time [h]	3	Yield [%]
1	1a	Pr	1	3aa	81
2 ^[b]	1b	cyclopentyl	3	3ba	71
3	1c	<i>t</i> Bu	5	3ca	0 ^[c]
4 ^[b]	1d	Ph	2	3da	62
5 ^[d]	1e	<i>p</i> -MeOC ₆ H ₄	6	3ea	62
6 ^[d]	1f	<i>p</i> -CF ₃ C ₆ H ₄	7	3fa	65

[a] The reaction was conducted with **1** and **2a** (1.2 equiv) in the presence of [Pd(PPh₃)₄] (1 mol %) and CuCl (4 mol %) in THF (1 M) at 100 °C for the time displayed in Table 3. [b] The reaction was conducted in 0.5 M THF solution. [c] **4ca** was obtained in 42 % yield. [d] Two equivalents of **2a** were used.

indoles **3aa** and **3ba** in 81 and 74 %, respectively (entries 1 and 2). However, the isocyanate with a terminal *tert*-butyl group (**1c**) afforded the corresponding *N*-allylaniline derivative **4ca** in 42 % yield and did not give the corresponding indole **3ca** at all (entry 3). The phenyl-substituted isocyanates **1d–1f** gave the corresponding indoles **3da–3fa** in good yields (entries 4–6). The above results indicate that the reaction proceeds irrespective of the electronic nature of the aromatic ring.

A proposed mechanism is shown in Scheme 2. Initially, the Pd⁰ center reacts with allyl carbonates **2** to give the π -allylpalladium alkoxide complex **A** with extrusion of CO₂. Then, the insertion of isocyanates **1** into complex **A** would form the π -allylpalladium intermediate **B**. This intermediate, with Pd–N bonding, could be in equilibrium with the Pd–O-bonded intermediate **C**, which should more probably be represented as the bis- π -allylpalladium analogue **D**.^[11] Insertion of the alkyne then occurs to form indole **3** and the Pd⁰ species is regenerated. The key step in this proposed mechanism is the insertion of the alkyne into the π -allylpalladium intermediates **B–D**, which occurs to give the *trans* indoles **3**. The role of CuCl is not understood completely at present, but we believe that the Cu^I center behaves as a Lewis acid for the C≡C bond to activate the alkyne through π coordination and catalyze the insertion of the alkyne into intermediates **B–D**, that is, it acts as a catalyst for the carboamination of the alkyne.^[12] An alternative mechanism through the intermediate **E** can be proposed, in which copper



Scheme 2. Proposed mechanism for the formation of indoles.

activates the nitrogen nucleophile and the alkyne bond coordinates to a π -allylpalladium cationic species.^[17] At the present time, it is impossible to distinguish between the two mechanisms; the alkyne bond is activated by copper or by palladium species. It should be emphasized that no carboamination takes place at all in the absence of CuCl.

In conclusion, we have developed a new [Pd(PPh₃)₄]/CuCl bimetallic catalyst system for the synthesis of indoles **3** from isocyanates **1** and allyl carbonates **2**. This is the first example of a palladium-catalyzed indole syntheses from isocyanates. Various kinds of indoles are now available, and deprotection of the alkoxycarbonyl group is relatively easy.^[18] Further studies on synthetic applications and a mechanistic investigation are now in progress.

Experimental Section

Allyl methyl carbonate (**2a**, 69 μ L, 0.6 mmol) was added to a solution of 2-(1-pentynyl)phenyl-1-isocyanate (**1a**, 92.7 mg, 0.5 mmol), [Pd(PPh₃)₄] (5.8 mg, 0.005 mmol), and CuCl (2.0 mg, 0.02 mmol) in THF (0.5 mL) under an argon atmosphere, and the mixture was stirred at 100 °C for 1 h. The reaction mixture was cooled to room temperature, filtered through a short Florisil pad, and concentrated. The residue was purified by column chromatography (silica gel, hexane/ethyl acetate 100/1 to 20/1) to afford 3-allyl-*N*-(methoxycarbonyl)-2-propylindole (**3aa**) in 81 % yield (104.2 mg).

Received: March 11, 2002 [Z18865]

[1] a) *Metal-catalyzed Cross-coupling Reactions*, (Eds.: F. Diederich, P. J. Stang), Wiley-VCH, Weinheim, 1998; b) *Transition Metal Reagents and Catalysts*, (Ed.: J. Tsuji), Wiley, New York, 2000.

[2] a) K. Sonogashira, Y. Tohda, N. Hagihara, *Tetrahedron Lett.* **1975**, 4467–4470; b) L. Anastasia, E.-i. Negishi, *Org. Lett.* **2001**, 3, 3111–

- 3113; c) A. Mori, J. Kawashima, T. Shimada, M. Suguro, K. Hirabayashi, Y. Nishimura, *Org. Lett.* **2000**, 2, 2935–2937.
- [3] a) M. Shibasaki, C. D. J. Boden, A. Kojima, *Tetrahedron*, **1997**, 53, 7371–7395, and references therein; b) M. M. Abelman, T. Oh, L. E. Overman, *J. Org. Chem.* **1987**, 52, 4133–4135; c) A. Ashimori, B. Bachand, L. E. Overman, D. J. Poon, *J. Am. Chem. Soc.* **1998**, 120, 6477–6487.
- [4] V. Farina, S. Kapadia, B. Krishnan, C. Wang, L. S. Liebeskind, *J. Org. Chem.* **1994**, 59, 5905–5911.
- [5] M. Sawamura, M. Sudoh, Y. Ito, *J. Am. Chem. Soc.* **1996**, 118, 3309–3310.
- [6] Reviews for the indole syntheses, see: a) *Indoles* (Ed.: R. J. Sundberg), Academic, London, **1996**; b) *Palladium in Heterocyclic Chemistry* (Eds.: J. J. Li, W. Gribble), Pergamon, Oxford, **2000**, chap. 3; c) J. A. Joule in *Science of Synthesis (Houben-Weyl Methods of Molecular Transformations)*, Vol. 10 (Ed.: E. J. Thomas), Thieme, Stuttgart, **2000**, pp. 361–652; d) G. W. Gribble, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1045–1075.
- [7] For a recent report for the palladium-catalyzed indole synthesis, see: A. Takeda, S. Kamijo, Y. Yamamoto, *J. Am. Chem. Soc.* **2000**, 122, 5662–5663, and references therein.
- [8] An indole synthesis from 2-alkynyl-*N*-(CO₂R)aniline derivatives has been reported but a cyclization with complete retention of the CO₂R group is rare, see: a) K. Iritani, S. Matsubara, K. Utimoto, *Tetrahedron Lett.* **1988**, 29, 1799–1802; b) K. Hiroya, S. Itoh, M. Ozawa, Y. Kanamori, T. Sakamoto, *Tetrahedron Lett.* **2002**, 43, 1277–1280; c) T. Sakamoto, Y. Kondo, S. Iwashita, T. Nagano, H. Yamanaka, *Chem. Pharm. Bull.* **1988**, 36, 1305–1308; d) A. Yasuhara, Y. Kanamori, M. Kaneko, A. Numata, Y. Kondo, T. Sakamoto, *J. Chem. Soc. Perkin Trans. 1* **1999**, 529–534.
- [9] For the formation of 3-allylindoles, see: a) S. Cacchi, G. Fabrizi, P. Pace, *J. Org. Chem.* **1998**, 63, 1001–1011; b) ref. [8a].
- [10] For Lewis acid catalyzed reactions of alkynes, see: N. Asao, Y. Yamamoto, *Bull. Chem. Soc. Jpn.* **2000**, 73, 1071–1087, and references therein.
- [11] For the chemistry on bis- π -allylpalladium, see: a) H. Nakamura, H. Iwama, Y. Yamamoto, *J. Am. Chem. Soc.* **1996**, 118, 6641–6647; b) H. Nakamura, J.-G. Shim, Y. Yamamoto, *J. Am. Chem. Soc.* **1997**, 119, 8113–8114; c) E. Yoshikawa, K. V. Radhakrishnan, Y. Yamamoto, *Tetrahedron Lett.* **2000**, 41, 729–731; d) S. Kamijo, T. Jin, Y. Yamamoto, *J. Am. Chem. Soc.* **2001**, 123, 9453–9454.
- [12] The complexation between an alkyne and CuCl was reported nearly two decades ago by D. W. Macomber and M. D. Rausch,^[13] and more recently such complexation has been applied to the synthesis of alkynyl–copper complexes.^[14,15] In addition, ZnX₂–alkyne complexes have also been isolated.^[16]
- [13] D. W. Macomber, M. D. Rausch, *J. Am. Chem. Soc.* **1983**, 105, 5325–5329.
- [14] H. Ito, K. Arimoto, H.-o. Sensui, A. Hosomi, *Tetrahedron Lett.* **1997**, 38, 3977–3980.
- [15] a) K. Ikegashita, Y. Nishihara, K. Hirabayashi, A. Mori, T. Hiyama, *Chem. Commun.* **1997**, 1039–1040; b) Y. Nishihara, K. Ikegashita, A. Mori, T. Hiyama, *Chem. Lett.* **1997**, 1233–1234; c) Y. Nishihara, K. Ikegashita, A. Mori, T. Hiyama, *Tetrahedron Lett.* **1998**, 39, 4075–4078; d) Y. Nishihara, K. Ikegashita, K. Hirabayashi, J.-i. Ando, A. Mori, T. Hiyama, *J. Org. Chem.* **2000**, 65, 1780–1787; e) Y. Nishihara, M. Takemura, A. Mori, K. Osakada, *J. Organomet. Chem.* **2001**, 620, 282–286.
- [16] H. Lang, N. Mansilla, G. Rheinwald, *Organometallics* **2001**, 20, 1592–1596.
- [17] A referee suggested this alternative mechanism. We appreciate this possibility being pointed out.
- [18] T. W. Green, P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Wiley, New York, **1999**, pp. 503–550.

Fluorescence Enhancement through Enzymatic Cleavage of Internally Quenched Dendritic Peptides: A Sensitive Assay for the AspN Endoproteinase**

John M. Ellard, Thomas Zollitsch, W. Jon Cummins, Alan L. Hamilton, and Mark Bradley*

Fluorescence resonance energy transfer (FRET) systems in which a peptide sequence contains both a fluorophore and an internal quencher are amongst the best methods available for protease analysis and characterization. Numerous proteases have been studied using this method, including trypsin which cleaves the FRET-based system Dabcyl-Gly-Pro-Ala-Lys-Leu-Ala-Ile-Gly-Edans,^[1] cathepsin B which cleaves Abz-Lys-Leu-X-Phe-Ser-Lys-Gln-EDDnp (where X is Cys(SBz), Thr(OBz), or Ser(OBz) and EDDnp is 2,4-dinitrophenylthylenediamine),^[2] leukotriene D₄ hydrolase which cleaves ϵ -DNP-L-Lys-D-Amp^[3] (where Amp is 2-amino-3-(7-methoxy-4-coumaryl)propanoic acid) and caspase 1 and 3 which cleave the FRET systems BFP-Tyr-Val-Ala-Asp-GFP and BFP-Asp-Glu-Val-Asp-GFP, respectively (where BFP is blue fluorescent protein and GFP is green fluorescent protein)^[4] to name but a few. A host of methods have been developed using this methodology to determine the most appropriate substrate for a particular protease. For example, Meldal et al. have reported a powerful general method for the use of a split-and-mix approach to libraries of FRET-based peptides for on-bead screening to determine optimal substrates for a range of proteases including subtilisin carlsberg,^[5] FRET-based hexapeptide libraries have been used to map the S'-subsite specificity of serine proteases using solution-based assays,^[6] and a positional scanning substrate library has been used to determine the specificity of ICE (= interleukin-converting enzyme).^[7]

In this paper a new method for the detection of proteolytic activity is presented using internal fluorescence quenching between fluorophores of the same type. These are attached to a dendrimer-type structure, thereby giving a high local concentration of fluorophore needed for quenching yet eliminating the need for a separate quenching moiety. This method simplifies the synthesis of substrates for assay compared to traditional FRET, while being very sensitive due to the amplification nature of the assay as multiple

[*] Prof. M. Bradley, Dr. J. M. Ellard
Department of Chemistry
University of Southampton
Highfield, Southampton SO17 1BJ (UK)
Fax: (+44)23-80596766
E-mail: mb14@soton.ac.uk

Dr. W. J. Cummins, Dr. A. L. Hamilton
Amersham Biosciences
Amersham Place, Little Chalfont, Buckinghamshire HP79NA (UK)
T. Zollitsch
Current address: Infineon Technologies AG
Wernerwerkstrasse 2, 39049 Regensburg (Germany)

[**] The authors gratefully acknowledge Mark S. J. Briggs for helpful comments and suggestions.

 Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.