


A Colorimetric and Ratiometric Fluorescent Chemosensor with Three Emission Changes: Fluoride Ion Sensing by a Triarylborane–Porphyrin Conjugate**

Yohei Kubo, Masashi Yamamoto, Masato Ikeda,
Masayuki Takeuchi,* Seiji Shinkai,*
Shigehiro Yamaguchi, and Kohei Tamao

The design of chemosensors for a specific analyte is of great importance for the facile monitoring of analytes.^[1,2] Most chemosensors developed so far are colorimetric and/or fluorescent sensors, which efficiently change their photo-physical properties (λ_{max} shift of the UV/Vis spectrum, change in the quantum yield or emission wavelength, etc) in the presence of the analyte.^[1,2] Ratiometric fluorescent probes, in particular, have found widespread use in biological, polymeric, and sensory materials chemistry, where the two emission bands from the covalently linked chromophores, or a chromophore with a dual channel or with a dual signaling pathway, were used to differentiate between the two signals. In the former case, the analyte recognition site was connected to two chromophores (an energy donor and an energy acceptor or chromophores forming an excimer/exciple) through a covalent bond, and the analyte changes the orientation and/or the dipolar interaction between the chromophores.^[2,3] In the latter cases, another signaling channel or signaling pathway is activated in the analyte binding site (which is frequently a part of a chromophore) only when the analyte is bound to the chromophore.^[3c,f] In these systems, the chromophores were designed so that the two emission changes would occur in the presence of the analyte. It thus occurred to us that if the analyte induced multiple emissions^[4] with different colors from a chemosensor, the perceived color change would be useful not only for the ratiometric method of detection but also for rapid

visual sensing. We report herein that a triarylborane–porphyrin conjugate **1** displays colorimetric (purple to green and red to bluish emission) and ratiometric fluorescent responses to fluoride ions with three emission bands at 356, 670, and 692 nm.^[5–9]



The triarylborane chromophore (an energy donor: D) and the porphyrin chromophore (an energy acceptor: A) in **1** share a Lewis acidic boron junction that can bind a fluoride ion with subsequent $\text{sp}^2\text{-sp}^3$ hybridization of the boron atom.^[10] The change in the hybridization of the boron atom interrupts both the electronic communication and the dipolar interaction between the triarylborane and porphyrin units as well as the internal charge-transfer state between the two chromophores so that each chromophore (D' and A') absorbs and emits light independently only in the presence of fluoride ions (Figure 1).^[11] This situation could lead to changes in the UV/Vis and fluorescence spectra, from which one could sense the fluoride ion colorimetrically and ratiometrically.


Compound **1** was prepared according to Scheme 1 and identified by ^1H NMR spectroscopy, MALDI-TOF mass spectrometry, and elemental analysis. Compounds **2** and **3**^[10b] were synthesized for control experiments. The Soret band of **1** ($[\text{1}] = 3.00 \mu\text{M}$) in THF appears at 436.5 nm, that is, slightly red-shifted relative to that of **2** (430.0 nm), whereas the absorption maxima of the triarylborane moiety (303.0, 322.5, and 392.5 nm) in **1** are identical with those of **3** (Figure 2 A). No band attributable to electronic transitions between mixed molecular levels of the triarylborane and porphyrin in **1** was observed. The similarity between the UV/Vis spectrum of **1** and the additive sum of **2** and **3** indicates that the electronic coupling between the two chromophores in the ground state is negligibly small.

[*] Dr. M. Takeuchi, Prof. S. Shinkai, Y. Kubo, Dr. M. Yamamoto, Dr. M. Ikeda
Department of Chemistry and Biochemistry
Graduate School of Engineering, Kyushu University
Fukuoka 812-8581 (Japan)
Fax: (+81) 92-642-3611
E-mail: taketcm@mbox.nc.kyushu-u.ac.jp
seijitcm@mbox.nc.kyushu-u.ac.jp

Dr. S. Yamaguchi, Prof. K. Tamao
Institute for Chemical Research, Kyoto University
Gokasho, Uji, Kyoto 611-0011 (Japan)

[**] Y.K. and M.Y. contributed equally to this work. We thank Dr. Midori Watanabe and Ms. Hongyue Li at Kyushu University for ^1H NMR measurements. M.T. and Y.K. thank prof. H. Yonemura for helpful discussions. M.I. thanks the JSPS Research Fellowship for Young Scientists for financial support. This work was partially supported by a grant-in-aid for COE research ("Design and Control of Advanced Molecular Assembly Systems") from the Ministry of Education, Science, and Culture, Japan (no. 08CE2005).

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



linked porphyrin moiety. In contrast to the porphyrin moiety, a decrease (to 392.5 nm) and an increase (to 323.0 and 343.4 nm) in the intensities of the triarylborane absorption bands in **1** were observed. These trends are the same as those observed for **3**, and the result is interpreted as a loss of π conjugation through the boron atom.^[9] These results support the view that the two chromophores, the aryl borane (D') and the porphyrin (A'), generate their absorption bands independently in the presence of fluoride ions. The change in the absorption spectra resulted in a change in the color of the solution from purple to green (Figure 3). The stoichiometry in the reaction between **1** and TBAF was confirmed to be 1:1




Figure 3. Color changes observed in samples of **1** and TBAF in THF. Left to right: **1**, **1**-TBAF, visible emission of **1**, and emission of **1**-TBAF.

from a plot of the mole ratio versus absorption. A standard curve-fitting method was utilized for the $A_{436.5}$ versus [TBAF] plot to evaluate the association constant for the 1:1 complex, which was estimated to be 99700 M^{-1} . This value is comparable with those previously determined for triarylboranes and fluoride ions.^[10] No change in the Soret band, Q band, or the tridurylborane absorption was observed upon exposure of **1** to larger halides, acetate, and hydroxide ions. Apparently, only the smaller anion, the fluoride ion, can fit in the space around the triduryl-surrounded Lewis acidic boron atom.

The coordination of the fluoride ion to the boron center was confirmed by NMR spectroscopy. A *meso*-H proton signal and pyrrole β -proton signals of the porphyrin unit as well as methyl proton signals of the duryl group are shifted to higher magnetic fields in the ^1H NMR spectra of [**1**]:[TBAF] = 1:0 and 1:2 in $[\text{D}_8]\text{THF}$ at 25°C ([**1**] = 1.00 mM; see the Supporting Information). A ^{19}F signal was shifted to a lower magnetic field ($\Delta\delta = 8.40$ ppm) relative to free TBAF in the ^{19}F NMR spectra as a result of its coordination to the boron center (see the Supporting Information).

The fluorescence spectrum of **1** ($0.5\ \mu\text{M}$) in THF was recorded with excitation at an isosbestic point (294 nm, see the Supporting Information) where the selectivity of the triarylborane absorption over that of the porphyrin ring was 3:1, as estimated from the absorbance of compounds **2** and **3**. Compound **1** gives the porphyrin emission band (670 nm) without the triarylborane emission band that should appear at 515 nm, as seen in the emission spectrum of **3** (see the Supporting Information). The same spectrum was obtained in a nonpolar solvent (benzene). The intermolecular energy transfer is negligible at such a low concentration, since the

fluorescence spectrum for an equimolar mixture ($0.5\ \mu\text{M}$ each) of **2** and **3** was equivalent to the sum of the two individual fluorescence spectra. Furthermore, the excitation spectrum of **1** shows an intense triarylborane absorption band as well as the porphyrin band. The findings clearly show that efficient transfer of energy from the triarylborane to the porphyrin chromophore takes place in the absence of fluoride ions.

The addition of fluoride ions to a solution of **1** in THF results in a substantial increase in the intensity of the two emission bands at 356 nm and 692 nm along with a decrease in the intensity of the emission band at 670 nm (Figure 4). The




Figure 4. Changes in the emission spectra upon addition of TBAF to solutions of THF at 25°C ; excitation wavelength 294 nm.

appearance of the emission band at 356 nm was the same response to fluoride ions as that of **3**. The longer wavelength shift of 22 nm in the porphyrin emission can be attributed to the lower energy of the Q band in the ground state. A visible change in the emission from a red to bluish color was also observed (Figure 3). Compound **1** showed a ratiometric response to fluoride ions at two different wavelengths (356 and 692 nm) when normalized against the emission at 670 nm (see the Supporting Information). The excitation spectra of **1** with added fluoride ions revealed that the absorbance of the aryl borane still participates in the appearance of the porphyrin emission at 692 nm. This observation means that the energy transfer from the aryl borane moiety to the porphyrin unit occurs even after complexation with a fluoride ion.^[14] The fluorescence decay of the emission band of the aryl borane moiety in **1** was evaluated to be shorter than 100 ps (515 nm) and 0.53 ns (380 nm) without and with the guest fluoride ion, respectively. A shorter lifetime at the 515 nm emission without fluoride ions than that of **3** (4.52 ns) could be attributed to an efficient Dexter-type energy transfer from the aryl borane to the porphyrin moiety in **1**. In the presence of fluoride ions, the lifetime of 0.53 ns is similar to that of the **3**-fluoride ion complex (monitored at 380 nm, 0.39 ns). This finding together with the excitation spectrum of the **1**-fluoride ion complex enables one to suppose that energy transfer from the aryl borane moiety to the porphyrin unit in **1** takes place to some extent in the presence of fluoride ions. The result suggests, therefore, that the guest fluoride ion can switch the energy-transfer pathway to make it possible to monitor the

binding event through the changes in the three emissions (Figure 5).

In conclusion, we have demonstrated that **1** displays a colorimetric (purple to green and red to bluish emission) and a ratiometric response (by utilizing changes in three emis-




Figure 5. Effects of the coordination of a fluoride ion on the emissions from **1** on excitation at 294 nm.

sions) that are useful for the detection of fluoride ions. The mechanism for the increase in the two emission bands and not the third emission band was assumed by the finding that the coordination of a fluoride ion to the boron center in **1** caused perturbation in the energy transfer pathway and the π conjugation. Thus, two chromophores sharing a recognition unit linked though a conjugate spacer would become a versatile molecular design scheme to achieve the monitoring of multiple emissions useful for signal differentiation.

Experimental Section

All starting materials and solvents were purchased from Tokyo Kasei Chemicals or Wako Chemicals, and used as received. The ^1H NMR spectra were recorded either on a Bruker AC 250 (250 MHz) or Bruker DRX 600 (600 MHz) spectrometer. Chemical shifts are reported in ppm downfield from tetramethylsilane as the internal standard. Mass spectral data were obtained by using a Perseptive Voyager RP MALDI-TOF spectrometer. UV/Vis and fluorescent spectra were recorded with a Shimadzu UV-2500 PC and Hitachi F-4500 spectrophotometer.

Received: December 17, 2002 [Z50788]

Keywords: fluorescent probes · fluorides · molecular recognition · porphyrinoids · sensors

- [1] a) A. P. de Silva, D. B. Fox, A. J. M. Huxley, T. S. Moody, *Coord. Chem. Rev.* **2000**, 205, 41; b) L. Prodi, F. Bolletta, M. Montaliti, N. Zaccaroni, *Coord. Chem. Rev.* **2000**, 205, 59; c) V. Amendola, L. Fabbrizzi, M. Licchelli, C. Mangano, P. Pallavicini, L. Parodi, A. Poggi, *Coord. Chem. Rev.* **2000**, 192, 649; d) J. P. Desvergne, A. W. Czarnik, *NATO ASI Ser. Ser. C* **1997**, 492.
- [2] J. R. Lakowicz, *Principles of Fluorescence Spectroscopy*, 2nd ed., Plenum, New York, **1999**.

- [3] For selected recent examples, see a) S. Maruyama, K. Kikuchi, T. Hirano, Y. Urano, T. Nagano, *J. Am. Chem. Soc.* **2002**, 124, 10650; b) J. Raker, T. E. Glass, *J. Org. Chem.* **2002**, 67, 6113; c) F. Würthner, S. Ahmed, C. Thalacker, T. Debaerdemaeker, *Chem. Eur. J.* **2002**, 8, 4742; d) J. Kim, D. T. McQuade, A. Rose, Z. Zhu, T. M. Swager, *J. Am. Chem. Soc.* **2001**, 123, 11488; e) J. V. Mello, N. S. Finney, *Angew. Chem.* **2001**, 113, 1584; *Angew. Chem. Int. Ed.* **2001**, 40, 1536; f) K. Choi, A. D. Hamilton, *Angew. Chem.* **2001**, 113, 4030; *Angew. Chem. Int. Ed.* **2001**, 40, 3912; g) J.-S. Yang, C.-S. Lin, C.-Y. Hwang, *Org. Lett.* **2001**, 3, 889; h) S. Deo, H. A. Godwin, *J. Am. Chem. Soc.* **2000**, 122, 174; i) A. P. de Silva, J. Eilers, G. Zlokarnik, *Proc. Natl. Acad. Sci. USA* **1999**, 96, 8336.
- [4] a) Y.-A. Lee, J. E. McGarrah, R. J. Lachicotte, R. Eisenberg, *J. Am. Chem. Soc.* **2002**, 124, 10662; b) H. Fu, B. H. Loo, D. Xiao, R. Xie, X. Ji, J. Yao, B. Zhang, L. Zhang, *Angew. Chem.* **2002**, 114, 1004; *Angew. Chem. Int. Ed.* **2002**, 41, 962.
- [5] For reviews on anion receptors, see a) P. D. Beer, P. A. Gale, *Angew. Chem.* **2001**, 113, 502; *Angew. Chem. Int. Ed.* **2001**, 40, 486; b) P. A. Gale, *Coord. Chem. Rev.* **2000**, 199, 181; c) T. S. Snoeden, E. V. Anslyn, *Curr. Opin. Chem. Biol.* **1999**, 3, 740; d) P. D. Beer, *Acc. Chem. Res.* **1998**, 31, 71; e) F. P. Schmidtchen, M. Berger, *Chem. Rev.* **1997**, 97, 1609; e) J. Scheerder, J. F. J. Engbersen, D. N. Reinhoudt, *Recl. Trav. Chim. Pays-Bas* **1996**, 115, 307.
- [6] For recent anion sensors, see a) P. Piatek, J. Jurczak, *Chem. Commun.* **2002**, 2450; b) J.-H. Liao, C.-Y. Chen, J.-M. Fang, *Org. Lett.* **2002**, 4, 561; c) A. Ojida, Y. Mito-oka, M. Inoue, I. Hamachi, *J. Am. Chem. Soc.* **2002**, 124, 6256; d) J. R. Blas, M. Marquez, J. L. Sessler, F. J. Luque, M. Orozco, *J. Am. Chem. Soc.* **2002**, 124, 12796; e) J. L. Sessler, H. Maeda, T. Mizuno, V. M. Lynch, H. Furuta, *Chem. Commun.* **2002**, 862; f) K. Choi, A. D. Hamilton, *J. Am. Chem. Soc.* **2001**, 123, 2456; g) D. H. Lee, K. H. Lee, J. I. Hong, *Org. Lett.* **2001**, 3, 5; h) P. D. Beer, N. Berry, M. G. B. Drew, O. D. Fox, M. E. Padilla-Tosta, S. Patell, *Chem. Commun.* **2001**, 199; i) M. J. Deetz, M. Shang, B. D. Smith, *J. Am. Chem. Soc.* **2000**, 122, 6201; j) Y. Kubo, M. Tsukahara, S. Ishihara, S. Tokita, *Chem. Commun.* **2000**, 653; k) S. Sasaki, M. Mizuno, K. Naemura, Y. Tobe, *J. Org. Chem.* **2000**, 65, 275; l) H. Miyaji, P. Anzenbachlar, Jr., J. L. Sessler, E. R. Bleasdale, P. A. Gale, *Chem. Commun.* **1999**, 1723.
- [7] Receptors for fluoride ions are of special importance for monitoring fluoride metabolism in nature, the analysis of drinking water, and the detection of chemical warfare agents.^[9]
- [8] For recent fluoride chemosensors, see a) J. L. Sessler, H. Maeda, T. Mizuno, V. M. Lynch, H. Furuta, *J. Am. Chem. Soc.* **2002**, 124, 13474; b) S. Aldridge, C. Bresner, I. A. Fallis, S. J. Colesb, M. B. Hursthouse, *Chem. Commun.* **2002**, 740; c) S. K. Kim, J. Yoon, *Chem. Commun.* **2002**, 770; d) T. Mizuno, W.-H. Wei, L. R. Eller, J. L. Sessler, *J. Am. Chem. Soc.* **2002**, 124, 1134; e) T. Gunnlaugsson, A. P. Davis, M. Glynn, *Chem. Commun.* **2001**, 2556; f) K. H. Lee, H.-Y. Lee, D. H. Lee, J.-I. Hong, *Tetrahedron Lett.* **2001**, 42, 5447; g) H. Furuta, H. Maeda, A. Osuka, *J. Am. Chem. Soc.* **2001**, 123, 6435; h) M. Takeuchi, T. Shioya, T. M. Swager, *Angew. Chem.* **2001**, 113, 3476; *Angew. Chem. Int. Ed.* **2001**, 40, 3372; i) C. R. Cooper, N. Spencer, T. D. James, *Chem. Commun.* **1998**, 1365.
- [9] H. Sohn, S. Létant, M. J. Sailor, W. C. Trogler, *J. Am. Chem. Soc.* **2000**, 122, 5399.
- [10] a) S. Yamaguchi, S. Akiyama, K. Tamao, *J. Am. Chem. Soc.* **2000**, 122, 6335; b) S. Yamaguchi, T. Shirasaka, K. Tamao, *Org. Lett.* **2000**, 2, 4129; c) S. Yamaguchi, S. Akiyama, K. Tamao, *J. Am. Chem. Soc.* **2001**, 123, 11372; d) S. Yamaguchi, T. Shirasaka, S. Akiyama, K. Tamao, *J. Am. Chem. Soc.* **2002**, 124, 8816.
- [11] The change in electron-transfer efficiency caused by the binding of the fluoride ion to a boronate ester has already been reported, see H. Shiratori, T. Ohno, K. Nozaki, A. Osuka, *Chem. Commun.* **1999**, 2181.

- [12] It was confirmed by ^1H NMR spectroscopy that **1** does not aggregate in THF both in the presence and in the absence of TBAF.
- [13] M. J. Crossley, L. A. Johnston, *Chem. Commun.* **2002**, 1122.
- [14] The emission intensity of **1** excited at 294 nm is 13 (at 670 nm) and 16 (at 692 nm) times larger than that of **2** in the absence and in the presence of fluoride ions, respectively, if it is assumed that the energy transfer occurs from the aryl borane moiety to the porphyrin ring.