


## Cationic and Neutral Four-Coordinate Alkylidene Complexes of Vanadium(IV) Containing Short V=C Bonds\*\*

Falguni Basuli, Uriah J. Kilgore, Xile Hu, Karsten Meyer, Maren Pink, John C. Huffman, and Daniel J. Mindiola\*

High-oxidation state transition-metal alkylidenes have received considerable attention in the past few years since these systems are capable of catalyzing important processes such as olefin metathesis, alkyne polymerization, and Wittig-type reactions.<sup>[1,2]</sup> Although transition-metal alkylidene complexes of the nucleophilic “Schrock-type” have been extensively studied for the 4d and 5d metals of Group 5, the chemistry for 3d vanadium systems has been far less explored. The first example of a vanadium(III)–alkylidene complex, [CpV(CH*t*Bu)(dmpe)] (dmpe = bis(dimethylphosphanyl)ethane), was reported by Hessen, Teuben, and co-workers and involved  $\alpha$ -hydrogen abstraction of the bis(alkyl) precursor.<sup>[3]</sup> In contrast to the heavier  $d^0$  alkylidene congeners, which are prepared by  $\alpha$ -hydrogen abstraction, the synthesis of the first vanadium(V)–alkylidene complex [CpV(NAr)(CHPh)(PMe<sub>3</sub>)] (Ar = 2,6-(CHMe<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>) involved a two-electron oxidation of the precursor [CpV(NAr)(PMe<sub>3</sub>)<sub>2</sub>] with the Wittig reagent Ph<sub>3</sub>P=CHPh by alkylidene transfer.<sup>[4]</sup> Since there are only a handful of vanadium–alkylidene complexes known,<sup>[3–6]</sup> we report the synthesis of the first cationic and neutral four-coordinate vanadium(IV)–neopentylidene complexes [(Nacnac)V=CH*t*Bu(thf)](BPh<sub>4</sub>) and [(Nacnac)V=CH*t*Bu(I)] (Nacnac<sup>−</sup> = [Ar]NC(Me)CHC(Me)N[Ar], Ar = 2,6-(CHMe<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>), and subsequent thermolysis of each system. These  $d^1$  paramagnetic species reported herein are kinetically stable, and contain the shortest V=C bonds reported so far.

Our recent strategy to prepare the first four-coordinate titanium–neopentylidene complex by an oxidatively induced  $\alpha$ -hydrogen abstraction reaction<sup>[7]</sup> motivated us to pursue other 3d transition metals containing this reactive motif.

Unlike titanium, vanadium displays more diverse redox chemistry, hence oxidation states ranging from +2 to +5 can occur. Using the precursor [(Nacnac)VCl<sub>2</sub>]<sup>[8]</sup> developed by Budzelaar and co-workers and two equivalents of LiCH<sub>2</sub>*t*Bu,<sup>[1g]</sup> we prepared the corresponding bis(neopentyl) complex [(Nacnac)V(CH<sub>2</sub>*t*Bu)<sub>2</sub>] (**1**) in 80 % yield as dark brown needles (Scheme 1). Attempts to obtain crystals suitable for a



**Scheme 1.** Synthesis of the vanadium(IV)–alkylidene complexes **2** and **4**.

single-crystal X-ray structure determination were unsuccessful, but elemental analysis, magnetic susceptibility measurements ( $\mu_{\text{eff}} = 2.96 \mu_{\text{B}}$ ), and reactivity (vide infra) are in agreement with the formation of **1**.<sup>[9]</sup> Complex **1** is a close analogue of the *n*Bu derivative prepared by Budzelaar and co-workers, differing only by the lack of  $\beta$ -hydrogen atoms.<sup>[8]</sup> Cyclic voltammograms of **1** (TBAH, THF; TBAH = tetra-*n*-butylammonium hexafluorophosphate) revealed reversible oxidation and reduction waves at  $-0.35$  and  $-2.68$  V, respectively, which correspond to the V<sup>III</sup>/V<sup>IV</sup> and V<sup>III</sup>/V<sup>II</sup> redox couple (referenced versus [FeCp<sub>2</sub>]/[FeCp<sub>2</sub>]<sup>+</sup>).<sup>[9]</sup> Chemical oxidation of **1** with AgBPh<sub>4</sub> leads to formation of a Ag<sup>0</sup> mirror concurrent with the vanadium(IV)–alkylidene complex [(Nacnac)V=CH*t*Bu(thf)](BPh<sub>4</sub>) (**2**) in 74 % yield (Scheme 1). Complex **2** is stable in the solid state, but its solutions (Et<sub>2</sub>O, C<sub>6</sub>H<sub>6</sub>, THF) gradually decompose over several hours at room temperature.


Dark brown single crystals of **2** were grown from THF solutions of the complex layered with Et<sub>2</sub>O at  $-35^\circ\text{C}$ . The structure of **2** (Figure 1) reveals a cationic and four-coordinate vanadium center with pseudotetrahedral geometry, a very short V=C bond length (V(1)–C(33) 1.795(3) Å), and an obtuse V(1)–C(33)–C(34) angle of  $159.8(4)^\circ$ .<sup>[9,10]</sup> The highly distorted neopentylidene ligand suggests that there is a significant  $\alpha$ -agostic CH interaction with the metal center. Since **2** is a  $d^1$  complex,  $\alpha$ -elimination is inhibited, and the complex distorts to form an unusually short V=C bond. The vanadium center deviates from the N(2)–N(6)–C(33) mean plane by 0.592(6) Å, and as consequence of the cationic nature of the metal center, a coordinated molecule of THF occupies the fourth site. EPR ( $g_{\text{iso}} = 1.982$ ,  $A_{\text{iso}} = 88 \times 10^{-4} \text{ cm}^{-1}$ , <sup>51</sup>V,  $I = 7/2$ , 99.6 %, Figure 2) and magnetic susceptibility measurements (SQUID,  $\mu_{\text{eff}} = 1.85 \mu_{\text{B}}$ ; Evans’,

[\*] Dr. F. Basuli, U. J. Kilgore, M. Pink, Dr. J. C. Huffman, Prof. Dr. D. J. Mindiola  
Department of Chemistry and Molecular Structure Center  
Indiana University, Bloomington, IN 47405 (USA)  
Fax: (+1) 812-855-8300  
E-mail: mindiola@indiana.edu

X. Hu, Prof. Dr. K. Meyer  
Department of Chemistry and Biochemistry, Mail Code 0358  
University of California, San Diego  
9500 Gilman Drive, La Jolla, CA 92093-0358 (USA)


[\*\*] This work was supported by Indiana University, the Camille and Henry Dreyfus Foundation, and the Ford Foundation. U.J.K was supported by NSF Grant CHE-0097388 under the auspices of the Research Experience for Undergraduates Program at Indiana University.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



**Figure 1.** Molecular structure of complexes [(Nacnac)V=CH*t*Bu(thf)]<sup>+</sup>(BPh<sub>4</sub>)<sup>−</sup> (**2**) and [(Nacnac)V=CH*t*Bu(I)] (**4**) with thermal ellipsoids at the 50% probability level. Hydrogen atoms with the exception of the alkylidene carbon atoms (C(33), **2**; C(34), **4**), and aryl groups on the nitrogen atoms with the exception of the *ipso*-carbon atoms (C(7), C(21), C(8), C(22)) are omitted for clarity. The [BPh<sub>4</sub>]<sup>−</sup> ion and two disordered THF solvent molecules are omitted for clarity in the structure of **2**. Selected bond lengths [Å] and angles [°]: for **2**: V(1)–C(33) 1.795(3), V(1)–N(2) 1.942(3), V(1)–N(6) 1.964(3), V(1)–O(38) 2.000(2); V(1)–C(33)–C(34) 159.8(4), O(38)–V(1)–C(33) 101.87(13), C(33)–V(1)–N(6) 112.82(15), C(33)–V(1)–N(2) 122.03(15), N(6)–V(1)–N(2) 96.12(11). For **4**: V(1)–C(34) 1.787(3), V(1)–N(3) 1.989(2), V(1)–N(7) 1.935(2), V(1)–I(2) 2.6478(6); V(1)–C(34)–C(35) 158.7(3), I(2)–V(1)–C(34) 110.65(12), C(34)–V(1)–N(3) 107.08(12), C(34)–V(1)–N(7) 123.41(15), N(7)–V(1)–N(3) 94.93(9).


$\mu_{\text{eff}} = 1.87 \mu_{\text{B}}$ ) are in accordance with a  $d^1$  paramagnetic species and unambiguously rule out the possibility that oxidation of a V–C bond in **1** occurred instead to give the V<sup>III</sup> cation [(Nacnac)V(CH*t*Bu)(thf)]<sup>+</sup>. Similar species have been prepared by protonation reactions of the corresponding bis(alkyl) systems.<sup>[11]</sup> The role of the anion in the oxidation of **1** to form **2** is key in the oxidatively induced  $\alpha$ -hydrogen abstraction process inasmuch as oxidation of **1** with AgOTf oxidizes the vanadium–carbon bond instead of the vanadium center. Hence, the V<sup>III</sup> complex [(Nacnac)V(CH*t*Bu)(OTf)] (**3**) is formed in good yields (78%) when **1** is treated with AgOTf. Elemental analysis, solution magnetic susceptibility data, and single-crystal X-ray diffraction studies are consistent with the connectivity proposed for complex **3**.<sup>[9]</sup> Complex **2** can undergo smooth anion exchange with 0.5 equivalents of I<sub>2</sub> (47% yield) or excess MgI<sub>2</sub> (69% yield) to produce the neutral, four-coordinate vanadium–alkylidene complex [(Nacnac)V=CH*t*Bu(I)] (**4**) (Scheme 1). In the reaction of **2** with I<sub>2</sub>, we also detect, by <sup>11</sup>B NMR spectroscopy, the formation of BPh<sub>3</sub> as a side product. This observation suggests that I<sub>2</sub> might be also oxidizing the B–C(phenyl) bond of the BPh<sub>4</sub><sup>−</sup> counter anion to afford **4**. Magnetic susceptibility measurements (Evans',  $\mu_{\text{eff}} = 1.94 \mu_{\text{B}}$ ), single-crystal X-ray diffraction study, and solution EPR ( $g_{\text{iso}} = 1.976$ ,  $A_{\text{iso}} = 64 \times 10^{-4} \text{ cm}^{-1}$ , <sup>51</sup>V,  $I = 7/2$ , 99.6%, Figure 2) spectra are consistent with the  $d^1$  vanadium(IV) complex **4** containing a terminal alkylidene functionality. Cyclic voltammograms of **4** (TBAH, THF) revealed irreversible oxidation and reduction waves at −0.18 and −2.28 V, respectively (referenced versus [FeCp<sub>2</sub>]/[FeCp<sub>2</sub>]<sup>+</sup>).<sup>[9]</sup> This suggests that on the cyclic voltammetry time scale the putative vanadium(III) and vanadium(V) systems of



**Figure 2.** X-band EPR spectrum of **2** (a) and **4** (b) recorded at room temperature in toluene. Experimental spectra (exp.) are above the simulated spectra (sim.).


**4**, generated by one-electron reduction and one-electron oxidation, respectively, appear to be unstable. In addition, chemical reduction and oxidation of complex **4** resulted in decomposition products. The molecular structure of **4** is depicted in Figure 1 and reveals the shortest V=C<sub>α</sub>{alkylidene} bond reported so far (V(1)–C(34) 1.787(3) Å) as well as an obtuse V–C<sub>α</sub>–C<sub>β</sub> angle (V(1)–C(34)–C(35), 158.7(3)°).<sup>[12]</sup> Complexes **2** and **4** are isostructural, where the bound THF molecule and cationic charge in **2** are replaced with I<sup>−</sup> to generate the neutral complex **4**.

Owing to the instability of the previously reported four-coordinate alkylidene complex [(Nacnac)Ti=CH*t*Bu(OTf)],<sup>[7]</sup> it was presaged that complexes **2** and **4** would also be prone to “Wittig-like” reactivity involving the metal–alkylidene functionality. Accordingly, when complex **2** is heated at 60 °C for 6 h in THF, extrusion of neopentane is observed (by <sup>1</sup>H NMR spectroscopy and GC MS analysis) concomitant with the formation of the zwitterion vanadium(II) product [(Nacnac)V(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>BPh<sub>3</sub>)] (**5**) (77% yield, Scheme 2). THF is dehydrogenated during the thermolysis inasmuch as [D<sub>2</sub>]neopentane is observed when the reaction is carried out in [D<sub>8</sub>]THF. Complex **5** is an orange, paramagnetic species, and the X-band EPR solution spectra in toluene, as well as solution and solid-state magnetic susceptibility measurements are in accordance with the molecule having a quartet ground



**Scheme 2.** Thermolysis of complexes **2** and **4** to prepare **5** and **6**, respectively.

state spin configuration (SQUID,  $\mu_{\text{eff}} = 3.93 \mu_{\text{B}}$ ; Evans',  $\mu_{\text{eff}} = 3.68 \mu_{\text{B}}$ ).<sup>[9]</sup> The structure determined for **5** by using suitable single crystals also substantiates the proposed connectivity and illustrates clearly a low-valent, two-coordinate (Nacnac)-V<sup>II</sup> scaffold coordinating to one of the phenyl groups attached to the borate (Figure 3).<sup>[13]</sup> In contrast to the transformation



**Figure 3.** Molecular structure of complexes  $[(\text{Nacnac})\text{V}(\eta^6\text{-C}_6\text{H}_5\text{BPh}_4)]$  (**5**) and  $[(\text{Ar})\text{NC}(\text{Me})\text{CHC}(\text{Me})\text{CH}t\text{Bu})\text{V}=\text{NAr}(\text{I})]$  (**6**) with thermal ellipsoids at the 50% probability level. Hydrogen atoms with the exception of the olefinic carbon atom (C(4), **6**), and aryl groups on the nitrogen atoms with the exception of the *ipso*-carbon atoms (C(7), C(21), C(5), C(5)) are omitted for clarity. Selected bond lengths [Å] and angles [°]: For **5**: V(1)–C(34) 2.445(2), V(1)–C(35) 2.363(2), V(1)–C(36) 2.340(2), V(1)–C(37) 2.322(2), V(1)–C(38) 2.353(2), V(1)–C(39) 2.383(2), V(1)–N(6) 2.066(8), V(1)–N(2) 2.082(8); N(6)–V(1)–N(2) 90.89(7). For **6**: V(1)–N(1) 1.985(7), V(1)–N(2) 1.687(8), V(1)–I(1) 2.6589(4), V(1)–C(4) 2.219(2), V(1)–C(3) 2.387(2), C(4)–C(3) 1.404(3); C(3)–C(4)–C(19) 127.2(9), I(1)–V(1)–C(4) 125.89(6), N(2)–V(1)–N(1) 112.69(8), C(4)–V(1)–N(2) 100.42(8), C(4)–V(1)–N(1) 89.76(8).

of **2** to **5**, when a solution of **4** in benzene is thermolyzed for 6 h at 60 °C, “Wittig-like” reactivity is observed, which is evidenced by the formation of the vanadium(IV)–imide complex  $[(\text{Ar})\text{NC}(\text{Me})\text{CHC}(\text{Me})\text{CH}t\text{Bu})\text{V}=\text{NAr}(\text{I})]$  (**6**) (57% yield; Scheme 2). Solution EPR and magnetic susceptibility measurements are consistent with a  $d^1$  system.<sup>[9]</sup> In addition, the molecular structure of **6** was obtained from single-crystal X-ray diffraction studies and disclosed a low-coordinate vanadium(IV) center supported by an imide, and an anilide with a  $\eta^2$ -coordinated diene pendant arm.<sup>[14]</sup> The low-coordination environment at the metal center promotes  $\eta^2$  coordination of the ene motif, which is now part of the former Nacnac<sup>−</sup> ligand (Figure 3). Similar transformations have been reported by our group for four-coordinate  $[(\text{Nacnac})\text{Ti}^{\text{IV}}]$  complexes with reactive alkylidene or phosphinidene functionalities.<sup>[7,15]</sup> Concentration-dependent experiments on the titanium–alkylidene system (by <sup>1</sup>H NMR spectroscopy) determined the cross-metathesis reaction to be first-order in titanium, hence we suspect the same to be valid for the thermolytic transformation of **4** to **6**.

In summary, the cationic and neutral four-coordinate complexes (**2** and **4**, respectively) derived from the one-electron oxidation of the bis(alkyl)vanadium(III) precursor **1** represent a new class of alkylidene systems in the context of the organometallic chemistry of vanadium. Further applications of these unique  $d^1$  metal radicals of vanadium containing metal–ligand multiple bonds to reactions such as group- or atom-transfer radical polymerization (ATRP),<sup>[16]</sup> and olefin metathesis will be described in due course.

## Experimental Section

Synthetic and characterization details for compounds **1–6** (including single-crystal X-ray data for **2–6**)<sup>[17]</sup> are described in the Supporting Information.

Received: October 31, 2003 [Z53210]

**Keywords:** alkylidenes · carbenes · N ligands · oxidation · vanadium

- a) R. R. Schrock, *Acc. Chem. Res.* **1979**, *12*, 98–104; b) R. R. Schrock, *Acc. Chem. Res.* **1990**, *23*, 158–165; c) R. R. Schrock, *Chem. Rev.* **2002**, *102*, 145–179; d) R. R. Schrock, *Reactions of Coordinated Ligands* (Ed.: P. R. Braterman), Plenum, New York, **1986**; e) J. Feldman, R. R. Schrock, *Prog. Inorg. Chem.* **1991**, *39*, 1–74; f) R. R. Schrock, *J. Organomet. Chem.* **1986**, *300*, 249–262; g) R. R. Schrock, J. D. Fellmann, *J. Am. Chem. Soc.* **1978**, *100*, 3359–3370; h) R. R. Schrock, P. R. Sharp, *J. Am. Chem. Soc.* **1978**, *100*, 2389–2399; i) G. A. Rupprecht, L. W. Messerle, J. D. Fellmann, R. R. Schrock, *J. Am. Chem. Soc.* **1980**, *102*, 6236–6244; j) D. N. Clark, R. R. Schrock, *J. Am. Chem. Soc.* **1978**, *100*, 6774–6776; k) J. D. Fellmann, G. A. Rupprecht, C. D. Wood, R. R. Schrock, *J. Am. Chem. Soc.* **1978**, *100*, 5964–5966.
- a) A. M. Rouhi, *Chem. Eng. News* **2002**, *80*, 29–38; b) A. Fürstner, *Adv. Synth. Catal.* **2002**, *344*, 567; c) R. R. Schrock, *Adv. Synth. Catal.* **2002**, *344*, 571–572.
- a) B. Hessen, A. Meetsma, J. H. Teuben, *J. Am. Chem. Soc.* **1989**, *111*, 5977–5978; b) B. Hessen, J.-K. F. Buijink, A. Meetsma,

- J. H. Teuben, G. Helgesson, M. Håkansson, S. Jagner, A. L. Spek, *Organometallics* **1993**, *12*, 2268–2276.
- [4] J.-K. F. Buijink, J. H. Teuben, H. Kooijman, A. L. Spek, *Organometallics* **1994**, *13*, 2922–2924.
- [5] a) B. Hessen, A. Meetsma, F. van Bolhuis, J. H. Teuben, G. Helgesson, S. Jagner, *Organometallics* **1990**, *9*, 1925–1936; b) M. Moore, S. Gambarotta, G. Yap, L. M. Liable-Sands, A. L. Rheingold, *Chem. Commun.* **1997**, 643–644; c) R. Milczarek, W. Russeler, P. Binger, K. Jonas, K. Angermund, C. Kruger, M. Regitz, *Angew. Chem.* **1987**, *99*, 957–958; *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 908–909.
- [6] For examples of vanadium(II) carbenes see: a) G. Erker, R. Lecht, R. Schlund, K. Angermund, C. Kruger, *Angew. Chem.* **1987**, *26*, 708–710; *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 666–668; b) G. Erker, R. Lecht, R. Schlund, K. Angermund, C. Kruger, **1987**, *99*, 708–709; c) M. Berlekamp, G. Erker, J. L. Petersen, *J. Organomet. Chem.* **1993**, *458*, 97–103; d) M. Grehl, M. Berlekamp, G. Erker, *Acta Crystallogr. Sect. C* **1995**, *51*, 1772–1774.
- [7] F. Basuli, B. C. Bailey, J. Tomaszewski, J. C. Huffman, D. J. Mindiola, *J. Am. Chem. Soc.* **2003**, *125*, 6052–6053.
- [8] P. H. M. Budzelaar, A. B. von Oort, A. G. Orpen, *Eur. J. Inorg. Chem.* **1998**, 1485–1494.
- [9] See Supporting Information for complete experimental, spectral, and crystallographic details.
- [10] Crystallographic details for **2·2 THF**: A dark crystal of approximate dimensions  $0.30 \times 0.30 \times 0.06 \text{ mm}^3$  was selected and mounted on a glass fiber. A total of 88763 reflections ( $-54 \leq h \leq 52$ ,  $-20 \leq k \leq 20$ ,  $-24 \leq l \leq 24$ ) was collected at  $T = 136(2) \text{ K}$  in the range of  $2.27$  to  $27.46^\circ$ , of which 8308 were observed ( $R_{\text{int}} = 0.0957$ );  $\text{MoK}\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). A direct-methods solution was calculated which provided most non-hydrogen atoms from the E-map. All non-hydrogen atoms were refined with anisotropic displacement parameters. The residual peak and hole electron densities were  $1.692$  and  $-0.604 \text{ e \AA}^{-3}$ , respectively. The absorption coefficient was  $0.213 \text{ mm}^{-1}$ . The least-squares refinement converged normally with residuals of  $R(F) = 0.0733$ ,  $wR(F^2) = 0.2286$ , and a GOF =  $0.934$  ( $I > 2\sigma(I)$ ).  $\text{C}_{70}\text{H}_{95}\text{BN}_2\text{O}_3\text{V}$ , space group  $C2/c$ , monoclinic,  $a = 42.3319(19)$ ,  $b = 15.8991(7)$ ,  $c = 19.1013(9) \text{ \AA}$ ,  $\beta = 110.2890(10)^\circ$ ,  $V = 12058.3(9) \text{ \AA}^3$ ,  $Z = 8$ ,  $\rho_{\text{calcd}} = 1.183 \text{ mg m}^{-3}$ ,  $F(000) = 4648$ .<sup>[17]</sup>
- [11] W.-K. Kim, M. J. Fevola, L. M. Liable-Sands, A. L. Rheingold, K. H. Theopold, *Organometallics* **1998**, *17*, 4541–4543.
- [12] Crystallographic details for **4**: A crystal of approximate dimensions  $0.20 \times 0.20 \times 0.15 \text{ mm}^3$  was selected and mounted on a glass fiber. A total of 29941 reflections ( $-13 \leq h \leq 13$ ,  $-36 \leq k \leq 36$ ,  $-13 \leq l \leq 15$ ) was collected at  $T = 112(2) \text{ K}$  in the range of  $2.92$  to  $25.35^\circ$ , of which 5437 were observed ( $R_{\text{int}} = 0.0804$ );  $\text{MoK}\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). A direct-methods solution was calculated which provided most non-hydrogen atoms from the E-map. Full-matrix least-squares/difference Fourier cycles were performed which located the remaining non-hydrogen atoms. The residual peak and hole electron densities were  $1.104$  and  $-0.608 \text{ e \AA}^{-3}$ , respectively. The absorption coefficient was  $1.215 \text{ mm}^{-1}$ . The least-squares refinement converged normally with residuals of  $R(F) = 0.0387$ ,  $wR(F^2) = 0.0810$  and a GOF =  $0.894$  ( $I > 2\sigma(I)$ ).  $\text{C}_{34}\text{H}_{51}\text{IN}_2\text{V}$ , space group  $P2(1)/n$ , monoclinic,  $a = 10.4492(9)$ ,  $b = 27.924(2)$ ,  $c = 12.3261(10) \text{ \AA}$ ,  $\beta = 107.904(2)^\circ$ ,  $V = 3422.4(5) \text{ \AA}^3$ ,  $Z = 4$ ,  $\rho_{\text{calcd}} = 1.292 \text{ mg m}^{-3}$ .<sup>[17]</sup>
- [13] Crystallographic details for **5·2 Et<sub>2</sub>O**: A crystal of approximate dimensions  $0.25 \times 0.25 \times 0.08 \text{ mm}^3$  was selected and mounted on a glass fiber. A total of 43991 reflections ( $-15 \leq h \leq 15$ ,  $-19 \leq k \leq 19$ ,  $-27 \leq l \leq 34$ ) was collected at  $T = 111(2) \text{ K}$  in the range of  $2.07$  to  $27.53^\circ$ , of which 5447 were observed ( $R_{\text{int}} = 0.1258$ );  $\text{MoK}\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). A direct-methods solution was calculated which provided most non-hydrogen atoms from the E-map. Full-matrix least-squares/difference Fourier cycles were performed which located the remaining non-hydrogen atoms. A disordered diethyl ether solvent molecule was located in the cell at an inversion center. All non-hydrogen atoms were refined with anisotropic displacement parameters. The residual peak and hole electron densities were  $0.267$  and  $-0.374 \text{ e \AA}^{-3}$ , respectively. The absorption coefficient was  $0.260 \text{ mm}^{-1}$ . The least-squares refinement converged normally with residuals of  $R(F) = 0.0461$ ,  $wR(F^2) = 0.0839$  and a GOF =  $0.835$  ( $I > 2\sigma(I)$ ).  $\text{C}_{55}\text{H}_{66}\text{N}_2\text{O}_{0.5}\text{V}$ , space group  $P2(1)/n$ , monoclinic,  $a = 11.5823(11)$ ,  $b = 15.0376(15)$ ,  $c = 26.496(2) \text{ \AA}$ ,  $\beta = 101.522(3)^\circ$ ,  $V = 4521.8(7) \text{ \AA}^3$ ,  $Z = 4$ ,  $\rho_{\text{calcd}} = 1.212 \text{ mg m}^{-3}$ .<sup>[17]</sup>
- [14] Crystallographic details for **6**: A crystal of approximate dimensions  $0.30 \times 0.30 \times 0.24 \text{ mm}^3$  was selected and mounted on a glass fiber. A total of 21477 reflections ( $-12 \leq h \leq 12$ ,  $-13 \leq k \leq 15$ ,  $-23 \leq l \leq 23$ ) was collected at  $T = 130(2) \text{ K}$  in the range of  $2.03$  to  $30.02^\circ$ , of which 7739 were observed ( $R_{\text{int}} = 0.0320$ );  $\text{MoK}\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). A direct-methods solution was calculated which provided most non-hydrogen atoms from the E-map. Full-matrix least-squares/difference Fourier cycles were performed which located the remaining non-hydrogen atoms. All non-hydrogen atoms were refined with anisotropic displacement parameters. The residual peak and hole electron densities were  $1.105$  and  $-0.614 \text{ e \AA}^{-3}$ , respectively. The absorption coefficient was  $1.295 \text{ mm}^{-1}$ . The least-squares refinement converged normally with residuals of  $R(F) = 0.0341$ ,  $wR(F^2) = 0.0869$  and a GOF =  $1.020$  ( $I > 2\sigma(I)$ ).  $\text{C}_{34}\text{H}_{51}\text{IN}_2\text{V}$ , space group  $P\bar{1}$ , triclinic,  $a = 9.0935(4)$ ,  $b = 11.0457(5)$ ,  $c = 17.0793(8) \text{ \AA}$ ,  $\alpha = 102.4790(10)^\circ$ ,  $\beta = 92.1090(10)^\circ$ ,  $\gamma = 98.7240(10)^\circ$ ,  $V = 1651.30(13) \text{ \AA}^3$ ,  $Z = 2$ ,  $\rho_{\text{calcd}} = 1.339 \text{ mg m}^{-3}$ .<sup>[17]</sup>
- [15] F. Basuli, J. Tomaszewski, J. C. Huffman, D. J. Mindiola, *J. Am. Chem. Soc.* **2003**, *125*, 10170–10171.
- [16] For an example of ATRP see: J. F. Lutz, D. Neugebauer, K. Matyjaszewski, *J. Am. Chem. Soc.* **2003**, *125*, 6986–6993.
- [17] CCDC-222957 (**2**), CCDC-222958 (**3**), CCDC-222959 (**4**), CCDC-222960 (**5**), and CCDC-222961 (**6**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html) (or from the Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB21EZ, UK; fax: (+44) 1223-336-033; or deposit@ccdc.cam.ac.uk).