

## Synthetic Methods


## Gold(I)-Catalyzed 5-endo-dig Carbocyclization of Acetylenic Dicarboxyl Compounds\*\*

Steven T. Staben, Joshua J. Kennedy-Smith, and F. Dean Toste\*

The importance of cyclopentanoid natural products<sup>[1]</sup> continues to inspire the development of methods for the synthesis of 5-membered rings.<sup>[2]</sup> For example, the Conia-ene reaction provides an atom-economical synthesis of methylenecyclopentanes by the thermal cyclization of  $\varepsilon$ -acetylenic carbonyl compounds.<sup>[3]</sup> While the classic thermal reaction is limited to the exocyclic cyclization mode, group 6 metal complexes catalyze the formal 5-endo-dig addition of  $\beta$ -ketoesters<sup>[4]</sup> and silyl enol ethers<sup>[5]</sup> to alkynes. These reactions proceed via intermediate metal vinylidenes and therefore are limited to substrates containing a terminal acetylene moiety. The scope and utility of this reaction would be greatly increased if nonterminal alkynes could be employed as electrophiles.<sup>[6]</sup> However, while examples of the transition-metal-catalyzed 5-endo-dig addition of heteroatom nucleophiles to nonterminal alkynes are common,<sup>[7–9]</sup> this class of cyclization employing carbon nucleophiles is rare.<sup>[10]</sup>

We have recently demonstrated that cationic gold(I) complexes catalyze the Conia-ene reaction by a mechanism that appears to involve formation of a gold(I) alkyne complex.<sup>[11]</sup> Thus, we reasoned that an endocyclic Conia-ene reaction might be feasible with group 11 metal complexes as catalysts for alkyne activation.<sup>[8,9]</sup> Furthermore, while the gold(I)-catalyzed Conia-ene reaction is limited to terminal alkynes, we antici-

pated that a 5-endo-dig variant would allow for carbocyclization onto nonterminal alkynes. To this end, we surveyed several cationic group 11 metal triflates as catalysts for the cyclization of  $\beta$ -ketoesters **1** with nonterminal  $\alpha$ -3'-alkynyl substituents [Eq. (1)]. Reaction of **1** with Cu<sup>I</sup> and Ag<sup>I</sup> triflate

Table 1: Scope of Au<sup>I</sup>-catalyzed 5-endo-dig carbocyclization.<sup>[a]</sup>

Entry	Substrate	Time	Product	Yield
1		10 min		90%
2		5 min		96%
3		5 h		90%
4		5 min		83%
5		45 min		80%
6		1 h		88% <sup>[b]</sup>
7		1 h		83%
8		15 min		74%
9		10 min		94%
10		20 h		80% <sup>[c]</sup>
11		6 min		99%
12		6 h		99% <sup>[c]</sup>

[a] Reaction conditions: see Experimental Section. [b] Together with about 15% of a 1,3-diene isomer. [c] 2 mol%  $[(\text{PPh}_3)_3\text{Au}]\text{Cl}$ , 2 mol%  $\text{AgOTf}$ .

[\*] S. T. Staben, J. J. Kennedy-Smith, Prof. Dr. F. D. Toste  
Department of Chemistry  
University of California  
Berkeley, CA 94720 (USA)  
Fax: (+1) 510-643-9480  
E-mail: fdtoste@uclink.berkeley.edu

[\*\*] We gratefully acknowledge the University of California, Berkeley, Merck Research Laboratories, and Amgen Inc. for financial support. J.J.K.-S. thanks Eli Lilly & Co. for a graduate fellowship. The Center for New Directions in Organic Synthesis is supported by Bristol-Myers Squibb as a Sponsoring Member and Novartis Pharma as a Supporting Member.


Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

complexes did not produce the desired cyclopentene adduct **2**. On the other hand, triphenylphosphinegold(I) triflate rapidly (10 min) converted alkyne **1** into cyclopentene **2** in 93% yield. Notably, these reactions were run under “open-flask” conditions at room temperature without the necessity for dry solvents or inert atmosphere.

Under these experimentally simple conditions, a wide range of substrates underwent rapid 5-endo-dig cycloisomerization to give cyclopentene products (Table 1, entries 1–6). In all cases no *exo*-cyclization was observed, including a substrate (**5**) having the opportunity to undergo competitive 5-*exo*-dig cyclization onto a propargyl ester (entry 2). Variation at the ester (entries 1 and 2) and ketone (entry 3)


moieties is tolerated, although cyclization of aryl ketones requires increased reaction times. The reaction is also amenable to a wide range of alkynyl substituents including alkyl, aryl (entry 9), vinyl (entry 6), and proton (entry 7), although the latter appears to react more sluggishly. Importantly, the mild reaction conditions allow for the use of acid-labile groups such as *tert*-butyl ester [Eq. (1)], tetrahydropyranyl ether (entry 5), and tertiary propargyl ether (entry 11). This cycloisomerization provides an alternative synthesis of 1,3-dienes often prepared by enyne metathesis.<sup>[12]</sup> For example, 1,3-enyne **13** underwent rapid cyclization to give 1,3-diene **14** in good yield (entry 6).

Having established the feasibility of the endocyclic carbocyclization, we sought to apply this method to the synthesis of bicyclic structures by a cyclopentene annulation<sup>[13]</sup> onto  $\alpha,\beta'$ -unsaturated  $\beta$ -ketoesters. Thus, conjugate addition of allenyltriphenylstannane<sup>[14]</sup> to **27**, followed by gold(I)-catalyzed cyclization afforded cyclopentene **28** as a single diastereomer [Eq. (2)]. This cyclopentene annulation




can also be applied to the diastereoselective formation of 5,5- (Table 1, entries 7–9) and 7,5-fused (entry 10) bicyclic ring systems. Additionally, bicyclo[3.2.1]octane **24** is available in excellent yield from the 5-*endo-dig* cyclization of  $\beta$ -ketoester **23** (entry 11). Lewis-basic groups, such as a tertiary amine, are tolerated and thus the gold-catalyzed reaction allows for the synthesis of heterocyclic ring systems. For example, the benzo-fused pyrrolizidine core of the mitosanes (**26**)<sup>[15]</sup> is available in excellent yield from a gold(I)-catalyzed cyclization of 3-hydroxyindole **25** (entry 12).

We have found that  $\beta$ -diketones<sup>[6]</sup> are also viable nucleophiles under the optimized reaction conditions. For example, 1 mol % triphenylphosphinegold(I) triflate rapidly and efficiently catalyzes the conversion of 1,3-dione **29** into cyclopentene **30** [Eq. (3)].




Carbocyclization onto 1-halo-1-alkynes would provide a facile entry into cyclopentenyl halides, however, transition-metal-catalyzed addition reactions to alkynyl halides are exceptionally rare.<sup>[16]</sup> We were, therefore, very pleased to find that 1-iodoalkynes **31** and **33** underwent rapid cyclization to

give cyclopentenyl iodides **32** and **34** as single diastereomers in 93 and 76 % yield, respectively [Eq. (4)].



We propose that these reactions proceed by a mechanism involving nucleophilic addition of an enol to a gold(I) alkyne complex (Scheme 1). Based on this mechanistic hypothesis, one of the potential explanations for the lack of reactivity of nonterminal alkynes in the gold-catalyzed 5-*exo-dig* cyclization<sup>[11]</sup> is that placement of the catalyst near an alkyl-substituted carbon atom is sterically unfavorable. However, in the transition state for the endocyclic reaction the gold center is located near an alkyl-substituted carbon atom without inhibiting the cyclization. We propose that the 5-*exo-dig* Conia-ene reaction is limited to terminal alkynes because of the development of 1,3-allylic strain in the transition state. This strain is absent in the transition state for the gold(I)-catalyzed 5-*endo-dig* cyclization allowing for the participation of nonterminal alkynes.



**Scheme 1.** Proposed mechanism for the gold(I)-catalyzed 5-*endo-dig* carbocyclization.

In conclusion, we have developed a gold(I)-catalyzed 5-*endo-dig* carbocyclization of dicarbonyl compounds onto appended alkynes. The reaction is carried out under open-flask conditions and shows excellent tolerance for variation in the ketone, ester, and alkyne substituents. As such, it provides entry into a wide range of cyclopentanoid structures including those containing 1,3-diene, vinyl iodide, and heterocyclic moieties. These results further highlight the potential of gold(I) complexes<sup>[17]</sup> to serve as catalysts for the formation of carbon–carbon bonds by alkyne activation. Applications of this strategy, including an asymmetric variant, are underway in our laboratory and will be reported in due course.

## Experimental Section

**General synthetic procedure:** To a small screw-cap scintillation vial equipped with a magnetic stir bar and charged with a solution of the  $\alpha$ -3'-alkynyl-substituted  $\beta$ -dicarbonyl compound (~150 mg, 1 equiv)

in  $\text{CH}_2\text{Cl}_2$  (0.4 M) was added  $[\text{Au}(\text{PPh}_3)]\text{Cl}$  (1 mol %) followed by  $\text{AgOTf}$  (1 mol %). The cloudy white reaction mixture was then stirred at room temperature and monitored periodically by thin layer chromatography. Upon completion of the reaction, the mixture was loaded directly onto a silica gel column and chromatographed with the appropriate mixture of hexanes and ethyl acetate to give the cycloisomerized products.

Received: June 2, 2004

**Keywords:** C–C coupling · cyclization · cyclopentenones · dicarbonyl compounds · gold

- [1] For examples, see: a) R. Noyori, M. Suzuki, *Science* **1993**, 259, 44–45; b) A. Nangia, G. Prasuna, P. B. Rao, *Tetrahedron* **1997**, 53, 14507–14545; c) M. H. Beale, J. L. Ward, *Nat. Prod. Rep.* **1998**, 15, 533–548; d) M. Seepersaud, Y. Al-Abed, *Tetrahedron Lett.* **2000**, 41, 4291–4293.
- [2] For reviews, see: a) B. M. Trost, *Chem. Soc. Rev.* **1982**, 11, 141–170; b) M. Ramaiah, *Synthesis* **1984**, 529–570; c) T. Hudlicky, J. D. Price, *Chem. Rev.* **1989**, 89, 1467–1486; d) L. Ghosez, *Pure Appl. Chem.* **1996**, 68, 15–22; e) M. A. Tius, *Acc. Chem. Res.* **2003**, 36, 284–290; f) T. R. Rheault, M. P. Sibi, *Synthesis* **2003**, 803–819.
- [3] For a review, see: J. M. Conia, P. Le Perche, *Synthesis* **1975**, 1–19.
- [4] F. E. McDonald, T. C. Olson, *Tetrahedron Lett.* **1997**, 38, 7691–7692.
- [5] N. Iwasawa, T. Miura, K. Kiyota, H. Kusama, K. Lee, P. H. Lee, *Org. Lett.* **2002**, 4, 4463–4466.
- [6] A related  $\text{Pd}^{\text{II}}$ -catalyzed 6-endo-trig addition of 1,3-diones to alkenes has been reported: a) T. Pei, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2001**, 123, 11290–11291; b) H. Qian, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2003**, 125, 2056–2057.
- [7] For a review, see: F. Alonso, I. P. Beletskaya, M. Yus, *Chem. Rev.* **2004**, 104, 3079–3160.
- [8] Au-catalyzed 5-endo-dig cyclization of O nucleophiles: a) A. S. K. Hashmi, T. M. Frost, J. W. Bats, *J. Am. Chem. Soc.* **2000**, 122, 11553–11554; b) A. S. K. Hashmi, L. Schwarz, J.-H. Choi, T. M. Frost, *Angew. Chem.* **2000**, 112, 2382–2385; *Angew. Chem. Int. Ed.* **2000**, 39, 2285–2288; c) A. S. K. Hashmi, P. Sinha, *Adv. Synth. Catal.* **2004**, 346, 432–438; of N nucleophiles: d) Y. Fukuda, K. Utimoto, *Synthesis* **1991**, 975–978; e) A. Arcadi, S. D. Giuseppe, F. Marinelli, E. Rossi, *Adv. Synth. Catal.* **2001**, 343, 443–446; f) A. Arcadi, D. Bianchi, F. Marinelli, *Synthesis* **2004**, 610–618.
- [9] Cu-catalyzed 5-endo-dig cyclization with N nucleophiles: a) C. E. Castro, E. J. Gaughan, D. C. Owsley, *J. Org. Chem.* **1966**, 31, 4071–4078; b) J. Fujiwara, Y. Fukutani, H. Sano, K. Maruoka, H. Yamamoto, *J. Am. Chem. Soc.* **1983**, 105, 7177–7179; Cu-catalyzed 5-endo-dig cyclization with O nucleophiles: c) I. N. Houpi, W. B. Choi, P. J. Reider, A. Molina, H. Churchill, J. Lynch, R. P. Volante, *Tetrahedron Lett.* **1994**, 35, 9355–9358; d) C. C. Bates, P. Saejueng, J. M. Murphy, D. Venkataraman, *Org. Lett.* **2002**, 4, 4727–4729; Ag-catalyzed addition of O nucleophiles: e) J. A. Marshall, M. A. Wolf, E. M. Wallace, *J. Org. Chem.* **1997**, 62, 367–371.
- [10] a) 5-endo-dig Michael addition: J. F. Lavallée, G. Berthiaume, P. Deslongchamps, *Tetrahedron Lett.* **1986**, 27, 5455–5458; b)  $\text{Hf}^{\text{IV}}$ -catalyzed alkyne/allylsilane 5-endo-dig cycloisomerization: K.-I. Imamura, E. Yoshikawa, V. Gevorgyan, Y. Yamamoto, *J. Am. Chem. Soc.* **1998**, 120, 5339–5340; c)  $\text{Rh}^{\text{I}}$ -catalyzed 5-endo-dig hydroacylation: K. Tanaka, G. C. Fu, *J. Am. Chem. Soc.* **2001**, 123, 11492–11493; d) radical 5-endo-dig carbocyclization: S. Amrein, A. Studer, *Chem. Commun.* **2002**, 1592–1593; e) Co-catalyzed cycloisomerization of 1,6-enynes: A. Ajamian, J. L. Gleason, *Org. Lett.* **2003**, 5, 2409–2411.
- [11] J. J. Kennedy-Smith, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2004**, 126, 4526–4528.
- [12] For a review of enyne metathesis, see: S. T. Diver, A. Giessert, *Chem. Rev.* **2004**, 104, 1317–1382.
- [13] For a related cyclopentene annulation onto  $\alpha,\beta$ -unsaturated ketones, see: a) R. L. Danheiser, D. J. Carini, A. Basak, *J. Am. Chem. Soc.* **1981**, 103, 1604–1606; b) R. L. Danheiser, D. J. Carini, D. M. Fink, A. Basak, *Tetrahedron* **1983**, 39, 935–947.
- [14] J. Haruta, K. Nishi, S. Matsuda, S. Akai, Y. Tamaura, Y. Kita, *J. Org. Chem.* **1990**, 55, 4853–4859.
- [15] For examples of transition-metal-mediated approaches to this ring system, see: a) J. Lee, J. D. Ha, J. K. Cha, *J. Am. Chem. Soc.* **1997**, 119, 8127–8128; b) H. Kusama, J. Takaya, N. Iwasawa, *J. Am. Chem. Soc.* **2002**, 124, 11592–11593.
- [16] For a 5-*exo-dig* acetoxymercuration of a 1-iodo-1-alkyne, see: a) G. A. Krafft, J. A. Katzenellenbogen, *J. Am. Chem. Soc.* **1981**, 103, 5459–5466; see also the W-mediated reaction of 1-iodo-1-alkynes involving a 1,2-iodide shift: b) T. Miura, N. Iwasawa, *J. Am. Chem. Soc.* **2002**, 124, 518–519; c) N. Iwasawa, T. Miura, K. Kiyota, H. Kusama, K. Lee, P. H. Lee, *Org. Lett.* **2002**, 4, 4463–4466.
- [17] For reviews of Au-catalyzed reactions, see: a) G. Dyker, *Angew. Chem.* **2000**, 112, 4407–4409; *Angew. Chem. Int. Ed.* **2000**, 39, 4237–4283; b) A. S. K. Hashmi, *Gold Bull.* **2003**, 36, 3–9; see also: C. Nieto-Oberhuber, M. P. Muñoz, E. Buñuel, C. Nevado, D. J. Cárdenas, A. M. Echavarren, *Angew. Chem.* **2004**, 116, 2456–2460; *Angew. Chem. Int. Ed.* **2004**, 43, 2402–2406.