


A Nickel(II) Gold(I) D-Penicillamate Coordination System with Multiple Switching in Color, Magnetism, and Chirality**

Mai Taguchi, Asako Igashira-Kamiyama, Takashi Kajiware, and Takumi Konno*

The design and creation of molecular systems that show chemical and physical changes in response to external stimuli is a topic of current interest because of their potential applications as sensing and switching devices.^[1] A common approach to create such systems is to combine multifunctional organic or inorganic ligands or both with transition-metal ions that adopt flexible geometrical and electronic structures.^[2] The Ni²⁺ ion is promising as a metal component of switching systems because it can adopt several coordination geometries such as octahedral and square-planar, which are interconvertible with concomitant changes in spin state and color. In fact, there are a number of reports on the chromotropic behavior of Ni^{II} complexes based on geometrical changes.^[3] However, most systems contain a single Ni^{II} center; therefore, the level of switching in such systems is limited. Herein we report a unique dinickel(II) coordination system derived from the combination of Ni^{II} and [Au(d-Hpen-S)₂][−] ions (d-Hpen = d-penicillamine). The anion serves as a chiral multidentate metalloligand by exploiting free amine and carboxylate groups in addition to coordinated thiolate groups.^[4] This system involved four different S-bridged Ni^{II}Au^I structures that are interconvertible in response to several external factors; changes in structure are accompanied by changes in color, magnetism, and chirality (Scheme 1). To our knowledge, such a multiple reversible switching system is unprecedented.

The reaction of an aqueous solution of NH₄[Au(d-Hpen-S)₂], neutralized by K₂CO₃, with Ni(NO₃)₂ in a 1:1 ratio gave a red solution, from which red crystals of K₂[1] were isolated. X-ray fluorescence spectroscopy suggested that K₂[1] contains Ni and Au atoms in a 1:1 ratio and its elemental-analysis data were in agreement with a 1:1 adduct of [Au(d-pen)₂]^{3−} and Ni²⁺ ions. The structure of K₂[1] was determined by



Scheme 1. Four S-bridged Ni^{II}Au^I structures constructed from [Au(d-Hpen-S)₂][−] and Ni²⁺ ions, and their interconversion as a consequence of changes in the solution pH (red arrows) and the Ni/Au ratio (blue arrows). The coordination environments and electronic configurations of the Ni^{II} centers are highlighted. Photographs of the isolated salts of these ionic structures show their different colors.

single-crystal X-ray analysis.^[5] As shown in Figure 1 a, the [1]^{2−} ion has an expected S-bridged Ni^{II}₂Au^I₂ structure in [Au₂{Ni(d-pen-N,S)₂}₂]^{2−}, in which two square-planar [Ni(d-pen-N,S)₂]^{2−} units bearing two bidentate N,S-d-pen moieties are linked by two linearly coordinated Au^I centers in a twisted chairlike form.^[6] The two bridging S atoms in each square-planar [Ni(d-pen-N,S)₂]^{2−} unit commonly adopt the opposite (*R* and *S*) configurations. The Vis/near-IR spectrum of K₂[1] in the solid state is dominated by a visible-light band at 488 nm, which is assigned to the ³A₂ ← ³T₁ transition for Ni^{II} centers with square-planar environments.^[7]

The reaction of NH₄[Au(d-Hpen-S)₂] with Ni(NO₃)₂ in a 3:2 ratio under similar conditions yielded a blue solution from which purple-blue crystals of K₅[2] were isolated. X-ray fluorescence and elemental-analysis data imply that this compound contains [Au(d-pen)₂]^{3−} and Ni²⁺ ions in a 3:2 ratio. While the overall IR spectrum of K₅[2] is very similar to that of K₂[1], its Vis/near-IR spectrum is different.^[8] K₅[2] gives a broad near-IR band (902 nm) and a visible-light band (577 nm) in the solid state, which are characteristic of octahedral Ni^{II} complexes as a result of ³T_{2g} ← ³A_{2g} and ³T_{1g} ← ³A_{2g} transitions.^[7] Based on these results, K₅[2] is assumed to contain octahedrally coordinated Ni^{II} centers surrounded by three bidentate d-pen-N,S moieties. Indeed, X-ray analysis demonstrated that [2]^{5−} has an S-bridged Ni^{II}₂Au^I₃ structure, in which two octahedral [Ni(d-pen-N,S)₃]^{4−} units are linked by three Au^I atoms with linear

[*] M. Taguchi, Dr. A. Igashira-Kamiyama, Prof. T. Konno
Department of Chemistry, Graduate School of Science
Osaka University
Toyonaka, Osaka 560-0043 (Japan)
Fax: (+81) 6-6850-5765
E-mail: konno@ch.wani.osaka-u.ac.jp

Dr. T. Kajiware
Department of Chemistry, Graduate School of Science
Tohoku University
Aoba, Sendai 980-8578 (Japan)

[**] This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (nos. 17036035 and 18033030) from the Ministry of Education, Culture, Sports, Science and Technology of Japan. The authors thank Dr. M. Nakano for the help with magnetic analyses.

Supporting information (experimental details and spectroscopic data) for this article is available on the WWW under <http://www.angewandte.org> or from the author.




Figure 1. Perspective views of a) $[1]^{2-}$, b) $[2]^{5-}$, c) $[3]^{2-}$, and d) $[4]^+$. Ni green, Au violet, C gray, H white (C-bound H omitted), N blue, O red, S yellow.

coordination environments (Figure 1b).^[5] As there are two kinds of chiralities, Δ or Λ , for the two octahedrally coordinated Ni^{II} atoms, and R or S configurations for the six bridging S atoms, a total of 21 stereoisomers are possible. However, the $[2]^{5-}$ ion gave only the $(\Lambda)_2(R)_6$ isomer that forms a triple helix from S–Au–S linkages.

Prompted by the formation of the different $\text{Ni}^{\text{II}}\text{Au}^{\text{I}}$ structures when changing the molar ratio from 1:1 to 3:2, we investigated the reaction of $\text{NH}_4[\text{Au}(\text{D-Hpen-S})_2]$ and $\text{Ni}(\text{NO}_3)_2$ with a molar ratio of 2:3 under similar conditions. As with the 1:1 reaction, the resulting solution was red and light blue crystals of $[\text{Ni}(\text{H}_2\text{O})_6][3]$, which is insoluble in

water, were grown from the red solution with a weakening of its color. X-ray fluorescence and elemental-analysis data imply that $[\text{Ni}(\text{H}_2\text{O})_6][3]$ contains $[\text{Au}(\text{D-pen})_2]^{3-}$ and Ni^{2+} ions in a 2:3 ratio, which is consistent with the reaction stoichiometry. The presence of octahedral Ni^{II} centers in $[\text{Ni}(\text{H}_2\text{O})_6][3]$ is indicated by the appearance of a broad near-IR band (1094 nm) and a visible-light band (638 nm) in the solid-state spectrum. The crystal structure of $[\text{Ni}(\text{H}_2\text{O})_6][3]$, determined by X-ray analysis,^[5] revealed the presence of a $\text{Ni}^{\text{II}}_2\text{Au}^{\text{I}}_2$ tetranuclear anion, $[3]^{2-}$, and an $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ ion (Figure 1c). In the $[3]^{2-}$ ion, two $[\text{Ni}(\text{D-pen-N,O,S})_2]^{2-}$ units are linked by two linearly coordinated Au^{I} centers to form an S-bridged $\text{Ni}^{\text{II}}_2\text{Au}^{\text{I}}_2$ structure. Each Ni^{II} center in the $[3]^{2-}$ ion has an octahedral coordination geometry, as with the $[2]^{5-}$ ion, but each center is surrounded by two tridentate D-pen-N,O,S moieties such that two N,S-chelate rings selectively form a Λ configurational skew pair with four bridging S atoms being fixed to the R configuration. It should be noted that in the crystal, each $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ ion is spanned by two neighboring cations by strong O–H...O hydrogen bonds ($\text{O}\cdots\text{O}$ 2.61(3) Å–2.81(3) Å), thus creating a rare example of a 1D chain composed of hexa-aqua metal species.^[8]

To investigate the influence of solution pH on the formation of $\text{Ni}^{\text{II}}\text{Au}^{\text{I}}$ structures, an aqueous solution of $\text{NH}_4[\text{Au}(\text{D-Hpen-S})_2]$ ($\text{pH} \approx 3.5$) was treated with $\text{Ni}(\text{NO}_3)_2$ in a 1:1 or 3:2 ratio without the addition of K_2CO_3 . Interestingly, a green reaction solution resulted, from which yellow-green crystals of $[4]\text{NO}_3$ were isolated, which are poorly soluble in water. Although the solid-state spectrum of this compound, which is dominated by a broad near-IR band (1102 nm) and a visible-light band (672 nm), is similar to that of the $[2]^{5-}$ ion, each band is considerably red-shifted, thus indicating a weaker ligand field around the octahedral Ni^{II} centers. X-ray analysis for $[4]\text{Br}$, prepared by using NiBr_2 instead of $\text{Ni}(\text{NO}_3)_2$, demonstrated that the $[4]^+$ ion has an S-bridged $\text{Ni}^{\text{II}}_2\text{Au}^{\text{I}}_3$ structure in $[\text{Au}_3\{\text{Ni}(\text{D-Hpen-O,S})_3\}_2]^+$ consisting of two octahedrally-coordinated $[\text{Ni}(\text{D-Hpen-O,S})_3]^-$ units linked by three Au^{I} centers with linear coordination environments (Figure 1d).^[5] This S-bridged structure in the $[4]^+$ ion resembles that in the $[2]^{5-}$ ion except that each D-Hpen moiety in the former coordinates to a Ni^{II} center through carboxylate and thiolate groups and a protonated NH_3^+ group does not participate in the coordination. In the $[4]^+$ ion, the two octahedral Ni^{II} units selectively take the Λ configuration, with six bridging S atoms being fixed to the R configuration as in the $[2]^{5-}$ ion. However, the three S–Au–S linkages in the $[4]^+$ ion are almost parallel to each other because of the steric demand of the O,S-chelating mode.

As expected, magnetic measurements indicate that $\text{K}_5[2]$, $[\text{Ni}(\text{H}_2\text{O})_6][3]$, and $[4]\text{Br}$ with octahedral Ni^{II} centers are all paramagnetic, while $\text{K}_2[1]$ with square-planar Ni^{II} centers is diamagnetic. From the temperature dependence of the magnetic susceptibility (2–300 K), a very weak antiferromagnetic interaction between two Ni^{II} centers was revealed for the $[2]^{5-}$ ($J = -4.3(1) \text{ cm}^{-1}$ and $g = 2.04(1)$) and $[3]^{2-}$ ions ($J = -1.1(1) \text{ cm}^{-1}$ and $g = 2.16(1)$), which is compatible with the large Ni...Ni separation (6.808(2) Å in $[2]^{5-}$ and 7.351(3) Å in $[3]^{2-}$).^[8] However, for the $[4]^+$ ion, an appreciable antiferromagnetic interaction was recognized between the two Ni^{II}

centers ($J = -11.6(1) \text{ cm}^{-1}$ and $g = 2.19(1)$) despite the Ni...Ni separation ($7.025(4) \text{ \AA}$) being comparable with those in the $[2]^{5-}$ and $[3]^{2-}$ ions.^[8] The presence of a relatively large M...M magnetic interaction through closed-shell $d^{10} \text{ Au}^I$ centers bridged by thiolate groups is noteworthy and is related to the parallel arrangement of S–Au–S linkages in the $[4]^+$ ion, which is distinct from the skewed arrangement in the $[2]^{5-}$ and $[3]^{2-}$ ions.

The most striking feature of this $\text{Ni}^{II}\text{Au}^I$ coordination system is the facile interconversion between the four S-bridged structures, accompanied by color change (Scheme 1).^[8,9] When 0.5 molar equivalents of $\text{Ni}(\text{NO}_3)_2$ were added to a blue aqueous solution of $\text{K}_5[2]$, the solution immediately turned red, and the absorption spectrum coincided with that of the $[1]^{2-}$ ion; further addition of $\text{Ni}(\text{NO}_3)_2$ led to the precipitation of a light blue powder, $[\text{Ni}(\text{H}_2\text{O})_6][3]$. In reverse, treatment of an aqueous suspension of $[\text{Ni}(\text{H}_2\text{O})_6][3]$ with 0.5 molar equivalents of $[\text{Au}(\text{p-pen-S})_2]^{3-}$ produced a red solution ($[1]^{2-}$), which turned blue ($[2]^{5-}$) on further addition of $[\text{Au}(\text{p-pen-S})_2]^{3-}$ ions. Furthermore, the addition of 0.1 M of aqueous HNO_3 to a blue aqueous solution of $\text{K}_5[2]$ ($\text{pH} \approx 8.5$) induced a change to red ($[1]^{2-}$) in a pH range of about 7 to 5, and this red solution turned green ($[4]^+$) on lowering the pH to about 3.5 by further addition of 0.1 M aqueous HNO_3 . The green solution of $[4]^+$ ions reverted back to a blue solution of $[2]^{5-}$ ions by way of a red solution on raising the pH to about 8.5 with aqueous NH_3 . Notably, the $[1]^{2-}$ and $[2]^{5-}$ ions are also interconvertible in response to solvent and temperature, thus demonstrating solvatochromism and thermochromism.^[10]

In summary, we have shown that four different, chiral-selective polynuclear structures are constructed from only Ni^{II} and $[\text{Au}(\text{p-pen-S})_2]^{3-}$ ions because of the flexible geometries of the Ni^{II} center and the versatile coordination modes of D-penicillamine bound to a Au^I center (N,S-bidentate, O,S-bidentate, and N,O,S-tridentate). Complexes $[2]^{5-}$, $[1]^{2-}$, and $[3]^{2-}$ are interconvertible in response to the $\text{Ni}^{II}/[\text{Au}(\text{p-pen-S})_2]^{3-}$ ratio, whereas $[2]^{5-}$, $[1]^{2-}$, and $[4]^+$ are interconvertible in response to solution pH. Note that $[1]^{2-}$ has Ni^{II} centers with a square-planar, achiral coordination environment and with a low-spin d^8 configuration, while $[2]^{5-}$, $[3]^{2-}$, and $[4]^+$ have Ni^{II} centers with an octahedral, chiral coordination environment and with a high-spin configuration. Thus, $[1]^{2-}$ is triply reversible with $[2]^{5-}$, $[3]^{2-}$, and $[4]^+$; the interconversion is accompanied not only by the readily detectable color change, but also by the drastic switching in magnetism and chirality. The creation of other dynamic switching systems through the use of multifunctional chiral metalloligands, such as $[\text{Au}(\text{p-pen-S})_2]^{3-}$ ions, in combination with two or more transition-metal ions with manifold geometrical and electronic properties is promising.

Received: November 21, 2006

Published online: February 19, 2007

Keywords: amino acids · coordination modes · ligand effects · molecular devices · sensors

- [1] a) *Molecular Switches* (Ed.: B. L. Feringa), Wiley-VCH, Weinheim, Germany, **2001**; b) G. W. Gokel, W. M. Leevy, M. E. Weber, *Chem. Rev.* **2004**, *104*, 2723–2750; c) K. Kinbara, T. Aida, *Chem. Rev.* **2005**, *105*, 1377–1400.
- [2] a) P. Gütllich, Y. Garcia, T. Woike, *Coord. Chem. Rev.* **2001**, *219–221*, 839–879; b) A. Dei, D. Gatteschi, C. Sangregorio, L. Sorace, *Acc. Chem. Res.* **2004**, *37*, 827–835; c) S. Kume, M. Murata, T. Ozeki, H. Nishihara, *J. Am. Chem. Soc.* **2005**, *127*, 490–491; d) N. Matsumoto, Y. Motoda, T. Matsuo, T. Nakashima, N. Re, F. Dahan, J.-P. Tuchagues, *Inorg. Chem.* **1999**, *38*, 1165–1173.
- [3] a) P. S. Pallavicini, A. Perotti, A. Poggi, B. Seghi, L. Fabbri, *J. Am. Chem. Soc.* **1987**, *109*, 5139–5144; b) W. Linert, Y. Fukuda, A. Camard, *Coord. Chem. Rev.* **2001**, *218*, 113–152; c) K. Miyamoto, M. Sakamoto, C. Tanaka, E. Horn, Y. Fukuda, *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1061–1071; d) A. Tamayo, J. Casabó, L. Escriche, C. Lodeiro, B. Covelo, C. D. Brondino, R. Kivekäs, R. Sillampää, *Inorg. Chem.* **2006**, *45*, 1140–1149.
- [4] A. Toyota, T. Yamaguchi, A. Igashira-Kamiyama, T. Kawamoto, T. Konno, *Angew. Chem.* **2005**, *117*, 1112–1116; *Angew. Chem. Int. Ed.* **2005**, *44*, 1088–1092.
- [5] Crystal data for $\text{K}_5[1] \cdot 6\text{H}_2\text{O}$: Bruker SMART CCD diffractometer, $T = 210 \text{ K}$, MoK_α radiation ($\lambda = 0.71073 \text{ \AA}$). The structure was solved by direct methods with SIR97 and refined by full-matrix least-squares techniques on F^2 by using SHELXL-97.^[11] $\text{C}_{80}\text{H}_{184}\text{Au}_8\text{K}_8\text{N}_{16}\text{Ni}_8\text{O}_{52}\text{S}_{16}$, $M_r = 5073.60$, crystal size $0.26 \times 0.07 \times 0.07 \text{ mm}^3$, trigonal, space group $R\bar{3}$, $a = 36.132(8) \text{ \AA}$, $c = 37.686(11) \text{ \AA}$, $V = 42607(19) \text{ \AA}^3$, $Z = 9$, $\rho_{\text{calcd}} = 1.780 \text{ g cm}^{-3}$, $\mu = 7.362 \text{ mm}^{-1}$, ω – 2θ scan mode, $2\theta_{\text{max}} = 55.0^\circ$, 102 126 reflections collected, 42 850 independent reflections, 25 612 observed reflections ($I > 2\sigma(I)$), 1568 parameters, semiempirical absorption corrections with SADABS, max./min. transmission 0.563/0.445, $R1$ ($I > 2\sigma(I)$) = 0.0458, $wR2$ (all data) = 0.0957, Flack parameter $-0.023(4)$, residual electron density $2.380/-0.894 \text{ e \AA}^{-3}$. Crystal data for $\text{K}_5[2] \cdot 23\text{H}_2\text{O} \cdot \text{KNO}_3$: Bruker SMART CCD diffractometer, $T = 220 \text{ K}$, MoK_α radiation ($\lambda = 0.71073 \text{ \AA}$). The structure was solved by direct methods with SIR97 and refined by full-matrix least-squares techniques on F^2 by using SHELXL-97.^[11] $\text{C}_{30}\text{H}_{100}\text{Au}_3\text{K}_6\text{N}_7\text{Ni}_2\text{O}_{38}\text{S}_6$, $M_r = 2302.45$, crystal size $0.25 \times 0.20 \times 0.10 \text{ mm}^3$, trigonal, space group $P3_2$, $a = 11.6927(13) \text{ \AA}$, $c = 48.269(89) \text{ \AA}$, $V = 5715.1(13) \text{ \AA}^3$, $Z = 3$, $\rho_{\text{calcd}} = 2.007 \text{ g cm}^{-3}$, $\mu = 6.819 \text{ mm}^{-1}$, ω – 2θ scan mode, $2\theta_{\text{max}} = 50.0^\circ$, 25 529 reflections collected, 11 364 independent reflections, 10 445 observed reflections ($I > 2\sigma(I)$), 812 parameters, semiempirical absorption corrections with SADABS, max./min. transmission 0.562/0.406, $R1$ ($I > 2\sigma(I)$) = 0.0338, $wR2$ (all data) = 0.0615, Flack parameter $-0.019(5)$, residual electron density $1.460/-1.075 \text{ e \AA}^{-3}$. Crystal data for $[\text{Ni}(\text{H}_2\text{O})_6][3] \cdot 3.5\text{H}_2\text{O}$: Bruker SMART CCD diffractometer, $T = 200 \text{ K}$, MoK_α radiation ($\lambda = 0.71073 \text{ \AA}$). The structure was solved by the Patterson method with DIRDIF99^[12] (PATY) and refined by full-matrix least-squares techniques on F^2 by using SHELXL-97.^[11] $\text{C}_{20}\text{H}_{55}\text{Au}_2\text{N}_4\text{Ni}_3\text{O}_{17.5}\text{S}_4$, $M_r = 1329.98$, crystal size $0.39 \times 0.12 \times 0.05 \text{ mm}^3$, monoclinic, space group $C2$, $a = 22.307(6) \text{ \AA}$, $b = 5.5472(14) \text{ \AA}$, $c = 17.234(4) \text{ \AA}$, $\beta = 133.135(5)^\circ$, $V = 1961.1(9) \text{ \AA}^3$, $Z = 2$, $\rho_{\text{calcd}} = 2.252 \text{ g cm}^{-3}$, $\mu = 9.153 \text{ mm}^{-1}$, ω – 2θ scan mode, $2\theta_{\text{max}} = 55.0^\circ$, 6803 reflections collected, 3202 independent reflections, 2936 observed reflections ($I > 2\sigma(I)$), 227 parameters, absorption corrections (numerical integration), max./min. transmission 0.629/0.191, $R1$ ($I > 2\sigma(I)$) = 0.0361, $wR2$ (all data) = 0.0897, Flack parameter $-0.021(10)$, residual electron density $2.224/-1.711 \text{ e \AA}^{-3}$. Crystal structure analysis for $[4]\text{Br} \cdot 9\text{H}_2\text{O}$: Bruker SMART CCD diffractometer, $T = 200 \text{ K}$, MoK_α radiation ($\lambda = 0.71073 \text{ \AA}$). The structure was solved by the direct method with SIR97 and refined by full-matrix least-squares techniques on F^2 by using SHELXL-97.^[11] $\text{C}_{30}\text{H}_{78}\text{Au}_3\text{BrN}_6\text{Ni}_2\text{O}_{21}\text{S}_6$, $M_r = 1839.57$, crystal

- size $0.40 \times 0.08 \times 0.08 \text{ mm}^3$, hexagonal, space group $P6_522$, $a = 11.8102(6)$, $c = 76.306(6) \text{ \AA}$, $V = 9217.3(9) \text{ \AA}^3$, $Z = 6$, $\rho_{\text{calcd}} = 1.988 \text{ g cm}^{-3}$, $\mu = 8.659 \text{ mm}^{-1}$, ω - 2θ scan mode, $2\theta_{\text{max}} = 55.0^\circ$, 100847 reflections collected, 7074 independent reflections, 6295 observed reflections ($I > 2\sigma(I)$), 317 parameters, semiempirical absorption corrections with SADABS, max./min. transmission 0.603/0.435, $R1$ ($I > 2\sigma(I)$) = 0.0722, $wR2$ (all data) = 0.1571, Flack parameter 0.067(18), residual electron density 1.207/−2.983 $\text{e \AA}^{-3}$. CCDC-627744–627747 (**1–4**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [6] The structure of $\text{K}_2[\mathbf{1}]$ is very similar to that of the previously reported $\text{Na}_2[\text{Au}_2\{\text{Ni}(\text{D-pen-}N,S)_2\}_2]$, which was isolated from a mixture of D-H₂pen, $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, and $[\text{Au}(\text{tu})_2\text{Cl}]$ (tu = thio-urea) in aqueous sodium acetate buffer. P. J. M. L. Birker, G. C. Verschoor, *Inorg. Chem.* **1982**, 21, 990–995.
- [7] F. Meyer, H. Kozłowski, *Comprehensive Coordination Chemistry II* (Eds.: J. A. McCleverty, T. J. Meyer), Elsevier, Amsterdam (The Netherlands), **2004**, chap. 6.3.
- [8] See the Supporting Information.
- [9] Compounds $\text{K}_2[\mathbf{1}]$, $\text{K}_5[\mathbf{2}]$, $[\text{Ni}(\text{H}_2\text{O})_6][\mathbf{3}]$, and $[\mathbf{4}]\text{X}$ ($\text{X} = \text{NO}_3, \text{Br}$) are all optically active, and their absorption and CD spectra in water, except those of insoluble $[\text{Ni}(\text{H}_2\text{O})_6][\mathbf{3}]$, are essentially the same as those in the solid state, which is indicative of the retention of their chiral structures in water.
- [10] Dissolution of $\text{K}_5[\mathbf{2}]$ in methanol gave a red solution characteristic of the presence of $[\mathbf{1}]^{2-}$ ions, although its aqueous solution is blue. In addition, a blue aqueous solution of $\text{K}_5[\mathbf{2}]$ turned red when heated to 60°C and reverted back to blue on cooling to room temperature.
- [11] G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures, University of Göttingen (Germany), **1998**.
- [12] P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, R. de Gelder, R. Israel, J. M. M. Smits, The DIRDIF99 Program System, Technical Report of the Crystallography Laboratory, University of Nijmegen (The Netherlands), **1999**.