


# Highly Enantioselective and Diastereoselective Cycloaddition of Cyclopropanes with Nitrones and Its Application in the Kinetic Resolution of 2-Substituted Cyclopropane-1,1-dicarboxylates\*\*

Yan-Biao Kang, Xiu-Li Sun, and Yong Tang\*

Trisoxazolines<sup>[1]</sup> have been applied widely in asymmetric catalysis<sup>[2]</sup> and molecular recognition.<sup>[3]</sup> In our efforts to develop superior catalysts that are cheap, readily accessible, air stable, and water tolerant, we designed and synthesized the pseudo- $C_3$ -symmetric trisoxazoline **1b** (Scheme 1) by the



**Scheme 1.** Trisoxazoline (tox) ligands. Py = pyridyl.

sidearm approach, and found that **1b**/ $\text{Cu}^{\text{II}}$  promoted smoothly the highly enantioselective Friedel–Crafts reaction of indoles with arylidene malonates<sup>[4]</sup> and the Kinugasa reaction.<sup>[5]</sup> We also described tox/ $\text{Co}^{\text{II}}$ -catalyzed [3+2] cycloaddition reactions between nitrones and alkylidene malonates in which the *endo/exo* selectivity could be controlled well by the reaction temperature. Thus, both *cis* and *trans* isoxazolidines could be prepared enantioselectively.<sup>[6]</sup> Herein, we report that the system tox/ $\text{Ni}^{\text{II}}$  catalyzes the [3+3] cycloaddition of racemic 2-substituted cyclopropane-1,1-dicarboxylates with nitrones to provide ready access to optically active tetrahydro-1,2-oxazine derivatives with high diastereoselectivity and enantioselectivity. Furthermore, this reaction can be employed for the kinetic resolution<sup>[7]</sup> of 2-substituted cyclopropane-1,1-dicarboxylates to furnish these compounds in optically active form with excellent *ee* values.

Tetrahydro-1,2-oxazine derivatives occur frequently in biologically active compounds<sup>[8]</sup> and are valuable synthetic intermediates.<sup>[9]</sup> Among the methods developed for the

preparation of such compounds,<sup>[10]</sup> the [3+3] cycloaddition of donor–acceptor cyclopropanes with nitrones, pioneered by Kerr and co-workers,<sup>[11]</sup> is a particularly elegant approach. They found that nitrones **3** react smoothly with cyclopropanes **2** in the presence of  $\text{Yb}(\text{OTf})_3 \cdot x \text{H}_2\text{O}$  to give the *cis* isomers **4** diastereospecifically (see Table 1). Recently, Sibi et al. developed a highly efficient asymmetric version of this reaction.<sup>[12]</sup> They reported that nitrones react with cyclopropane-1,1-dicarboxylates to afford tetrahydro-1,2-oxazine derivatives with high enantioselectivity in the presence of dbfox/ $\text{Ni}^{\text{II}}$  (10 mol %; dbfox = 4,6-dibenzofurandiyl-2,2'-bisoxazoline). When 2-substituted cyclopropane-1,1-dicarboxylates were used as substrates, and the quantity of the catalyst was increased to 30 mol %, the reactions proceeded smoothly to give the products with high *ee* values but with low diastereoselectivities (the *cis/trans* ratio ranged from 1.0:1.4 to 1.0:0.8).


Gratifyingly, we found that diethyl 2-phenylcyclopropane-1,1-dicarboxylate (**2a**) reacted smoothly with nitrone **3a** to furnish the cycloaddition adduct in 99 % yield with excellent diastereoselectivity (d.r. > 99:1) in the presence of a catalytic amount of **1b**/ $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ . Further studies showed that ligands derived from *i*Pr-box (box = bisoxazoline) influenced strongly both the enantioselectivity and diastereoselectivity of the reaction. The bisoxazoline **1a** without a pendant group gave the desired product with good diastereoselectivity (d.r. 97:3) but only 20 % *ee* (Table 1, entry 1). However, when a pyrid-2-yl or oxazolinyl group was introduced as a pendant group, the enantioselectivity was improved to a moderate level, and the high diastereoselectivity was maintained (Table 1, entries 2, 3, and 5). Both excellent enantioselectivity and high diastereoselectivity were observed when the substituent on the nitrogen atom of the nitrone was changed from a phenyl group to a methyl group (Table 1, entry 6). In this case, **4b** was obtained with 95 % *ee* and d.r. 11:1 when a catalytic amount of **1e**/ $\text{Ni}(\text{ClO}_4)_2$  was used. Thus, both high enantioselectivity and high diastereoselectivity could be attained by using **1e**/ $\text{Ni}(\text{ClO}_4)_2$  as the catalyst. The ester groups of the 2-phenyl cyclopropane-1,1-dicarboxylate also influenced the enantioselectivity. For example, when **2b** was used instead of **2a**, the *ee* value of the product decreased from 95 to 90 % (Table 1, entry 7).

We studied the scope of this reaction in terms of the substrate under the optimal conditions by investigating a variety of nitrones and cyclopropanes with different structures. As shown in Table 2, the ester groups of cyclopropanes **2** slightly influenced the enantioselectivity of the reaction. Benzyl and ethyl diesters reacted with higher enantioselectivity than the corresponding methyl diester (Table 2,

[\*] Y.-B. Kang, Dr. X.-L. Sun, Prof. Dr. Y. Tang  
The State Key Laboratory of Organometallic Chemistry  
Shanghai Institute of Organic Chemistry  
Chinese Academy of Sciences  
354 Fenglin Lu, Shanghai, 200032 (P.R. China)  
Fax: (+86) 215-492-5078  
E-mail: tangy@mail.sioc.ac.cn


[\*\*] We are grateful for the financial support of the Natural Sciences Foundation of China, the Major State Basic Research Development Program (Grant No. 2006CB806105), the Chinese Academy of Sciences, and The Science and Technology Commission of Shanghai Municipality.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

**Table 1:** Effect of the ligand on the cycloaddition.<sup>[a]</sup>


Entry	R <sup>1</sup>	R <sup>2</sup>	1	t [days]	4	Yield [%] <sup>[b]</sup>	d.r. <sup>[c]</sup>	ee [%] <sup>[d]</sup>
1	Et	Ph	1a	4	4a	96	97:3	20
2	Et	Ph	1b	4	4a	99	99:1	55
3	Et	Ph	1c	4	4a	91	99:1	66
4	Et	Ph	1d	4	4a	99	94:6	13
5	Et	Ph	1e	4	4a	99	99:1	57
6	Et	Me	1e	4	4b	88 <sup>[e]</sup>	11:1	95
7	Me	Me	1e	3	4c	82 <sup>[e]</sup>	13:1	90
8	Me	Me	1c	3	4c	77 <sup>[e]</sup>	6:1	88

[a] Reaction conditions: Ni(ClO<sub>4</sub>)<sub>2</sub> (0.040 mmol), 1 (0.044 mmol), 2 (0.44 mmol), 3 (0.20 mmol), 4-Å molecular sieves; DME = 1,2-dimethoxyethane. [b] Yield of the isolated product. [c] Ratio of isomers (*cis/trans*) as determined by <sup>1</sup>H NMR spectroscopy. [d] Determined by HPLC on a chiral phase. [e] Yield of the isolated *cis* isomer.

**Table 2:** Asymmetric cycloaddition of cyclopropanes with nitrones.<sup>[a]</sup>


Entry	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	t [days]	Yield [%] <sup>[b]</sup>	d.r. <sup>[c]</sup>	ee [%] <sup>[d]</sup>
1	Ph	Me	Ph	3	82 <sup>[e]</sup>	13:1	90
2 <sup>[f]</sup>	Ph	Bn	Ph	4	62 <sup>[e]</sup>	10:1	97
3	Ph	Et	Ph	4	88 <sup>[e]</sup>	11:1	95
4	Ph	Et	4-BrC <sub>6</sub> H <sub>4</sub>	4	85 <sup>[e]</sup>	12:1	97
5	Ph	Et	4-MeO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub>	4	97	11:1	97
6	Ph	Et	4-MeC <sub>6</sub> H <sub>4</sub>	4	80 <sup>[e]</sup>	12:1	96
7	Ph	Et	4-MeOC <sub>6</sub> H <sub>4</sub>	4	92 <sup>[e]</sup>	13:1	90
8	Ph	Et	2-furyl	4	99	13:1	93
9	Ph	Et	styryl	3	76 <sup>[e]</sup>	4:1	92
10 <sup>[f]</sup>	vinyl	Et	Ph	3	88 <sup>[e]</sup>	6:1	80
11 <sup>[f]</sup>	styryl	Et	Ph	5 <sup>[g]</sup>	84	5:1	80
12	Ph	Et	Ph <sup>[h]</sup>	5	74	11:1	93

[a] Reaction conditions: Ni(ClO<sub>4</sub>)<sub>2</sub>/1e = 0.040 mmol/0.044 mmol, 2/3 = 0.44 mmol/0.20 mmol, 4-Å molecular sieves. [b] Combined yield of the two diastereomers based on 3. [c] Ratio of isomers (*cis/trans*), determined by <sup>1</sup>H NMR spectroscopy. [d] Determined by HPLC on a chiral phase. [e] Yield of the isolated *cis* isomer. [f] T = -40 °C. [g] t = 5 h. [h] PhCH=N(O)Bn was used.


entries 1–3). The electronic character of the α-aryl group on the nitron had a slight effect on the enantioselectivity of the reaction (Table 2, entries 4–7). All reactions of both electron-deficient and electron-rich α-aryl nitrones proceeded with excellent enantio- and diastereoselectivities. A nitron with an α-furyl group underwent the cycloaddition smoothly with high diastereoselectivity to afford the corresponding 1,2-oxazine product in quantitative yield with a high *ee* value (Table 2, entry 8). A styryl-substituted nitron reacted to give the cycloaddition product with high enantioselectivity and good diastereoselectivity (Table 2, entry 9). Diethyl 2-vinyl- and 2-styrylcyclopropane-1,1-dicarboxylates reacted smoothly with 3b to give the desired products with good


enantio- and diastereoselectivities (Table 2, entries 10 and 11). Finally, the reaction of an *N*-Bn-substituted nitron also gave the cycloaddition adduct in good yield with excellent stereoselectivity (Table 2, entry 12).

In the [3+3] cycloaddition reactions described above, an excess of the cyclopropane derivative (2.2 equiv) was used. Interestingly, the remaining cyclopropane could be recovered with a good *ee* value. For example, in the case of entry 1 in Table 2, dimethyl 2-phenylcyclopropane-1,1-dicarboxylate was recovered in 57% yield with 59% *ee* (Scheme 2). These results, together with the importance of cyclopropane-1,1-dicarboxylates as intermediates as a result of their wide application in organic synthesis,<sup>[13]</sup> encouraged us to explore the kinetic resolution of 2-substituted cyclopropane-1,1-dicarboxylates. It was found that under optimized conditions the trisoxazoline-Ni(ClO<sub>4</sub>)<sub>2</sub> catalyst 1e/Ni(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O was also an excellent catalyst for the kinetic resolution of 2-substituted cyclopropane-1,1-dicarboxylates. Simply by changing the ratio of (±)-2b to 3b, a variety of 2-aryl cyclopropane-1,1-dicarboxylates could be resolved efficiently in high yields in the presence of a catalytic amount of 1e/Ni(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O (Table 3). This method provides ready access to optically active 2-substituted cyclopropane-1,1-dicarboxylates with excellent enantioselectivities (91–97% *ee*).<sup>[14]</sup>

Interestingly, optically active 2b reacted smoothly with nitron 3b in the presence of a catalytic amount of Ni(ClO<sub>4</sub>)<sub>2</sub> without the


ligand 1e to give the tetrahydro-1,2-oxazine 4c in high yield with the same level of enantiomeric purity as that of the starting material 2b (Scheme 3). This result further supports the mechanism proposed by Young and Kerr for the [3+3]


**Scheme 2.** Reaction of 3b with excess cyclopropane 2b.

**Table 3:** Kinetic resolution of 2-substituted cyclopropane-1,1-dicarboxylates.<sup>[a]</sup>


X	(±)-2 [mmol]	3b [mmol]	s <sup>[b]</sup>	t [h]	ee [%] <sup>[c]</sup>	Recovery of 2 [%] <sup>[d]</sup>
Ph	0.40	0.40	16	30	91	43
p-MeC <sub>6</sub> H <sub>4</sub>	0.40	0.25	97	48	96	49
p-BrC <sub>6</sub> H <sub>4</sub>	0.40	0.23	81	67	95	49
p-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	0.40	0.23	97	168	96	49
p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	0.40	0.23	36	96	97	45
p-ClC <sub>6</sub> H <sub>4</sub>	0.40	0.23	70	72	94	49
p-MeOC <sub>6</sub> H <sub>4</sub>	0.40	0.23	13	48	92	40

[a] Reaction conditions: Ni(ClO<sub>4</sub>)<sub>2</sub> (0.040 mmol), 1e (0.044 mmol), 4-Å molecular sieves. For details, see the Supporting Information. [b]  $s = \ln[(1-C)(1-ee)]/\ln[(1-C)(1+ee)]$ ; C refers to the conversion of (±)-2 (1 – (yield of recovered 2)). [c] Determined by HPLC on a chiral phase. [d] Yield of isolated recovered 2.

**Scheme 3.** Synthesis of both enantiomers of the 1,2-oxazine 4c.

cycloaddition of nitrones with cyclopropanes.<sup>[11a]</sup> Thus, both enantiomers of the tetrahydro-1,2-oxazine could be prepared from the racemic 2-phenyl cyclopropane-1,1-dicarboxylate: by direct cycloaddition with the nitrone in the presence of a catalytic amount of 1e/Ni(ClO<sub>4</sub>)<sub>2</sub> or by 1e/Ni(ClO<sub>4</sub>)<sub>2</sub>-catalyzed resolution followed by cycloaddition with the nitrone (Scheme 3).

In conclusion, we have developed a highly diastereoselective and enantioselective catalytic cycloaddition of 2-substituted cyclopropane-1,1-dicarboxylates with nitrones in the presence of a trisoxazoline–Ni<sup>II</sup> catalyst. The diastereoselectivity of the cycloaddition has been improved greatly relative to that of the existing method. The same catalyst 1e/Ni(ClO<sub>4</sub>)<sub>2</sub> also proved to be excellent for the kinetic resolution of 2-substituted cyclopropane-1,1-dicarboxylates. Furthermore, a combination of the asymmetric cycloaddition and the kinetic resolution/cycloaddition provided straightforward access to both enantiomers of tetrahydro-1,2-oxazines. The mild reaction conditions, atom economy, and high stereoselectivity make this reaction potentially useful for organic synthesis.

## Experimental Section

Typical procedure for the enantioselective [3+3] cycloaddition: A mixture of Ni(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O (0.040 mmol) and the trisoxazoline (0.044 mmol) in dimethoxyethane (1 mL) was stirred at 50 °C for 2 h under nitrogen. The mixture was then cooled to room temperature and added to the cyclopropane diester 2 (0.44 mmol) with a syringe. Activated 4-Å molecular sieves (100 mg) were added to the resulting solution. The mixture was stirred at –30 °C for 30 min, and then the nitrone (0.20 mmol) was added. When the reaction was complete (monitored by TLC), the mixture was passed rapidly through a glass funnel with a thin layer (20 mm) of silica gel (300–400 mesh), which was then washed with CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The filtrate was concentrated under reduced pressure, and the residue was purified by flash chromatography to afford the product.

The general procedure for the kinetic resolution of 2-substituted cyclopropane 1,1-diester is similar to that for the enantioselective [3+3] cycloaddition. The reaction was carried out at the desired temperature, and the ratio of the nitrone 3b to the 2-substituted cyclopropane diester (±)-2 was changed (see Table 3). The reaction was quenched when the conversion of (±)-2 was equal to or higher than 50 % (monitored by <sup>1</sup>H NMR spectroscopy).<sup>[15]</sup>

Received: November 15, 2006

Revised: January 26, 2007

Published online: April 17, 2007

**Keywords:** asymmetric catalysis · cycloaddition · cyclopropanes · kinetic resolution · trisoxazolines

- [1] a) J. Zhou, Y. Tang, *Chem. Soc. Rev.* **2005**, 34, 664; b) S. E. Gibson, M. P. Castaldi, *Angew. Chem.* **2006**, 118, 4834; *Angew. Chem. Int. Ed.* **2006**, 45, 4718; c) C. Moberg, *Angew. Chem.* **2006**, 118, 4838; *Angew. Chem. Int. Ed.* **2006**, 45, 4721.
- [2] For selected recent reports on the use of trisoxazolines in asymmetric catalysis, see: a) S. Bellemin-Lapponnaz, L. H. Gade, *Chem. Commun.* **2002**, 1286; b) S. Bellemin-Lapponnaz, L. H. Gade, *Angew. Chem.* **2002**, 114, 3623; *Angew. Chem. Int. Ed.* **2002**, 41, 3473; c) C. Dro, S. Bellemin-Lapponnaz, R. Welter, L. H. Gade, *Angew. Chem.* **2004**, 116, 4579; *Angew. Chem. Int. Ed.* **2004**, 43, 4479; d) B. D. Ward, S. Bellemin-Lapponnaz, L. H. Gade, *Angew. Chem.* **2005**, 117, 1696; *Angew. Chem. Int. Ed.* **2005**, 44, 1668; e) C. Foltz, B. Stecker, G. Marconi, S. Bellemin-Lapponnaz, H. Wadepohl, L. H. Gade, *Chem. Commun.* **2005**, 5115; f) M. Seitz, C. Capacchione, S. Bellemin-Lapponnaz, H. Wadepohl, B. D. Ward, L. H. Gade, *Dalton Trans.* **2006**, 193.
- [3] For selected recent reports on the use of trisoxazolines for molecular recognition, see: a) S.-G. Kim, K.-H. Kim, J. Jung, S. K. Shin, K. H. Ahn, *J. Am. Chem. Soc.* **2002**, 124, 591; b) J. Kim, S.-G. Kim, H. R. Seong, K. H. Ahn, *J. Org. Chem.* **2005**, 70, 7227; c) J. Kim, B. Raman, K. H. Ahn, *J. Org. Chem.* **2006**, 71, 38; d) J. Kim, D. Ryu, Y. Sei, K. Yamaguchi, K. H. Ahn, *Chem. Commun.* **2006**, 1136.
- [4] a) J. Zhou, Y. Tang, *J. Am. Chem. Soc.* **2002**, 124, 9030; b) J. Zhou, M.-C. Ye, Z.-Z. Huang, Y. Tang, *J. Org. Chem.* **2004**, 69, 1309; c) M.-C. Ye, B. Li, J. Zhou, X.-L. Sun, Y. Tang, *J. Org. Chem.* **2005**, 70, 6108.
- [5] a) M.-C. Ye, J. Zhou, Z.-Z. Huang, Y. Tang, *Chem. Commun.* **2003**, 2554; b) M.-C. Ye, J. Zhou, Y. Tang, *J. Org. Chem.* **2006**, 71, 3576.
- [6] Z.-Z. Huang, Y.-B. Kang, J. Zhou, M.-C. Ye, Y. Tang, *Org. Lett.* **2004**, 6, 1677.
- [7] For reviews on kinetic resolution, see: a) H. B. Kagan, J. C. Fiaud, *Top. Stereochem.* **1988**, 18, 249; b) J. M. Keith, J. F. Larrow, E. N. Jacobsen, *Adv. Synth. Catal.* **2001**, 343, 5; c) F. Cardona, A. Goti, A. Brandi, *Eur. J. Org. Chem.* **2001**, 2999;

- d) D. E. J. E. Robinson, S. D. Bull, *Tetrahedron: Asymmetry* **2003**, *14*, 1407; e) E. Vedejs, M. Jure, *Angew. Chem.* **2005**, *117*, 4040; *Angew. Chem. Int. Ed.* **2005**, *44*, 3974; for selected reports, see: f) M. Frohn, X. Zhou, J.-R. Zhang, Y. Tang, Y. Shi, *J. Am. Chem. Soc.* **1999**, *121*, 7718; g) A. Berkessel, F. Cleemann, S. Mukherjee, T. N. Müller, J. Lex, *Angew. Chem.* **2005**, *117*, 817; *Angew. Chem. Int. Ed.* **2005**, *44*, 807; h) A. Suárez, C. W. Downey, G. C. Fu, *J. Am. Chem. Soc.* **2005**, *127*, 11244; i) J. Paetzold, J. E. Bäckvall, *J. Am. Chem. Soc.* **2005**, *127*, 17620; j) K. Tanaka, Y. Hagiwara, M. Hirano, *Angew. Chem.* **2006**, *118*, 2800; *Angew. Chem. Int. Ed.* **2006**, *45*, 2734; k) A. Berkessel, M. Brandenburg, *Org. Lett.* **2006**, *8*, 4401; l) M. Tokunaga, J. Kiyosu, Y. Obora, Y. Tsuji, *J. Am. Chem. Soc.* **2006**, *128*, 4481; m) T. Murai, D. Matsuoka, K. Morishita, *J. Am. Chem. Soc.* **2006**, *128*, 4584; n) V. B. Birman, H. Jiang, X. Li, L. Guo, E. W. Uffman, *J. Am. Chem. Soc.* **2006**, *128*, 6536; o) S.-Y. Tosaki, K. Hara, V. Gnanadesikan, H. Morimoto, S. Harada, M. Sugita, N. Yamagiwa, S. Matsunaga, M. Shibasaki, *J. Am. Chem. Soc.* **2006**, *128*, 11776; p) S. Hoffmann, M. Nicoletti, B. List, *J. Am. Chem. Soc.* **2006**, *128*, 13074; q) F. O. Arp, G. C. Fu, *J. Am. Chem. Soc.* **2006**, *128*, 14264.
- [8] a) I. Uchida, S. Takase, H. Kayakiri, S. Kiyoto, M. Hashimoto, *J. Am. Chem. Soc.* **1987**, *109*, 4108; b) H. Terano, S. Takase, J. Hosoda, M. Kohsaka, *J. Antibiot.* **1989**, *42*, 145; c) Q.-S. Yu, X. Zhu, H. W. Holloway, N. F. Whittaker, A. Brossi, N. H. Greig, *J. Med. Chem.* **2002**, *45*, 3684; d) T. Katoh, E. Itoh, T. Yoshino, S. Terashima, *Tetrahedron* **1997**, *53*, 10229; e) T. C. Judd, R. M. Williams, *Angew. Chem.* **2002**, *114*, 4877; *Angew. Chem. Int. Ed.* **2002**, *41*, 4683; f) M. Suzuki, J. Kambe, H. Tokuyama, T. Fukuyama, *Angew. Chem.* **2002**, *114*, 4880; *Angew. Chem. Int. Ed.* **2002**, *41*, 4686.
- [9] a) R. Pulz, A. Al-Harrasi, H.-U. Reißig, *Org. Lett.* **2002**, *4*, 2353; b) A. A. Tishkov, H.-U. Reißig, S. L. Ioffe, *Synlett* **2002**, 863; c) M. Buchholz, H.-U. Reißig, *Eur. J. Org. Chem.* **2003**, 3524; d) A. Al-Harrasi, H.-U. Reißig, *Angew. Chem.* **2005**, *117*, 6383; *Angew. Chem. Int. Ed.* **2005**, *44*, 6227; for the total synthesis of (+)-phyllanthidine from a cyclopropane, see: e) C. A. Carson, M. A. Kerr, *Angew. Chem.* **2006**, *118*, 6710; *Angew. Chem. Int. Ed.* **2006**, *45*, 6560.
- [10] For reviews on the synthesis of 1,2-oxazines, see: a) T. L. Gilchrist, J. E. Wood in *Comprehensive Heterocyclic Chemistry II*, Vol. 6, Elsevier, Oxford, UK, **1996**, pp. 279–299 and 1177–1307; b) P. G. Tsoungas, *Heterocycles* **2002**, *57*, 1149; for the synthesis of 1,2-oxazines, see: c) S. C. Yoon, K. Kim, Y. J. Park, *J. Org. Chem.* **2001**, *66*, 7334; d) R. Pulz, S. Cicchi, A. Brandi, H.-U. Reißig, *Eur. J. Org. Chem.* **2003**, 1153; e) Y. Yamamoto, N. Momiyama, H. Yamamoto, *J. Am. Chem. Soc.* **2004**, *126*, 5962; f) S. Kumarn, D. M. Shaw, D. A. Longbottom, S. V. Ley, *Org. Lett.* **2005**, *7*, 4189; g) M. Helms, W. Schade, R. Pulz, T. Watanabe, A. Al-Harrasi, L. Fisira, I. Hlobilová, G. Zahn, H.-U. Reißig, *Eur. J. Org. Chem.* **2005**, 1003; h) S. Kumarn, D. M. Shaw, S. V. Ley, *Chem. Commun.* **2006**, 3211.
- [11] a) I. S. Young, M. A. Kerr, *Angew. Chem.* **2003**, *115*, 3131; *Angew. Chem. Int. Ed.* **2003**, *42*, 3023; b) I. S. Young, M. A. Kerr, *Org. Lett.* **2004**, *6*, 139; c) I. S. Young, J. L. Williams, M. A. Kerr, *Org. Lett.* **2005**, *7*, 953.
- [12] M. P. Sibi, Z. Ma, C. P. Jasperse, *J. Am. Chem. Soc.* **2005**, *127*, 5764.
- [13] For excellent reviews on the chemistry of donor-acceptor cyclopropanes, see: a) M. Yu, B. L. Pagenkopf, *Tetrahedron* **2005**, *61*, 321; b) H.-U. Reißig, R. Zimmer, *Chem. Rev.* **2003**, *103*, 1151; c) H. N. C. Wong, M.-Y. Hon, C.-W. Tse, Y.-C. Yip, J. Tanko, T. Hudlicky, *Chem. Rev.* **1989**, *89*, 165; d) S. Danishefsky, *Acc. Chem. Res.* **1979**, *12*, 66; for synthetic applications, see: e) E. J. Corey, T. G. Gant, *Tetrahedron Lett.* **1994**, *35*, 5373; f) P. E. Harrington, M. A. Kerr, *Tetrahedron Lett.* **1997**, *38*, 5949; g) M. A. Kerr, R. G. Keddy, *Tetrahedron Lett.* **1999**, *40*, 5671; h) D. B. England, T. D. O. Kuss, R. G. Keddy, M. A. Kerr, *J. Org. Chem.* **2001**, *66*, 4704; i) D. B. England, T. K. Woo, M. A. Kerr, *Can. J. Chem.* **2002**, *80*, 992; j) P. D. Pohlhaus, J. S. Johnson, *J. Org. Chem.* **2005**, *70*, 1057; k) P. D. Pohlhaus, J. S. Johnson, *J. Am. Chem. Soc.* **2005**, *127*, 16014; l) Y. B. Kang, Y. Tang, X. L. Sun, *Org. Biomol. Chem.* **2006**, *4*, 299.
- [14] For the asymmetric synthesis of 2-substituted cyclopropane-1,1-dicarboxylates, see: P. Müller, A. Ghanem, *Org. Lett.* **2004**, *6*, 4347.
- [15] For details, see the Supporting Information.