


A Chiral Pool Based Synthesis of Platensimycin**

K. C. Nicolaou,* Doron Pappo, Kit Y. Tsang, Romelo Gibe, and David Y.-K. Chen*

The fight against infectious disease rages on as scientists continue their struggle to overcome the emergence of drug resistance through new discoveries in chemistry and biology.^[1] The recent discovery of platensimycin (**1**, Scheme 1) by a Merck research team is a beautiful illustration of how natural products chemistry combined with modern biological techniques can lead to important developments in the search for new antibiotics.^[2] Platensimycin (**1**) was identified by high-throughput screening and RNA silencing technologies as a potent and selective inhibitor of the β -ketoacyl-(acyl-carrier-protein) synthase (FabF). Flexible synthetic routes to platensimycin may be particularly useful in accessing novel analogues that may prove superior to the natural product with regard to its pharmacological properties.^[3] Herein we report a new synthetic strategy that leads from the readily available and inexpensive (*R*)-(-)-carvone (**7**) to the tetracyclic enone **2**—a convenient precursor to platensimycin (**1**).

Scheme 1 outlines, in retrosynthetic format, the overall plan for the construction of the required enone **2** starting from (*R*)-(-)-carvone (**7**). Thus, enone **2** was expected to arise from tricyclic ketone **3**, whose connection to bicyclic keto aldehyde **4** could be recognized through a retro-anionic or radical 1,4-addition reaction. The origins of **4** were then traced to hydroxy keto acetal **5** by standard functional-group manipulations. Finally, **5** was envisioned to arise from enone **6** through a radical-based ring closure, with the latter intermediate being readily traced back to (*R*)-(-)-carvone (**7**) through a retro-Grignard reaction.



Scheme 1. Structure and retrosynthetic analysis of (-)-platensimycin (**1**).


According to the synthetic blueprint, the synthesis of **2** began from (*R*)-(-)-carvone (**7**, Scheme 2). Thus, 1,2-addition of Grignard reagent **8** to (-)-carvone (**7**) under Luche conditions (CeCl_3)^[4] led to the corresponding tertiary alcohol, which was directly oxidized (PCC) without further purification to afford enone **6** in 90% yield for the two steps. Intramolecular radical cyclization within the substituted carvone **6** with concomitant construction of the C8 quaternary center was accomplished through application of the sequential oxymercuration/reductive-alkylation methodology pioneered by Giese.^[5] In this instance, treatment of **6** with $\text{Hg}(\text{OAc})_2$ resulted in selective Markovnikov oxymercuration of the geminal disubstituted alkene. The intermediate organomercurial species was then reduced with NaBH_4 , presumably leading to the corresponding primary radical, which underwent 1,4-addition onto the enone system, thereby furnishing a mixture (ca. 1:1) of *exo* (**5a**) and *endo* (**5b**), existing predominantly as its hemiketal form **5b'**; **5b'**/**5b** ca. 6:1 alcohols in 61% yield. Dehydration of a mixture of **5a** + **5b/5b'** with Martin's sulfurane^[6a,b] led to exocyclic alkene **9**,^[6c] which was subjected to a regioselective silyl enol ether formation (TMSI, HMDS) followed by an electrophilic quench with PhSeCl , and subsequent oxidative elimination (H_2O_2) to give enone **10** in 36% yield from **5a** + **5b/5b'**. Finally, unveiling of the dioxane-masked aldehyde in **10** was achieved under micro-

[*] Prof. Dr. K. C. Nicolaou, Dr. D. Pappo, Dr. K. Y. Tsang, Dr. R. Gibe, Dr. D. Y.-K. Chen
Chemical Synthesis Laboratory@Biopolis
Institute of Chemical and Engineering Sciences (ICES)
Agency for Science, Technology and Research (A*STAR)
11 Biopolis Way, The Helios Block, no. 03-08, Singapore 138667 (Singapore)
Fax: (+1) 858-784-2469
E-mail: kcn@scripps.edu
david_chen@ices.a-star.edu.sg

Prof. Dr. K. C. Nicolaou
Department of Chemistry and
The Skaggs Institute for Chemical Biology
The Scripps Research Institute
10550 North Torrey Pines Road, La Jolla, CA 92037 (USA)
and
Department of Chemistry and Biochemistry
University of California, San Diego
9500 Gilman Drive, La Jolla, CA 92093 (USA)

[**] We thank Doris Tan (ICES) for high-resolution mass-spectrometric (HRMS) assistance. Financial support for this work was provided by A*STAR, Singapore.


Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



Scheme 2. Synthesis of dienone aldehyde **4**. Reagents and conditions: a) CeCl_3 (1.1 equiv), THF, 1 h; then **8** (1.0 equiv), $0 \rightarrow 23^\circ\text{C}$, 3 h; b) PCC (2.2 equiv), silica gel, CH_2Cl_2 , 2 h, 90% for the two steps; c) $\text{Hg}(\text{OAc})_2$ (1.1 equiv), THF/ H_2O (1:1), 23°C , 30 min, NaBH_4 (0.8 equiv), 20°C , 45 min, **5a**: 31%, **5b/5b'**: 30%, **6**: 20%; d) Martin's sulfurane (1.1 equiv), CH_2Cl_2 , 23°C , 12 h; e) Lil (2.5 equiv), TMSCl (2.5 equiv), HMDS (5.0 equiv), CH_2Cl_2 , $-35 \rightarrow 23^\circ\text{C}$, 3 h; then PhSeCl (1.15 equiv), CH_2Cl_2 , -78°C , 30 min, 42% for the two steps; f) 30% aq H_2O_2 (10.0 equiv), Py (2.5 equiv), CH_2Cl_2 , 0°C , 1.5 h, 85%; g) $\text{AcOH}/\text{H}_2\text{O}$ (5:1), 80°C , microwave, 65 min, 85%. PCC = pyridinium chlorochromate; TMS = trimethylsilyl; HMDS = hexamethyldisilazane; Py = pyridine.


wave-irradiation conditions in $\text{AcOH}/\text{H}_2\text{O}$ (5:1) at 80°C to furnish dienone aldehyde **4** in 85% yield.

The next objective was the forging of the final carbocyclic ring within the tetracyclic skeleton of **2**. For the accomplishment of this goal, an intramolecular Stetter reaction^[3d,7] and a SmI_2 -mediated^[3a,8] radical cyclization were the available options. Thus, a Stetter reaction of aldehyde **4** catalyzed by thiazolium salt **11**^[7] in an ethanolic solution containing Et_3N yielded an inseparable mixture of C9 epimeric diketones **3a** and **3b** in 65% combined yield (**3a/3b** ca. 5:1; scheme 3). Interestingly, however, treatment of **4** with SmI_2 in THF/MeOH at room temperature resulted in the formation of hydroxy ketone **12** possessing the undesired C9 configuration as a single stereoisomer in 57% yield. A chelated transition state (**18**, Scheme 4a) is postulated as a plausible explanation for this highly stereoselective process. Accessing greater quantities of the desired C9 stereoisomer was realized through a base-mediated (KOH) equilibration process.



Scheme 3. Synthesis of tetracyclic enone **2**. Reagents and conditions: a) **11** (0.2 equiv), Et_3N (1.2 equiv), EtOH, 80°C , 18 h, 65% (**3a/3b** ca. 5:1); b) SmI_2 (0.1 M in THF, 2.2 equiv), MeOH (2.5 equiv), 23°C , THF (0.032 M), 2 min, 57%; c) *p*-nitrobenzoic acid (1.5 equiv), PPh_3 (1.5 equiv), DIAD (3.4 equiv), benzene, 23°C , 2 h, 67%; d) KOH (10% in MeOH), 50°C , 5 h, **14a**: 46%, **14b**: 45%; e) L-selectride (1.0 M in THF, 4.0 equiv), THF, $-78 \rightarrow 23^\circ\text{C}$, 2 h; then 1 M aq HCl, $0 \rightarrow 23^\circ\text{C}$, 2 h, 80%; f) PCC (2.0 equiv), CH_2Cl_2 , 23°C , 1.5 h, 95%; g) TMSCl (3.3 equiv), Lil (3.3 equiv), HMDS (5.0 equiv), CH_2Cl_2 , 0°C , 3 h; h) 1. IBX (1.5 equiv), MPO (1.5 equiv), DMSO, 23°C , 2 h, **2**: 53%, **17**: 26%; or 2. $\text{Pd}(\text{OAc})_2$ (1.0 equiv), CH_3CN , 23°C , 5 h, **2**: 60%, **17**: 30%; DIAD = diisopropyl azodicarboxylate; IBX = *o*-iodoxybenzoic acid; MPO = 4-methoxy-pyridine-*N*-oxide.

While diketone **3a** proved unstable and hydroxy ketone **12** appeared resistant to epimerization under the basic conditions employed (KOH/MeOH), ester ketone **13**, obtained through Mitsunobu inversion of the C5 hydroxy group (*p*- $\text{NO}_2\text{C}_6\text{H}_4\text{CO}_2\text{H}/\text{DIAD}/\text{PPh}_3$, 67% yield) within **12**, underwent smooth and facile equilibration in the presence of KOH with concomitant *p*-nitrobenzoate ester hydrolysis, to afford a readily separable mixture of C9 epimers (**14a/14b** ca. 1.1:1, 91% yield). As seen in Scheme 4, this substrate-dependent epimerization could be rationalized by the apparent slight preference for the equatorially oriented C5 hydroxy group,



Scheme 4. a) Postulated samarium-templated ring closure of radical **18** to form hydroxy ketone **12**; and b) base-mediated equilibration of hydroxy ketones **14a** and **14b**.

hence the reluctance of hydroxy ketone **12** to undergo the similar equilibration process (see Scheme 4b). In preparation for the casting of the remaining tetrahydrofuran system, stereoselective reduction of the C10 ketone within **14b** was carried out with L-selectride, which resulted, upon acidic workup (1N aq HCl), in the formation of the desired cage hydroxy compound **15**, in 80% yield. Sequential oxidation of the latter intermediate with PCC, followed by IBX^[9] or Saegusa^[10] oxidation of the silyl enol ether of **16** (TMSCl, LiI, HMDS; then IBX or Pd(OAc)₂), furnished a readily separable mixture of enone **2** and its regioisomer **17** (IBX: 79% overall yield, **2/17** ca. 2:1; Pd(OAc)₂: 90% overall yield, **2/17** ca. 2:1).^[11] The recycling of **17** by conversion into **2** has been demonstrated,^[3f,g] as has the conversion of **2** into **1**.^[3a-c] Enone **2** exhibited identical physical properties (¹H and ¹³C NMR spectra, [α]_D values, and mass spectra) to those reported previously.^[3]

In conclusion, an efficient and stereocontrolled synthesis of the advanced intermediate **2** en route to (–)-platensimycin (**1**) has been accomplished by using (R)-(–)-carvone as an inexpensive chiral starting material. This expedient and flexible entry, which is applicable to both enantiomers of platensimycin, should offer opportunities for the synthesis of designed platensimycin analogues for chemical biology investigations.^[12]

Received: November 2, 2007

Keywords: antibiotics · natural products · radicals · synthesis design

[1] D. Niccolai, L. Tarsi, R. J. Thomas, *Chem. Commun.* **1997**, 2333–2342.

- [2] a) J. Wang, S. M. Soisson, K. Young, W. Shoop, S. Kodali, A. Galgoci, R. Painter, G. Parthasarathy, Y. S. Tang, R. Cummings, S. Ha, K. Dorso, M. Motyl, H. Jayasuriya, J. Ondeyka, K. Herath, C. Zhang, L. Hernandez, J. Allocco, A. Basilio, J. R. Tormo, O. Genilloud, F. Vicente, F. Pelaez, L. Colwell, S. H. Lee, B. Michael, T. Felcetto, C. Gill, L. L. Silver, J. D. Hermes, K. Bartizal, J. Barrett, D. Schmatz, J. W. Becker, D. Cully, S. B. Singh, *Nature* **2006**, *441*, 358–361; b) S. B. Singh, H. Jayasuriya, J. G. Ondeyka, K. B. Herath, C. Zhang, D. L. Zink, N. N. Tsou, R. G. Ball, A. Basilio, O. Genilloud, M. T. Diez, F. Vicente, F. Pelaez, K. Young, J. Wang, *J. Am. Chem. Soc.* **2006**, *128*, 11916–11920; c) D. Häbich, F. von Nussbaum, *ChemMedChem* **2006**, *1*, 951–954.
- [3] For the total and formal syntheses of platensimycin, in racemic and enantiopure forms, see a) K. C. Nicolaou, A. Li, D. J. Edmonds, *Angew. Chem.* **2006**, *118*, 7244–7248; *Angew. Chem. Int. Ed.* **2006**, *45*, 7086–7090; b) K. C. Nicolaou, D. J. Edmonds, A. Li, G. S. Tria, *Angew. Chem.* **2007**, *119*, 4016–4019; *Angew. Chem. Int. Ed.* **2007**, *46*, 3942–3945; c) Y. Zou, C.-H. Chen, C. D. Taylor, B. M. Foxman, B. B. Snider, *Org. Lett.* **2007**, *9*, 1825–1828; d) K. C. Nicolaou, Y. Tang, J. Wang, *Chem. Commun.* **2007**, 1922–1923; e) P. Li, J. N. Payette, H. Yamamoto, *J. Am. Chem. Soc.* **2007**, *129*, 9534–9535; f) K. Tiefenbacher, J. Mulzer, *Angew. Chem.* **2007**, *119*, 8220–8221; *Angew. Chem. Int. Ed.* **2007**, *46*, 8074–8075; *Angew. Chem.* **2007**, *119*, 8220–8221; *Angew. Chem. Int. Ed.* **2007**, *46*, 8074–8075; g) G. Lalic, E. J. Corey, *Org. Lett.* **2007**, *9*, 4921–4923; for other synthetic studies towards the tetracyclic core of platensimycin, see h) K. P. Kaliappan, V. Ravikumar, *Org. Lett.* **2007**, *9*, 2417–2419; i) A. K. Ghosh, K. Xi, *Org. Lett.* **2007**, *9*, 4013–4016.
- [4] G. A. Molander, *Chem. Rev.* **1992**, *92*, 29–68.
- [5] a) B. Giese, *Radicals in Organic Synthesis: Formation of Carbon-Carbon Bonds*, Pergamon Press, Oxford, **1986**; b) K. Weinges, H. Reichert, *Synlett* **1991**, 785–786.
- [6] a) J. C. Martin, R. J. Arhart, *J. Am. Chem. Soc.* **1971**, *93*, 4327–4329; b) R. J. Arhart, J. C. Martin, *J. Am. Chem. Soc.* **1972**, *94*, 5003–5010; c) from the several dehydration protocols examined, the one involving Martin's sulfurane was the only one that gave exclusively the exocyclic olefin **9**, and thus able to effect this transformation on both **5a** and **5b/5b'**.
- [7] a) D. Enders, K. Breuer, J. Runsink, J. H. Teles, *Acc. Chem. Res.* **2004**, *37*, 534–541; b) J. S. Johnson, *Angew. Chem.* **2004**, *116*, 1348–1350; *Angew. Chem. Int. Ed.* **2004**, *43*, 1326–1328.
- [8] D. J. Edmonds, D. Johnston, D. J. Procter, *Chem. Rev.* **2004**, *104*, 3371–3403.
- [9] K. C. Nicolaou, D. L. F. Gray, T. Montagnon, S. T. Harrison, *Angew. Chem.* **2002**, *114*, 1038–1042; *Angew. Chem. Int. Ed.* **2002**, *41*, 996–1000.
- [10] Y. Ito, T. Hirao, T. Saegusa, *J. Org. Chem.* **1978**, *43*, 1011–1013.
- [11] An alternative silyl enol ether formation (Me₃N, TMSOTf (Tf = trifluoromethanesulfonyl)) has been reported, which upon subsequent oxidation (IBX) yielded enone **2** with higher regioselectivity (**2/17** ca. 7:1), see Ref. [3g].
- [12] a) K. C. Nicolaou, T. Lister, R. M. Denton, A. Montero, D. J. Edmonds, *Angew. Chem.* **2007**, *119*, 4796–4798; *Angew. Chem. Int. Ed.* **2007**, *46*, 4712–4714; b) K. C. Nicolaou, Y. Yang, A. F. Stepan, A. Li, A. Montero, *J. Am. Chem. Soc.* **2007**, *129*, 14850–14851.