

Dihydrogen Activation

Heterolytic Dihydrogen Activation by a Sulfido- and Oxo-Bridged Dinuclear Germanium–Ruthenium Complex**


Tsuyoshi Matsumoto, Yukiko Nakaya, and Kazuyuki Tatsumi*

Heterolytic cleavage of dihydrogen on transition metal sulfides/thiolates has been regarded as the key reaction in hydrogen metabolism in nature^[1] and in catalytic desulfurization of fossil fuel.^[2] Although there are numerous reports of heterolytic dihydrogen activation,^[3,4] that occurring at metal–sulfur bonds is still limited. A representative example is the reaction of $[\{\text{Rh}(\text{triphos})\}_2(\mu\text{-S})]^{2+}$ (triphos = tris(diphenylphosphanylmethyl)methane) with H_2 to generate $[\{\text{RhH}(\text{triphos})\}_2(\mu\text{-SH})]^{2+}$.^[4a,b] Heterolytic H_2 cleavage was also reported to be promoted by sulfido/thiolato-bridged dinuclear Mo–Mo,^[4c–e] Ir–Ir,^[4f] and W–Ir^[4g] complexes and mononuclear sulfido/thiolato complexes of Ti,^[4h–j] Ni, Ru, and Rh.^[4j–n] In the course of our studies on transition metal sulfide/thiolate complexes,^[5] we found that sulfido-bridged W–Ru complexes activate H_2 in a heterolytic manner.^[5c]

Here we report heterolytic cleavage of H_2 by the sulfido- and oxo-bridged heterodinuclear germanium–ruthenium complex $[(\text{dmp})(\text{dep})\text{Ge}(\mu\text{-S})(\mu\text{-O})\text{Ru}(\text{PPh}_3)]$ (**1**; dmp = 2,6-dimesitylphenyl, dep = 2,6-diethylphenyl). As we showed previously, the $\mu\text{-S}$ ligand of **1** prefers softer acids, and the $\mu\text{-O}$ ligand harder acids.^[6] The synergetic $\mu\text{-sulfide}$ and $\mu\text{-oxide}$ pair plays an important role in H_2 heterolysis by **1**.

Heterodinuclear complex **1** was prepared from $[(\text{dmp})(\text{dep})\text{Ge}(\text{SH})(\text{OH})]$, $[\text{RuCl}_2(\eta^6\text{-p-cymene})]$, and PPh_3 according to Scheme 1.^[6] No H_2 activation by **1** took place under an atmospheric pressure of dihydrogen even at 90 °C. However, complex **1** was converted slowly to *anti* and *syn* isomers of hydroxy hydride complex **2** when heated to 75 °C in toluene under 10 atm of H_2 (Scheme 1, Table 1). As is obvious from the structures of *anti*-**2** and *syn*-**2**, H_2 was cleaved heterolytically by **1** into a hydroxy-bound proton on Ge and a hydride ligand on Ru.

Dihydrogen activation was examined under various conditions to obtain insight into the reaction mechanism. Intriguingly, *anti*-**2** is the favored product in the early stage of the reaction (Table 1, entry 1). The relative ratio of *syn*-**2** to



Scheme 1. Synthesis of **1** and the H_2 activation.

Table 1: Product ratio for H_2 activation by **1**.

Entry	Reaction conditions				Product ratio ^[a]			
	<i>T</i> [°C]	<i>t</i> [h]	<i>p</i> (H_2) [atm]	Additive	1	<i>anti</i> - 2	<i>syn</i> - 2	<i>anti</i> - 2 / <i>syn</i> - 2
1	75	6	10	–	32	52	16	3.3
2	75	6	10	PPh_3 ^[b]	36	49	15	3.3
3	75	24	10	–	2	73	25	2.9
4	75	24	10	PPh_3 ^[b]	2	71	27	2.6
5	90	72	7.5	–	0	0	100	0.0
6	90	72	7.5	PPh_3 ^[b]	0	34	66	0.5

[a] The product ratio was estimated by ^1H NMR spectroscopy. Complexes **1** and/or *anti*-**2**, *syn*-**2** were exclusively observed by ^1H NMR spectroscopy. [b] 10 equiv PPh_3 .

anti-**2** slowly increases as the reaction proceeds, and eventually *syn*-**2** becomes the exclusive product at 90 °C after 3 days under 7.5 atm H_2 (Table 1, entry 5). The results suggest that the kinetically favored product is *anti*-**2**, which gradually isomerizes to the more thermodynamically stable *syn*-**2**.


Addition of 10 equiv PPh_3 hardly affects the reaction of **1** with H_2 . No significant deceleration of the consumption of **1** was observed, either for the reactions at 75 °C for 6 h (Table 1, entries 1 and 2) or for those at 24 h (Table 1, entries 3 and 4), that is, the rate-determining step does not include PPh_3 dissociation. On the other hand, the subsequent isomerization from *anti*-**2** to *syn*-**2** is clearly decelerated by addition of PPh_3 , as is manifested in the results at 90 °C and 72 h (Table 1, entries 5 and 6). The isomerization appears to be accompanied by dissociation of PPh_3 .

[*] Dr. T. Matsumoto, Y. Nakaya, Prof. Dr. K. Tatsumi
Department of Chemistry
Graduate School of Science, and
Research Center for Materials Science
Nagoya University
Furo-cho, Chikusa-ku, Nagoya 464-8602 (Japan)
Fax: (+81) 52-789-2943
E-mail: i45100a@nucc.cc.nagoya-u.ac.jp

[**] This work was supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 18065013, “Chemistry of Concerto Catalysis”) from Ministry of Education, Culture, Sports, Science and Technology (Japan).

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


On the basis of these facts, we propose the mechanism of H_2 heterolysis summarized in Scheme 2. As a premise, H_2 activation between two molecules of **1** is excluded, because of the steric bulk of **1**.^[6] The activation of H_2 may occur stepwise



Scheme 2. Proposed mechanism of H_2 heterolysis by **1**. Some of the substituents on Ge and Ru are omitted for clarity.

via initial coordination to the Ru atom with slippage of the η^6 -arene and successive heterolytic cleavage.^[7] On the other hand, the H_2 molecule could undergo a straightforward σ -bond metathesis.^[8] We do not have any evidence to support either pathway.

If H_2 heterolysis occurs at the Ru–O bond (path b), it straightforwardly affords *syn*-**2**. However, path b does not account for the formation of *anti*-**2** as the kinetic product. The most conceivable pathway affording *anti*-**2** proceeds by H_2 heterolysis at the Ru–S bond (path a). This pathway should form **X**, which would be further transformed into *anti*-**2** by migration of the SH proton to the μ -oxo ligand with concomitant Ru–S bond formation and Ru–O bond dissociation. The proposed intermediate **X** has not been detected. We examined H_2 activation with a bis(μ -S) analogue of **1**, namely, $[(\text{dmp})(\text{dep})\text{Ge}(\mu\text{-S})_2\text{Ru}(\text{PPh}_3)]$ (**3**), at 90°C under 10 atm H_2 , anticipating the formation of the μ -S analogue of **X**. However, complex **3** was recovered quantitatively [Eq. (1)]. This result suggests that H_2 activation at Ru–S bonds is reversible, and furthermore that intermediate **X** and its μ -S analogue are considerably less stable than **1** and **3**, respectively. It seems that the key to H_2 activation via path a is subsequent proton migration onto the hard μ -oxo ligand to generate stable *anti*-**2**.



The hard nature of the μ -oxo ligand of **1** was demonstrated by μ -O-protonation of **1**.^[6] The preference for path a could be due to the softness of the μ -S compared to the μ -O moiety, as was also observed in the reaction of **1** with MeOTf.^[6]

Isomerization of *anti*-**2** to *syn*-**2** proceeds via dissociation of the phosphine ligand. Although the mechanism of this isomerization process is not entirely clear, it may involve weak H_2 coordination at Ru to compensate for PPh_3 dissociation. In fact, isomerization of preformed *anti*-**2** is even slower in the absence of H_2 , for example, requiring 9 days in toluene at 90°C . A basis for the relative thermodynamic stability of *syn*-**2** compared to that of *anti*-**2** can be deduced from their crystal structures (Figure 1).^[9] The




Figure 1. ORTEP drawings of a) *anti*-**2** and b) *syn*-**2**.

strained *m*-terphenyl arrangement of dmp in *anti*-**2** is reflected in the significantly smaller dihedral angle of 67° between the central arene ring of dmp and its Ru-coordinated mesityl ring, while those for both **1** and *syn*-**2** are 87° . The strained dmp conformation in *anti*-**2** is attributable to intramolecular steric congestion between the triphenylphosphine ligand and the hydroxy group, whereas the conformational strain in dmp is relieved in the structure of *syn*-**2**. The $\text{GeOH}\cdots\text{HRu}$ interaction for *syn*-**2** also contributes to its relative thermodynamic stability. The $\text{H}\cdots\text{H}$ interaction is indicated by the H54–H55 and O1–H54 distances of 2.25(3) and 2.95(2) Å, respectively, as derived from X-ray data.^[10] This nonclassical hydrogen bonding is also manifested in the ^1H NMR spectrum. An nOe measurement on *syn*-**2** showed 17% enhancement of GeOH at $\delta = 2.73$ ppm on irradiation of RuH at $\delta = -9.04$ ppm in C_6D_6 . In the IR spectrum, the lower frequency of the Ru–H stretching band of *syn*-**2** at

1940 cm⁻¹ relative to that of *anti*-**2** at 2015 cm⁻¹ may also indicate a nonclassical hydrogen-bonding interaction.

In conclusion, the S/O-bridged dinuclear germanium–ruthenium complex **1** activates H₂ heterolytically at the ruthenium–chalcogen bonds. Cooperation of the μ-S and μ-O atoms is significant in the mechanism of H₂ activation.

Received: October 23, 2007

Revised: November 23, 2007

Published online: January 25, 2008

Keywords: germanium · heterometallic complexes · oxo ligands · ruthenium · sulfido ligands

- [1] a) M. W. W. Adams, E. I. Steifel, *Science* **1998**, 282, 1842; b) R. Cammack, *Nature* **1999**, 397, 214.
- [2] H. Topsoe, B. S. Clausen, F. E. Massoth, *Hydrotreating Catalysis, Science and Technology*, Springer, Berlin, **1996**.
- [3] Recent reports: a) S. Ogo, R. Kabe, K. Uehara, B. Kure, T. Nishimura, S. C. Menon, R. Harada, S. Fukuzumi, Y. Higuchi, T. Ohhara, T. Tamada, R. Kuroki, *Science* **2007**, 316, 585; b) C. M. Nagaraja, P. Parameswaran, E. D. Jemmis, B. R. Jagirdar, *J. Am. Chem. Soc.* **2007**, 129, 5587; c) A. D. Wilson, R. K. Shoemaker, A. Miedaner, J. T. Muckerman, D. L. Dubois, M. Rakowski Dubois, *Proc. Natl. Acad. Sci. USA* **2007**, 104, 6951; d) A. D. Wilson, R. H. Newell, M. J. McNevin, J. T. Muckerman, M. Rakowski Dubois, D. L. Dubois, *J. Am. Chem. Soc.* **2006**, 128, 358; e) G. C. Welch, R. R. S. Juan, J. D. Masuda, D. W. Stephan, *Science* **2006**, 314, 1124; f) G. C. Welch, D. W. Stephan, *J. Am. Chem. Soc.* **2007**, 129, 1880.
- [4] a) C. Bianchini, C. Mealli, A. Meli, M. Sabat, *Inorg. Chem.* **1986**, 25, 4617; b) C. Bianchini, A. Meli, *Inorg. Chem.* **1987**, 26, 4268; c) J. C. R. Laurie, L. Duncan, R. C. Haltiwanger, R. T. Weberg, M. Rakowski DuBois, *J. Am. Chem. Soc.* **1980**, 102, 7456; d) M. Rakowski DuBois, *Chem. Rev.* **1989**, 89, 1; e) J. C. V. Laurie, L. D. Duncan, R. C. Haltiwanger, R. T. Weberg, M. Rakowski DuBois, *J. Am. Chem. Soc.* **1986**, 108, 6234; f) R. C. Linck, R. J. Pafford, T. B. Rauchfuss, *J. Am. Chem. Soc.* **2001**, 123, 8856; g) H. Kato, H. Seino, Y. Mizobe, M. Hidai, *J. Chem. Soc. Dalton Trans.* **2002**, 1494; h) Z. K. Sweeney, J. L. Polse, R. A. Andersen, R. G. Bergman, M. G. Kubinec, *J. Am. Chem. Soc.* **1997**, 119, 4543; i) Z. K. Sweeney, J. L. Polse, R. G. Bergman, R. A. Andersen, *Organometallics* **1999**, 18, 5502; j) D. Sellmann, R. Prakash, F. W. Heinemann, M. Moll, M. Klimowicz, *Angew. Chem.* **2004**, 116, 1913; *Angew. Chem. Int. Ed.* **2004**, 43, 1877; k) D. Sellmann, F. Geipel, M. Moll, *Angew. Chem.* **2000**, 112, 570; *Angew. Chem. Int. Ed.* **2000**, 39, 561; l) D. Sellmann, A. Fürsattel, *Angew. Chem.* **1999**, 111, 2142; *Angew. Chem. Int. Ed.* **1999**, 38, 2023; m) D. Sellmann, T. Gottschalk-Gaudig, F. W. Heinemann, *Inorg. Chem.* **1998**, 37, 3982; n) D. Sellmann, G. H. Rackelmann, F. W. Heinemann, *Chem. Eur. J.* **1997**, 3, 2071.
- [5] a) Z. Li, Y. Ohki, K. Tatsumi, *J. Am. Chem. Soc.* **2005**, 127, 8950; b) Y. Ohki, H. Sadohara, Y. Takikawa, K. Tatsumi, *Angew. Chem.* **2004**, 116, 2340; *Angew. Chem. Int. Ed.* **2004**, 43, 2290; c) Y. Ohki, N. Matsuura, T. Marumoto, H. Kawaguchi, K. Tatsumi, *J. Am. Chem. Soc.* **2003**, 125, 7978; d) Y. Ohki, Y. Sunada, M. Honda, M. Katada, K. Tatsumi, *J. Am. Chem. Soc.* **2003**, 125, 4052.
- [6] T. Matsumoto, Y. Nakaya, K. Tatsumi, *Organometallics* **2006**, 25, 4835.
- [7] The mechanism of H₂ heterolysis over [[Rh(triphos)]₂(μ-S)₂]²⁺ was studied by DFT calculations: A. Ienco, M. J. Calhorda, J. Reinhold, F. Reineri, C. Bianchini, M. Peruzzini, F. Vizza, C. Mealli, *J. Am. Chem. Soc.* **2004**, 126, 11954.
- [8] S. Niu, M. B. Hall, *Chem. Rev.* **2000**, 100, 353.
- [9] The metric parameters are included in the Supporting Information. CCDC-610035 (*anti*-**2**) and -610036 (*syn*-**2**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif
- [10] The typical range of nonclassical hydrogen-bond lengths is reported to be 1.7–2.2 Å: “Hydrides and Hydrogen Bonding: Combining Theory with Experiment”: E. Clot, O. Eisenstein, D.-H. Lee, R. H. Crabtree in *Recent Advances in Hydride Chemistry* (Eds.: M. Peruzzini, R. Poli), Elsevier, Amsterdam, **2001**, p. 75.