


Enzyme Catalysis

K. Kano,* Y. Ishida

Regulation of α -Chymotrypsin Catalysis by Ferric Porphyrins and Cyclodextrins

Chem. Asian J.

DOI: 10.1002/asia.200700383



Put the stopper on: The anionic ferric porphyrin FeTPPS inhibits the catalysis of α -chymotrypsin (ChT) by covering the catalytic site with its rigid porphyrin ring. The inhibited catalysis is completely recovered by per-*O*-methylated β -cyclodextrin (TMe- β -CD), which detaches FeTPPS from the protein surface. These results led to the preparation of a more useful inhibitor.


Directed Evolution

M. T. Reetz,* M. Rentzsch, A. Pletsch, A. Taglieber, F. Hollmann, R. J. G. Mondière, N. Dickmann, B. Höcker, S. Cerrone, M. C. Haeger, R. Sterner

A Robust Protein Host for Anchoring Chelating Ligands and Organocatalysts

ChemBioChem

DOI: 10.1002/cbic.200700413



Robust hybrid catalysts: A platform for directed evolution of hybrid catalysts has been optimized and miniaturized by using a thermostable enzyme, tHisF (see figure) from *Thermotoga maritima*. A tHisF mutant with a strategically placed cysteine was identified and subjected to bioconjugation by introducing a variety of ligands for metal ligation, a ligand/metal moiety, and several organocatalysts such as flavin and thiazolium entities.


Atomic Force Microscopy

J. Sotres, A. Lostao, L. Wildling, A. Ebner, C. Gómez-Moreno, H. J. Gruber, P. Hinterdorfer, A. M. Baró*

Unbinding Molecular Recognition Force Maps of Localized Single Receptor Molecules by Atomic Force Microscopy

ChemPhysChem

DOI: 10.1002/cphc.200700597



Simultaneous recordings: Molecular recognition forces on ligand-receptor pairs (biotin-avidin) are measured by AFM simultaneously with the topography of the receptor molecules (see figure). Spatially resolved ligand unbinding events in single molecules of 5 nm diameter are obtained.


Antibiotics Synthesis

F. von Nussbaum,* S. Anlauf, C. Freiberg, J. Benet-Buchholz, J. Schamberger, T. Henkel, G. Schiffer, D. Häbich


Total Synthesis and Initial Structure–Activity Relationships of Longicatenamycin A

ChemMedChem

DOI: 10.1002/cmdc.200700297



What is the active principle of the “S-520 longicatenamycin antibiotic complex”? Longicatenamycin A (shown) is the first defined longicatenamycin congener that has been totally synthesized and tested in pure form.



4,4,5,5-Tetraphenyl-1,3,2-dioxaborolane (HBBzpin) has been prepared in high yield and used in the catalysed hydroboration of alkenes to give air- and chromatography-stable organoboronate ester products. Reactions with metal complexes have also been investigated. Addition of HBBzpin to $\text{RhCl}(\text{PPh}_3)_3$ gave the borylrhodium complex $\text{Rh}(\text{H})\text{Cl}(\text{BzBpin})\text{-(PPh}_3)_2$.


Metal-Catalysed Hydroboration

C. B. Fritsch, S. M. Wernitz,
C. M. Vogels, M. P. Shaver, A. Decken,
A. Bell,* S. A. Westcott*

4,4,5,5-Tetraphenyl-1,3,2-dioxaborolane:
A Bulky Borane for the Transition Metal
Catalysed Hydroboration of Alkenes

Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.200701066

Synthetic strategies for oseltamivir phosphate, an important orally active anti-influenza drug, are reviewed.




Anti-influenza Drugs

M. Shibasaki,* M. Kanai

Synthetic Strategies for Oseltamivir
Phosphate

Eur. J. Org. Chem.
DOI: 10.1002/ejoc.200800033



Electron-spin relaxation is a key parameter in the design of advanced MRI contrast agents. EPR has been used to probe the relationship between this key


parameter and the coordination environment of gadolinium in complexes with DOTA-type ligands (see figure).

Contrast Agents

A. Borel,* J. F. Bean, R. B. Clarkson,
L. Helm, L. Moriggi, A. D. Sherry,
M. Woods*

Towards the Rational Design of MRI
Contrast Agents: Electron Spin
Relaxation Is Largely Unaffected by the
Coordination Geometry of
Gadolinium(III)–DOTA-Type Complexes

Chem. Eur. J.
DOI: 10.1002/chem.200701747



An ionic liquid: Investigation of the mechanism of the Beckmann rearrangement of cyclohexanone oxime to ϵ -caprolactam (**1**) in sulfuric acid or oleum has led to the conclusion that the manufac-

turing process for ϵ -caprolactam is in fact the largest-scale industrial technology that has been using an ionic liquid, caprolactamium hydrogen sulfate (**2**), as the reaction medium for decades.

Industrial Chemistry

V. Fábos, D. Lantos, A. Bodor,
A.-M. Bálint, L. T. Mika, O. E. Sielcken,
A. Cuiper, I. T. Horváth*

ϵ -Caprolactamium Hydrogen Sulfate: An
Ionic Liquid Used for Decades in the
Large-Scale Production of ϵ -Caprolactam

ChemSusChem
DOI: 10.1002/cssc.200700135