

The Synthesis and Isolation of a Metal-Substituted Bis-silene**


Dmitry Bravo-Zhivotovskii,* Roman Dobrovetsky, Dmitry Nemirovsky, Victoria Molev, Michael Bendikov, Gregory Molev, Mark Botoshansky, and Yitzhak Apeloig*

Dedicated to Professor Robert West

Since the isolation of the first stable silene ($R_2Si=CR_2$)^[1] and disilene ($R_2Si=SiR_2$)^[2], the field of multiply bonded silicon compounds has developed rapidly and led to the isolation of many novel compounds.^[3] Yet this field of chemistry is still in its infancy. In contrast to metal-substituted olefins, which are common reagents,^[4] only few metal-substituted unsaturated silicon compounds have been isolated and characterized,^[5] and found to have interesting reactivity.^[6] Few silene-metal complexes, in which the transition metal is coordinated to a Si=C bond have been reported.^[7] Metal-substituted silenes are still unknown.^[8]

In this paper we report the synthesis, isolation, and X-ray molecular structure of the first metal-substituted silene **1**, which was obtained by a Brook-type thermal rearrangement (Scheme 1). Compound **1** is also the first isolated and fully characterized bis-silene, a compound with two Si=C bonds.

Bis-silene **1** was synthesized in a single-pot reaction by mixing bis(lithiosilyl)mercury **2**^[9] with two equivalents of 1-adamantoyl chloride (**3**) in toluene at -78°C . After the reaction mixture was warmed to room temperature and stirred for 1 h, **1** was obtained as the only silicon-containing product. The reaction probably proceeds by means of a spontaneous Brook rearrangement of short-lived bis-acylsilane **4** (Scheme 1). Although **4** was not directly observed, we believe that **4** is an intermediate in the reaction, based on the isolation of an analogous compound, **7** (see below). Bis-silene **1** was isolated (80% yield) by crystallization from hexane at room temperature, and its molecular structure was determined by X-ray crystallography (Figure 1).^[10]



Scheme 1. Synthesis of **1**.

The average Si=C (1.764(15) Å) and Si-Si (2.369(6) Å) bond lengths in **1** are similar to the analogous bond lengths in Brook's silene ($[(Me_3Si)_2Si=C(OSiMe_3)Ad]$ (**5**) (Si=C 1.76 Å, Si-Si 2.34 Å).^[11] This shows that replacement of a R_3Si group by mercury has a small effect on the C=Si bond length. The Si1-Hg1 and Si4-Hg1 bond lengths in **1** (2.417(4) Å and 2.477(4) Å, respectively) are similar to that in $[(Me_3Si)_3Si]_2Hg$ (2.47 Å).^[11] The double-bonded Si1 and Si4 atoms in **1** are essentially planar (sum of bond angles around Si is 359.9°), suggesting sp^2 hybridization. The Si-Hg-Si bond angle is $177.4(1)^\circ$. The two C=Si bonds are nearly *gauche* to each other ($\angle C1-Si1-Si4-Si30 = -67.9^\circ$) to minimize steric repulsions between the large substituents. Overall **1** has the structure expected for a bis-silene in which two essentially noninteracting C=Si bonds are linked through a mercury atom.

The ^{13}C NMR chemical shift of the C=Si units (C1 and C30) appears at 225 ppm, slightly downfield from the corresponding signal of **5** (214 ppm^[11]). On the other hand, the ^{29}Si chemical shift of the double-bonded Si1 and Si4 atoms appears at 107.6 ppm and is strongly deshielded compared with the analogous signal of **5** (41.4 ppm^[11]). The difference (Δ) between $\delta(^{29}\text{Si})$ in **5** and in **1**, of -66.3 ppm, is similar in magnitude to the $\Delta\delta(^{29}\text{Si})$ value of $(Me_3Si)_4Si$ and $[(Me_3Si)_3Si]_2Hg$,^[12] (-80 ppm), indicating that Hg substitution has a similar effect on $\delta(^{29}\text{Si})$ regardless of whether the Si atom is single or double bonded.

Bis-silene **1** exhibits surprising high stability towards water, methanol, and acetone. Thus, **1** can be kept in these

[*] Dr. D. Bravo-Zhivotovskii, R. Dobrovetsky, D. Nemirovsky, V. Molev, G. Molev, Dr. M. Botoshansky, Prof. Y. Apeloig
Schulich Faculty of Chemistry and The Lise Meitner-Minerva Center for Computational Quantum Chemistry
Technion-Israel Institute of Technology, Haifa 32000 (Israel)
Fax: (+972) 4-829-4601
E-mail: chrbrzh@tx.technion.ac.il
chrapel@tx.technion.ac.il

Dr. M. Bendikov
Department of Organic Chemistry
Weizmann Institute of Science
Rehovot 76100 (Israel)

[**] This research was supported by the Israel Science Foundation, the Fund for the Promotion of Research at the Technion, and the Minerva Foundation in Munich. D.B.-Z. and Mark B. are grateful to the Ministry of Immigrant Absorption (Israel) for a Kamea scholarship.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.





Figure 1. Molecular structure of **1** in the crystal (ORTEP drawing). The hydrogen atoms have been omitted for clarity, and the thermal ellipsoids are set at the 20% probability level. Selected bond lengths [Å], bond angles [°], and bond dihedral angles [°]: Si1–C1 1.770(15), Si4–C30 1.758(15), Si1–Hg1 2.477(4), Si4–Hg1 2.417(4), Si1–Si2 2.337(6), Si4–Si5 2.401(6), C1–O1 1.384(16), C1–C2 1.553(19), C30–O2 1.390(17), C30–C31 1.521(2); Si4–Hg1–Si1 177.39(14), C1–Si1–Si2 127.3(5), C1–Si1–Hg1 113.0(5), Si2–Si1–Hg1 119.5(5), C30–Si4–Si5 124.5(6), C30–Si4–Hg1 116.6(5), Si5–Si4–Hg1 118.5(2), O1–C1–C2 110.3(11), O1–C1–Si1 121.5(10), C2–C1–Si1 128.0(10), O2–C30–C31 109.5(12), O2–C30–Si4 118.6(11), C31–C30–Si4 131.7(11); C1–Si1–Hg–Si4 –29.2(4), O2–C30–Si4–Hg1 –15.5(6), Si5–Si4–C30–O2 166.2(7), Si6–O2–C30–Si4 –65.4(9), O1–C1–Si1–Hg1 –17.3(8), Si2–Si1–C1–O1 165.9(6), Si3–O1–C1–Si1 –65.0(13), C1–Si1–Si4–C30 –67.9(4).

solvents for weeks without any noticeable transformation! However, when exposed to air, **1** reacts instantaneously yielding a complex mixture of products. The low reactivity of **1** probably results from effective steric protection of the Si=C bonds by the surrounding large *t*Pr₃Si and adamantyl groups, although electronic effect from the mercury substituent may also contribute. The low reactivity of sterically protected Si=C bond was reported previously, e.g., for (RR')Si=C=CR''₂ (R = 1-adamantyl, R' = 2,4,6-tri-*tert*-butylphenyl, CR''₂ = 1,3,6,8-tetraisopropyl-2,7-dimethoxy-9H-fluorenyl).^[13]


In an attempt to prepare more-reactive analogues of **1**, we synthesized bis(lithiosilyl)mercury **6**, an analogue of **2** carrying smaller *t*BuMe₂Si substituents. The coupling reaction of **6** with 1-adamantoyl chloride (**3**) yields the corresponding mercury bis-acylsilane **7** (Scheme 2), which was isolated




Scheme 2. Synthesis of **7**.

(87% yield) by crystallization from hexane at room temperature, and its structure was determined by X-ray crystallography.^[14]

Disappointingly, heating **7** to 200°C does not lead to rearrangement to the desired bis-silene, in contrast to **4** which rearranges spontaneously at room temperature. A possible explanation for the dramatic difference in the reactivity of **4** and **7** is the higher steric congestion in **4** (R₃Si = *i*Pr₃Si) relative to that in **7** (R₃Si = *t*BuMe₂Si). Consequently rearrangement of **4** to the corresponding silene is favored, but the less crowded **7** does not rearrange. DFT calculations at the B3LYP/6-31G* + ZPVE level of theory^[15] support this explanation. Thus, the rearrangement of **4** (R₃Si = *i*Pr₃Si) to the bis-silene **1** is endothermic by 1.5 kcal mol^{–1}. On the other hand, the rearrangement of **7** to **7a** is endothermic by 11.3 kcal mol^{–1}. With the smaller Me₃Si substituent the rearrangement **8** to **8a** is even more endothermic (15.8 kcal mol^{–1}) [Eq. (1)].



A similar steric enhancing effect is calculated also for the classic Brook rearrangement. The rearrangement of acylsilane **9** (R₃Si = Me₃Si) to silene **9a** is endothermic by 10.8 kcal mol^{–1}, decreasing to 5.2 kcal mol^{–1} for R₃Si = *t*BuMe₂Si substituents (**10**→**10a**) [Eq. (2)]. This insight is important for the synthesis of new silenes by means of the Brook rearrangement.



In conclusion, we have synthesized and characterized by X-ray crystallography the first bis-silene, a compound with two Si=C bonds, as well as the first metal-substituted silene, and have demonstrated that the size of the substituents in the precursor acylsilane controls the occurrence of its thermal rearrangement to a silene. We continue to explore the chemistry of **1** including its photochemistry (silyl mercurials are photosensitive yielding silyl radicals^[16]) which may lead to novel silicon compounds and reactive intermediates.

Received: February 28, 2008

Published online: April 30, 2008

Keywords: anions · lithium · rearrangements · silenes · structure elucidation

- [1] A. G. Brook, S. C. Nyburg, F. Abdesaken, B. Gutekunst, G. Gutekunst, R. Krishna, M. R. Kallury, Y. C. Poon, Y. M. Chang, W. N. Winnie, *J. Am. Chem. Soc.* **1982**, *104*, 5667.
- [2] R. West, M. J. Fink, J. Michl, *Science* **1981**, *214*, 1343.
- [3] For reviews see: a) R. West, *Polyhedron* **2002**, *21*, 467; b) R. West, *ACS Symp. Ser.* **2006**, *917*, 166.
- [4] For review see: N. Chinkov, H. Chechik, S. Majumdar, A. Liard, I. Marek, *Synthesis* **2002**, *17*, 2473.
- [5] a) D. Scheschkewitz, *Angew. Chem.* **2004**, *116*, 3025; *Angew. Chem. Int. Ed.* **2004**, *43*, 2965; b) M. Ichinohe, K. Sanuki, S. Inoue, A. Sekiguchi, *Organometallics* **2004**, *23*, 3088; c) S. Inoue, M. Ichinohe, A. Sekiguchi, *Chem. Lett.* **2005**, *34*, 1564; d) T.-L. Nguyen, D. Scheschkewitz, *J. Am. Chem. Soc.* **2005**, *127*, 10174; e) M. Driess, S. Block, M. Brym, M. T. Gamer, *Angew. Chem.* **2006**, *118*, 2351; *Angew. Chem. Int. Ed.* **2006**, *45*, 2293.
- [6] a) I. Bejan, D. Gueclue, S. Inoue, M. Ichinohe, A. Sekiguchi, D. Scheschkewitz, *Angew. Chem.* **2007**, *119*, 3413; *Angew. Chem. Int. Ed.* **2007**, *46*, 3349; b) S. Inoue, M. Ichinohe, A. Sekiguchi, *Angew. Chem.* **2007**, *119*, 3410; *Angew. Chem. Int. Ed.* **2007**, *46*, 3346; c) I. Bejan, D. Scheschkewitz, *Angew. Chem.* **2007**, *119*, 5885; *Angew. Chem. Int. Ed.* **2007**, *46*, 5783; d) S. Yao, S. Block, M. Brym, M. Driess, *Chem. Commun.* **2007**, 3844.
- [7] a) D. Bravo-Zhivotovskii, H. Peleg-Vasserman, M. Kosa, G. Molev, M. Botoshanskii, Y. Apeloig, *Angew. Chem.* **2004**, *116*, 763; *Angew. Chem. Int. Ed.* **2004**, *43*, 745; b) B. K. Campion, R. Heyn, T. D. Tilley, *J. Am. Chem. Soc.* **1990**, *112*, 4079.
- [8] A metal-substituted silene was proposed as a reactive intermediate but was not isolated, see: J. Ohshita, H. Sakurai, S. Masaoka, M. Tamai, A. Kunai, M. Ishikawa, *J. Organomet. Chem.* **2001**, *633*, 131.
- [9] D. Bravo-Zhivotovskii, M. Yuzefovich, N. Sigal, G. Korogodsky, K. Klinkhammer, B. Tumanskii, A. Shames, Y. Apeloig, *Angew. Chem.* **2002**, *114*, 671; *Angew. Chem. Int. Ed.* **2002**, *41*, 649.
- [10] Crystallographic data for **1**: $C_{58}H_{116}HgO_2Si_6$, $M_r = 1214.64$, monoclinic, space group $C2/c$, $a = 39.649(8)$, $b = 12.719(2)$, $c = 27.119(5)$ Å, $\beta = 109.37(3)^\circ$, $V = 12902(4)$ Å³, $Z = 8$, Nonius Kappa CCD, $Mo_{K\alpha}$ radiation (0.71073 Å), 220 K, $2\theta_{max} = 25^\circ$, $R = 0.0994$ ($I > 2\sigma I$), $wR2 = 0.2585$ ($I > 2\sigma I$), $Rw = 0.1709$ (all data), GOF = 0.847. CCDC 645497 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [11] K. Klinkhammer, J. Z. Weidlein, *Z. Anorg. Allg. Chem.* **1996**, *622*, 1209.
- [12] J. Arnold, T. D. Tilley, A. L. Rheingold, S. J. Geib, *Inorg. Chem.* **1987**, *26*, 2106.
- [13] G. E. Miracle, J. L. Ball, D. R. Powell, R. West, *J. Am. Chem. Soc.* **1993**, *115*, 11598.
- [14] For more information about the crystal structure of **7** see the Supporting Information.
- [15] Gaussian 03 was used. All molecules were fully optimized using a small core ECP on Hg; frequency calculations were performed at the same level. For full details see the Supporting Information.
- [16] a) D. Bravo-Zhivotovskii, I. Ruderfer, M. Yuzefovich, M. Kosa, M. Botoshansky, B. Tumanskii, Y. Apeloig, *Organometallics* **2005**, *24*, 2698; b) D. Bravo-Zhivotovskii, I. Ruderfer, S. Melamed, M. Botoshansky, B. Tumanskii, Y. Apeloig, *Angew. Chem.* **2005**, *117*, 749; *Angew. Chem. Int. Ed.* **2005**, *44*, 739.