


Carbocycle Synthesis through Facile and Efficient Palladium-Catalyzed Allylative De-aromatization of Naphthalene and Phenanthrene Allyl Chlorides**

Shirong Lu, Zhanwei Xu, Ming Bao,* and Yoshinori Yamamoto*


De-aromatization reactions of arenes have attracted considerable attention because they provide a simple way to synthesize functionalized alicyclic compounds, which can be used as intermediates for the preparation of natural products and bioactive compounds.^[1] Over the past four decades, many methods, including oxidation,^[2] reduction,^[3] photocycloaddition,^[4] [2,3] σ -rearrangement,^[5] electrophilic addition,^[6] nucleophilic addition,^[7] and other approaches^[8] have been developed for breaking up the conjugated π system. The complexation of aromatic system to transition metals leads to the activation of arenes and thus facilitates the electrophilic addition of $[M(\eta^2\text{-arene})]$ ($M = \text{Os, Re, Mo, and W}$) complexes and the nucleophilic addition of $[M(\eta^6\text{-arene})]$ ($M = \text{Cr, Mn, and Ru}$) complexes.

We recently reported the facile palladium-catalyzed allylative de-aromatization reaction of benzylic chlorides **1** with allyltributyltin.^[9] This process appears to involve the formation and isomerization of the $\eta^3\text{-allyl-}\eta^3\text{-benzylpalladium}$ intermediate **2** to give **2'**, which led to **3**, where an allyl group is linked *para* to the exocyclic methylene group [Eq. (1)]. Our interest in extending the scope of this de-



aromatization reaction led us to examine the cinnamyl chloride **4**. We assumed that, if the bis($\eta^3\text{-allyl}$)palladium

intermediate **5**, generated from the cinnamyl chlorides and allyltributyltin in the presence of a palladium catalyst, could undergo rearrangement to give the $\eta^3\text{-benzylpalladium}$ intermediate **5'**, and **6'** (or its *para* isomer) would be produced. However, only the Stille cross-coupling product **6** was obtained [Eq. (2)].^[10] Thus, extension of the de-aromatization to the cinnamyl chlorides **4** was not feasible.



Herein, we report the facile and efficient allylative de-aromatization of naphthalene and phenanthrene derivatives bearing an allyl chloride unit (Tables 1 and 2). These reactions did not form the corresponding Stille cross-coupling products, but pleasingly gave the *ortho* allylated product **8**; this regioselectivity is in marked contrast to the *para* selectivity of the de-aromatization of benzylic chlorides. This result introduces the possibility of synthesizing six-membered ring systems with three or four fused rings from naphthalenes or phenanthrenes, respectively.



















The allylative de-aromatization reactions of naphthalene derivatives **7a–i** with allyltributylstannane were performed in the presence of $[\text{Pd}_2(\text{dba})_3]$ (5 mol %) and PPh_3 (20 mol %; Table 1). The simple substrates **7a** and **7h** underwent the de-aromatization reaction smoothly to afford **8a** and **8h** in high yields (84 % and 78 %, respectively; Table 1, entries 1 and 8). Neither the electron-donating group nor the electron-withdrawing group on the aromatic ring exerted a strong influence on the reaction (except in terms of the reaction times). The yields of **8b–g** and **8i** were in the range of 74 to 89 % (Table 1, entries 2–7 and 9); a longer reaction time was needed for **7b** than for **7a** and **7c–i**. The lower reactivity of **7b** is perhaps due to steric hindrance from the β methyl group. The de-aromatization reactions of **7e** and **7f** were accelerated by the electron-withdrawing groups (Br or PhCOO , respectively) at the *para* position, and were completed in shorter reaction times compared with **7a–d** and **7g–i**. Products **8a** and **8c–i** were sensitive to acid, but were comparatively more stable than **3** derived from benzylic chlorides. Product **8b** with a quaternary carbon center was very robust, and could be purified by standard column chromatography on silica gel.

[*] S. Lu, Z. Xu, Prof. Dr. M. Bao
State Key Laboratory of Fine Chemicals
Dalian University of Technology, Dalian 116012 (China)
Fax: (+86) 411-8899-3687
E-mail: mingbao@dlut.edu.cn
Prof. Dr. Y. Yamamoto
Department of Chemistry, Graduate School of Science
Tohoku University, Sendai 980-8578 (Japan)
Fax: (+81) 22-795-6784
E-mail: yoshi@mail.tains.tohoku.ac.jp

[**] We are grateful to the National Natural Science Foundation of China (20572010) for financial support. This work was partly supported by the Program for Changjiang Scholars and Innovative Research Teams in Universities (IRT0711).

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


Table 1: Palladium-catalyzed de-aromatization reaction of naphthalene derivatives **7a–i** with allyltributylstannane.^[a]

Entry	Substrate	<i>t</i> [h]	Product	Yield [%] ^[b]
1		12		84
2		24		82
3		12		82
4		14		87
5		2		89
6		6		87
7		14		80
8		18		78
9		12		74


[a] A solution of naphthalene derivative **7** (0.5 mmol), allyltributylstannane (0.5 mmol), [Pd₂(dba)₃] (5 mol%), and Ph₃P (20 mol%) in dichloromethane (3 mL) was stirred at room temperature under N₂ for the period indicated. The reaction progress was monitored by TLC.
[b] Yields of isolated product. dba = *trans,trans*-dibenzylideneacetone.

Each of the new products was characterized through NMR^[11] and IR spectroscopy as well as HRMS.


It occurred to us that the two terminal olefinic groups of **8** might undergo a metathesis reaction to construct a new alicyclic ring. Indeed, when **8b** was treated with the ruthenium-carbene catalyst **9**,^[12] the cyclized product **10** was isolated in 62 % yield [Eq. (3)]. This result further confirmed



that the configuration of the 1,3-butadiene moiety of **8b** was identical to that presented in Table 1. The de-aromatization products were very stable under basic conditions, thus suggesting that **8f** could be hydrolyzed in an NaOH solution to produce a cyclic ketone derivative without the formation of any isomerization product. In fact, the desired product **11** was isolated in 77 % yield [Eq. (4)]; it seemed that the functionalized product **11** might be useful for further manipulation.



A plausible mechanism for the allylative de-aromatization reaction is shown in Scheme 1. The oxidative addition of **7a** to a Pd⁰ species would produce the η³-allylpalladium chloride intermediate **12**, which would react with allyltributylstannane to generate a bis(η³-allyl)palladium intermediate **13** upon ligand exchange. Isomerization of **13** would occur to give a bis(η³-allyl)palladium intermediate **13'**, which could undergo










Scheme 1. Proposed mechanism for the de-aromatization reaction.

reductive elimination to form the de-aromatization product **8a** and regenerate the Pd⁰ catalyst.

The successful extension of the allylative de-aromatization to the naphthalene derivatives **7a–i** encouraged us to examine the phenanthrene derivatives **7j–l**, and the results are summarized in Table 2. The simple substrate **7j** gave the

Table 2: Palladium-catalyzed de-aromatization reaction of phenanthrene derivatives **7j–l** with allyltributylstannane.^[a]



Entry	Substrate	t [h]	Product	Yield [%] ^[b]
1		12		87
2		9		84
3		21		86

[a] A solution of phenanthrene derivative **7** (0.5 mmol), allyltributylstannane (0.5 mmol), [Pd₂(dba)₃] (5 mol%), and Ph₃P (20 mol%) in dichloromethane (3 mL) was stirred at room temperature under N₂ for the period indicated. The reaction progress was monitored by TLC. [b] Yields of isolated product.

corresponding product **8j** in 87% yield (Table 2, entry 1). Even **7k**, with a C(sp³)-H β-hydrogen atom, underwent the reaction smoothly and afforded **8k** as the only product in 84% yield (Table 2, entry 2).^[13] The substrate **7l**, bearing an *n*Pr group α to the chlorine atom, also provided the de-aromatization product **8l** in 86% yield, without the formation of any β-hydride elimination product (Table 2, entry 3). The products **8j–l** are very stable, and each was purified by standard column chromatography on silica gel. No re-aromatized compounds were observed.

Even though the de-aromatization of the benzene derivatives **4** bearing allyl chloride was not successful, that of the naphthalene and phenanthrene derivatives **7** proceeded readily and efficiently. Perhaps, this difference is due to lower resonance energies of the aromatic rings of **7** compared to those of **4**.^[14] The facile and efficient construction of **8** and their rather unexpected stability allows the synthesis of extended fused and functionalized ring systems from naphthalenes and phenanthrenes.

Experimental Section

General procedure for the allylative de-aromatization reaction: Allyltributylstannane (165.6 mg, 0.5 mmol) and **7a** (101.3 mg, 0.5 mmol) were added to a solution of [Pd₂(dba)₃] (22.8 mg, 0.025 mmol) and PPh₃ (26.2 mg, 0.1 mmol) in dichloromethane (3 mL) at room temperature, and the reaction mixture was stirred for 12 h under N₂. After the allyltributylstannane was consumed (as evident by TLC), the solvent was removed under reduced pressure. The residue was filtered through a short column of basic alumina to remove palladium by-products and eluted with pentane to give **8a** in 84% yield (87.5 mg) as a colorless liquid.


Received: February 1, 2008

Published online: April 29, 2008

Keywords: de-aromatization · naphthalenes · palladium · phenanthrenes · synthetic methods

- [1] For recent selected references, see: a) F. López Ortiz, M. J. Iglesias, I. Fernández, C. M. Andújar Sánchez, G. Ruiz Gómez, *Chem. Rev.* **2007**, *107*, 1580–1691; b) J. Gagnepain, F. Castet, S. Quideau, *Angew. Chem.* **2007**, *119*, 1555–1557; *Angew. Chem. Int. Ed.* **2007**, *46*, 1533–1535; c) E. N. Pitsinos, V. I. Moutsos, O. Vageli, *Tetrahedron Lett.* **2007**, *48*, 1523–1526; d) N. Lebrasseur, J. Gagnepain, A. Ozanne-Beaudenon, J.-M. Léger, S. Quideau, *J. Org. Chem.* **2007**, *72*, 6280–6283; e) L. H. Mejorado, T. R. R. Pettus, *J. Am. Chem. Soc.* **2006**, *128*, 15625–15631; f) D. Crich, V. Krishnamurthy, *Tetrahedron* **2006**, *62*, 6830–6840; g) J. Zhu, N. P. Grigoriadis, J. P. Lee, J. A. Porco, Jr., *J. Am. Chem. Soc.* **2005**, *127*, 9342–9343.
- [2] For the oxidation of phenol and its derivatives, see: a) P. J. Stang, V. V. Zhdankin, *Chem. Rev.* **1996**, *96*, 1123–1178; b) N. Lebrasseur, G.-J. Fan, M. Oxoby, M. A. Looney, S. Quideau, *Tetrahedron* **2005**, *61*, 1551–1562; c) E. N. Pitsinos, A. Cruz, *Org. Lett.* **2005**, *7*, 2245–2248; d) S. Quideau, L. Pouységu, A. Ozanne, J. Gagnepain, *Molecules* **2005**, *10*, 201–216; e) L. H. Mejorado, C. Hoarau, T. R. R. Pettus, *Org. Lett.* **2004**, *6*, 1535–1538; f) R. W. Van De Water, C. Hoarau, T. R. R. Pettus, *Tetrahedron Lett.* **2003**, *44*, 5109–5113; g) S. Mandal, D. Macikenas, J. D. Protasiewicz, L. M. Sayre, *J. Org. Chem.* **2000**, *65*, 4804–4809; h) J. S. Swenton, K. Carpenter, Y. Chen, M. L. Kerns, G. W. Morrow, *J. Org. Chem.* **1993**, *58*, 3308–3316; also see Ref. [1g]; for microbial oxidation, see: i) V. Bui, T. V. Hansen, Y. Stenström, D. W. Ribbons, T. Hudlicky, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1669–1672; j) D. R. Boyd, N. D. Sharma, N. I. Bowers, J. Duffy, J. S. Harrison, H. Dalton, *J. Chem. Soc. Perkin Trans. 1* **2000**, 1345–1350; k) D. R. Boyd, N. D. Sharma, S. A. Barr, H. Dalton, J. Chima, G. Whited, R. Seemayer, *J. Am. Chem. Soc.* **1994**, *116*, 1147–1148.
- [3] For catalytic hydrogenation, see: a) E. Heller, W. Lautenschläger, U. Holzgrabe, *Tetrahedron Lett.* **2005**, *46*, 1247–1249; b) T. J. Donohoe, R. Garg, C. A. Stevenson, *Tetrahedron: Asymmetry* **1996**, *7*, 317–344; for Birch reduction, see: c) A. J. Birch, *Pure Appl. Chem.* **1996**, *68*, 553–556.
- [4] a) For a review, see: J. Cornélisse, *Chem. Rev.* **1993**, *93*, 615–669; b) S. Kohmoto, H. Masu, C. Tatsuno, K. Kishikawa, M. Yamamoto, K. Yamaguchi, *J. Chem. Soc. Perkin Trans. 1* **2000**, 4464–4468, and references therein; c) A. Yokoyama, K. Mizuno, *Org. Lett.* **2000**, *2*, 3457–3459, and references therein.
- [5] For nitrogen-ylide rearrangement, see: a) N. Shirai, Y. Watanabe, Y. Sato, *J. Org. Chem.* **1990**, *55*, 2767–2770, and references therein; for sulfur-ylide rearrangement, see: b) C. C. McComas, D. L. Van Vranken, *Tetrahedron Lett.* **2003**, *44*, 8203–8205, and references therein; c) M. G. Burdon, J. G. Moffatt, *J. Am. Chem. Soc.* **1967**, *89*, 4725–4735; d) C. R. Hauser, S. W. Kantor, W. R.

- Brasen, *J. Am. Chem. Soc.* **1953**, 75, 2660–2663; for Still–Wittig rearrangement, see: e) P. Monje, P. Grana, M. R. Paleo, F. J. Sardina, *Org. Lett.* **2006**, 8, 951–954; f) P. A. Caruana, A. J. Frontier, *Tetrahedron* **2004**, 60, 10921–10926.
- [6] For reviews, see: a) J. M. Keane, W. D. Harman, *Organometallics* **2005**, 24, 1786–1798; b) W. D. Harman, *Coord. Chem. Rev.* **2004**, 248, 853–866; c) A. R. Pape, K. P. Kaliappan, E. P. Kündig, *Chem. Rev.* **2000**, 100, 2917–2940; also see: d) D. A. Delafuente, W. H. Myers, M. Sabat, W. D. Harman, *Organometallics* **2005**, 24, 1876–1885.
- [7] Arenes were activated by electron-withdrawing groups, see: a) A. M. Ramallal, I. Fernández, F. López Ortiz, J. González, *Chem. Eur. J.* **2005**, 11, 3022–3031, and references therein; b) Z. Wang, Z. Xi, *Synlett* **2006**, 1275–1277; c) J. Clayden, F. E. Knowles, I. R. Baldwin, *J. Am. Chem. Soc.* **2005**, 127, 2412–2413; d) J. Clayden, R. Turnbull, M. Helliwell, I. Pinto, *Chem. Commun.* **2004**, 2430–2431; e) J. Clayden, R. Turnbull, I. Pinto, *Org. Lett.* **2004**, 6, 609–611; f) J. Clayden, M. N. Kenworthy, M. Helliwell, *Org. Lett.* **2003**, 5, 831–834; g) J. Barluenga, S. K. Nandy, Y. R. S. Laxmi, J. R. Suárez, I. Merino, J. Flórez, S. García-Granda, J. Montejó-Bernardo, *Chem. Eur. J.* **2003**, 9, 5725–5736; arenes can be activated by complexation to transition metals, see: h) L. Zhou, L.-Z. Wu, L.-P. Zhang, C.-H. Tung, *Organometallics* **2006**, 25, 1707–1711; i) E. P. Kündig, A. Bellido, K. P. Kaliappan, A. R. Pape, S. Radix, *Org. Biomol. Chem.* **2006**, 4, 342–351; j) F. C. Pigge, J. J. Coniglio, R. Dalvi, *J. Am. Chem. Soc.* **2006**, 128, 3498–3499; k) F. C. Pigge, J. J. Coniglio, N. P. Rath, *Org. Lett.* **2003**, 5, 2011–2014; also see Ref. [6c].
- [8] For organolanthanide-induced de-aromatization, see: a) R. Liu, C. Zhang, Z. Zhu, J. Luo, X. Zhou, L. Weng, *Chem. Eur. J.* **2006**, 12, 6940–6952; for radical addition, see: b) D. Crich, M. Sannigrahi, *Tetrahedron* **2002**, 58, 3319–3322; c) J. Boivin, M. Yousfi, S. Z. Zard, *Tetrahedron Lett.* **1997**, 38, 5985–5988; for a de-aromatization reaction involving naphthylborate anions, see: d) E. Negishi, R. E. Merrill, *J. Chem. Soc. Chem. Commun.* **1974**, 860–861.
- [9] M. Bao, H. Nakamura, Y. Yamamoto, *J. Am. Chem. Soc.* **2001**, 123, 759–760.
- [10] H. Nakamura, M. Bao, Y. Yamamoto, *Angew. Chem.* **2001**, 113, 3308–3310; *Angew. Chem. Int. Ed.* **2001**, 40, 3208–3210.
- [11] The structural assignment of product **8** was based on 1D (^1H and ^{13}C) and 2D (^1H – ^1H COSY and NOESY) spectroscopic analysis. For example, the connection and configuration of products **8c** and **8h** were obtained from ^1H – ^1H COSY and NOESY spectra.



- [12] W. Zhang, C. Bai, X. Lu, R. He, *J. Organomet. Chem.* **2007**, 692, 3563–3567.
- [13] In the Stille cross-coupling reaction of $\text{C}(\text{sp}^3)\text{-X}$ electrophiles, the β hydride elimination often gave alkenes in preference to coupling products, see: a) D. J. Cárdenas, *Angew. Chem.* **2003**, 115, 398–401; *Angew. Chem. Int. Ed.* **2003**, 42, 384–387; b) T.-Y. Luh, M.-k. Leung, K.-T. Wong, *Chem. Rev.* **2000**, 100, 3187–3204.
- [14] DFT computation of intermediates and products are now under investigation, and a detailed mechanistic discussion will appear in due course.