


Hybrid Nanoparticles

S. Guo, J. Zhai, Y. Fang, S. Dong,
E. Wang*

Nanoelectrocatalyst Based on
High-Density Au/Pt Hybrid
Nanoparticles Supported on a Silica
Nanosphere

Chem. Asian J.
DOI: 10.1002/asia.200700422

Good things come in small packages: A high-efficiency nanoelectrocatalyst based on high-density Au/Pt hybrid nanoparticles supported on a silica nanosphere (Au-Pt/SiO₂) can be prepared easily by a wet chemical process. This Au-Pt/SiO₂ nanostructure exhibits a high electrocatalytic activity for oxygen reduction and methanol oxidation, making it an interesting candidate for application in fuel cells. APTMS = (3-aminopropyl)trimethoxysilane.




Combinatorial Libraries

S. Mitra, A. M. Barrios*

Identifying Selective Protein Tyrosine
Phosphatase Substrates and Inhibitors
from a Fluorogenic, Combinatorial
Peptide Library

ChemBioChem
DOI: 10.1002/cbic.200800046



Protein tyrosine phosphatases are increasingly recognized as enzymes that exhibit exquisite substrate selectivity with important roles in cellular signaling, and have been identified as attractive therapeutic targets in human diseases includ-


ing autoimmunity, obesity, diabetes, and cancer. A new approach was developed to rapidly and efficiently profile the substrate selectivity of protein tyrosine phosphatase and is described herein.

DNA Dimers

E. Y. M. Bonnist, A. C. Jones*

Long-Wavelength Fluorescence from
2-Aminopurine-Nucleobase Dimers in
DNA

ChemPhysChem
DOI: 10.1002/cphc.200700813



Illuminating interactions: A fluorescent analogue of adenine illuminates inter-base interactions in DNA. Long-wavelength spectra arise from the formation

of a ground-state heterodimer with an adjacent π -stacked, natural base, in addition to the familiar short-wavelength spectra (see figure).


Imaging Agents


H. W. Chen, C. D. Medley, K. Sefah,
D. Shangguan, Z. Tang, L. Meng,
J. E. Smith, W. Tan*

Molecular Recognition of Small-Cell
Lung Cancer Cells Using Aptamers


ChemMedChem
DOI: 10.1002/cmdc.200800030

Early diagnosis is the key for lung cancer survival. Novel aptamer-based molecular probes were developed for the recognition of specific small-cell lung cancer (SCLC) cell-surface molecular markers. They show high affinity and specificity in various assay formats. This approach shows the potential for early lung cancer detection.





a bent imido



a linear imido

Structural characterisation of (arylimido)-triethanolaminovanadium(V) compounds was carried out by single-crystal X-ray structure determinations to elucidate the substituent effect on the imido structures in which the imido structures are strongly controlled through π conjugation by the *para* substituents of the aryl moieties.


(Arylimido)vanadium(V) Species

T. Moriuchi,* T. Beppu, K. Ishino, M. Nishina, T. Hirao*

Structural Control of (Arylimido)vanadium(V) Compounds through π Conjugation

Eur. J. Inorg. Chem.

DOI: 10.1002/ejic.200701252



Oxidative homocoupling of lithiated 2-substituted indenenes furnishes *racemic* dimers, while radical reaction leads to a mixture of *racemic* and *meso* diastereomers. Internal rotation in these dimers was studied by NMR and molecular modelling.


Triptycenes

K. Nikitin,* H. Müller-Bunz, Y. Ortin, W. Risse, M. J. McGlinchey*

Twin Triptycyl Spinning Tops: A Simple Case of Molecular Gearing with Dynamic C_2 Symmetry

Eur. J. Org. Chem.

DOI: 10.1002/ejoc.200800202



Novel catalytic features: This paper describes the development of a new unsymmetrical PCS-pincer palladium complex (see scheme) for catalytic aldol reactions and coupling of allyl chlorides and vinyloxiranes with aldehyde and imine electrophiles. The synergistic electronic effects of the phosphorus and

sulfur side-arms of this pincer complex generate novel catalytic features, one of the most interesting of which was tandem catalytic activity found for the coupling reactions of allyl chlorides with electrophiles in the presence of hexamethylditin.

Pincer Complexes


M. Gagliardo, N. Selander, N. C. Mehendale, G. van Koten, R. J. M. Klein Gebbink,* K. J. Szabó*

Catalytic Performance of Symmetrical and Unsymmetrical Sulfur-Containing Pincer Complexes: Synthesis and Tandem Catalytic Activity of the First PCS-Pincer Palladium Complex

Chem. Eur. J.

DOI: 10.1002/chem.200800350

Scratching the surface: The reactions of oxygenates such as glycerol are important for the production of H_2 . Temperature-programmed desorption experiments have revealed an increased production of H_2 on the Ni surface monolayer on Pt(111) (Ni-Pt-Pt(111)). Glycerol reforming activity trends are similar to previous results for ethylene glycol and ethanol, demonstrating that smaller oxygenates can be used as good models for reforming of larger, biomass-derived oxygenates.



Glycerol Conversion

O. Skoplyak, M. A. Barteau,* J. G. Chen

Enhancing H_2 and CO Production from Glycerol Using Bimetallic Surfaces

ChemSusChem

DOI: 10.1002/cssc.200800053