

Living Radical Polymerization


M. Ouchi,* M. Ito, S. Kamemoto,
M. Sawamoto*

Highly Active and Removable Ruthenium Catalysts for Transition Metal-Catalyzed Living Radical Polymerization: Design of Ligands and Cocatalysts

Chem. Asian J.

DOI: 10.1002/asia.200800142

Slim and Active. The systematic search and design of phosphine ligands (PR_3) and amine cocatalysts (additives) resulted in obtaining highly active and removable catalysts, pentamethyl-cyclopentadienyl (Cp^*) ruthenium(II) phosphine complexes $[\text{RuCp}^*\text{Cl}(\text{PR}_3)_2]$, for the transition metal-catalyzed living radical polymerization of methyl methacrylate (MMA).



Oligosaccharides


M. Vasan, J. Rauvolfova, M. A. Wolfert,
C. Leoff, E. L. Kannenberg, C. P. Quinn,
R. W. Carlson,* G.-J. Boons*

Chemical Synthesis and Immunological Properties of Oligosaccharides Derived from the Vegetative Cell Wall of *Bacillus anthracis*

ChemBioChem

DOI: 10.1002/cbic.200800210


Sweet relief: Sera from rabbits exposed either to live and irradiation-killed spores of *Bacillus anthracis* Sterne 34F₂ or immunized with *B. anthracis* polysaccharide conjugated to keyhole limpet hemocyanin (KLH) were found to contain antibodies that recognized isolated polysaccharide (shown in scheme) and two synthetic trisaccharides. This provides proof-of-concept towards the development of vegetative and spore-specific reagents for detection and targeting of nonprotein structures of *B. anthracis*.



Ionic Liquids

Y. Gao, A. Voigt,* L. Hilfert,
K. Sundmacher

Nanodroplet Cluster Formation in Ionic Liquid Microemulsions



Drop by drop: A unique self-assembled cluster nanostructure is observed in reverse bmimBF₄-in-toluene ionic liquid microemulsions (see picture). The large nanodroplet clusters consist of small

ionic liquid microemulsion droplets. These novel nanodroplet clusters may have some unusual and unique properties with a number of interesting possibilities for potential applications.

ChemPhysChem

DOI: 10.1002/cphc.200800157


Antitumor Agents

K. V. Kong, W. K. Leong,* S. P. Ng,
T. H. Nguyen, L. H. K. Lim


Osmium Carbonyl Clusters: A New Class of Apoptosis Inducing Agents

ChemMedChem

DOI: 10.1002/cmdc.200800069



Putting the 'Os' in apoptosis: Osmium carbonyl clusters were found to induce apoptosis in four cancer cell lines, namely, ER+ breast carcinoma (MCF-7), ER- breast carcinoma (MDA-MB-231), metastatic colorectal adenocarcinoma (SW620), and hepatocarcinoma (Hep G2). The metal clusters are more cytotoxic towards these cancer cells than they are towards normal epithelial cells.



Photophysical properties and surface structures of silica-gel-supported photo-functional (phthalocyaninato)silicon complexes (SiPc-SiO_2) have been investigated. By changing the amount of SiPcs, we have succeeded in preparing solid materials based on Pc complexes having both monomeric photophysical properties and a high ability for singlet-oxygen generation.


Photoactive-Solid Phthalocyanines

K. Ishii,* Y. Kikukawa, M. Shiine, N. Kobayashi,* T. Tsuru, Y. Sakai, A. Sakoda

Synthesis and Photophysical Properties of Silica-Gel-Supported Photofunctional (Phthalocyaninato)silicon Complexes

Eur. J. Inorg. Chem.

DOI: 10.1002/ejic.200800208



Discotic liquid crystalline π -acceptor compounds were synthesized by using *anti*- and *syn*-pyrenediones. The liquid crystalline phases were assigned to be D_{L2} from POM, DSC, and XRD measurements and single-crystal study of the model crystalline compounds.


Liquid Crystals

M. Yasutake, T. Fujihara, A. Nagasawa, K. Moriya, T. Hirose*

Synthesis and Phase Structures of Novel π -Acceptor Discotic Liquid Crystalline Compounds Having a Pyrenedione Core

Eur. J. Org. Chem.

DOI: 10.1002/ejoc.200800360



Two electrons, three new stereocentres: The *cis*-hydrindane motif is found in a number of natural products that display important biological activity. A flexible, stereoselective approach to the framework has been developed that features

highly diastereoselective, SmI_2 -mediated cyclisations. The strategy has been exploited in the first synthesis of the proposed structure of faurinone, a sesquiterpene ketone isolated from *Valeriana officinalis*.

Natural Product Synthesis


T. J. K. Findley, D. Sucunza, L. C. Miller, D. T. Davies, D. J. Procter*

A Flexible, Stereoselective Approach to the Decorated *cis*-Hydrindane Skeleton: Synthesis of the Proposed Structure of Faurinone

Chem. Eur. J.

DOI: 10.1002/chem.200800930

It's not easy being green! New value chains and catalytic systems must be developed to decrease the cost of biomass processing to bioproducts as those employed for hydrocarbons are not adapted to biomolecules. However, the extensive use of biomass for industrial production raises environmental and ethical issues, which in turn raise doubts on the sustainability of these processes.



Biomass Conversion

P. Gallezot*

Catalytic Conversion of Biomass: Challenges and Issues

ChemSusChem

DOI: 10.1002/cssc.200800091



Auf diesen beiden Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie die

Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley InterScience leicht online zugänglich.