

Reply**

Daniel Himmel, Nils Trapp, Ingo Krossing,* Sandra Altmannshofer, Verena Herz, Georg Eickerling, and Wolfgang Scherer*

ab initio calculations · alkynes · silver ·
structure elucidation · weakly coordinating anions

We acknowledge the comments of Krapp and Frenking (KF)^[1] on our recent publication “Homoleptic Silver(I) Acetylene Complexes”^[2] and respond to the issues they raised: KF criticize the use of the term “pseudo-gas phase conditions” as misleading, discussing the difference between solid-state structures and gas-phase structures in HCN–BF₃ and a “high level” (i.e. MP2/def2-SVP)^[3] calculated gas-phase structure of pentaphenylborole^[4] as examples that intermolecular interactions cannot be neglected. Indeed, the term pseudo-gas phase conditions was only defined for cations in salts with large weakly coordinating anions (WCAs).^[5] Such

anions separate the cations under investigation considerably and the large, poorly polarizable fluorinated “Teflon”-surface of the anions effectively eliminates interactions between them. Clearly, directed interactions, such as hydrogen bonds and π – π interactions, have a strong influence on the solid-state structures and distort them from the gas-phase geometry. In this respect we agree with KF that there are no pseudo-gas phase conditions in the solid-state structures of HCN–BF₃ and the mentioned borole. However, as detailed analyses of the solid-state structures showed, there are almost no such interactions in the compounds discussed in our publication.^[2]

We generally compare the environment provided by salts of good WCAs with the gas-phase conditions inside a mass spectrometer. In agreement with this, we were able to synthesize and characterize compounds in condensed phase that were initially investigated by advanced mass spectrometry (MS) methods, for example, [Ag(P₄)]⁺, [Ag(P₄)₂]⁺,^[6–8] [Ag(S₈)]⁺, [Ag(S₈)₂]⁺,^[9,10] [Cu(CO)_n]⁺ ($n = 1–4$),^[11] and Cl₃⁺.^[12–14] In part we were even able to characterize species prior to MS (cf. [Ag(C₂H₄)]⁺ and [Ag(C₂H₄)₂]⁺ were identified by MS in 1991,^[15] but no evidence for [Ag(C₂H₄)₃]⁺ was found until 2005;^[16] yet [Ag(C₂H₄)₃]⁺[Al(OC(CF₃)₃)₄][–] could be prepared in 2003^[17]).

The thermodynamic principles of pseudo-gas phase conditions follow from the Born–Haber cycle for the formation of [AgL_x]⁺[A][–] ($x = 3, 4$) from Ag⁺[A][–] and L (L = C₂H₂) for the two differently sized anions [A][–] = [BF₄][–] ($V_{\text{ion}} = 0.077 \text{ nm}^3$) and the ten-times larger [Al(OC(CF₃)₃)₄][–] ($V_{\text{ion}} = 0.758 \text{ nm}^3$; Figure 1).^[18]

Using volume-based thermodynamics developed by Jenkins and Glasser^[20,21] and the molecular volumes of [Ag(C₂H₂)₃]⁺[Al(OR^F)₄][–] (**1**) and [Ag(C₂H₂)₄]⁺[Al(OR^F)₄][–] (**2**) derived from crystal data (OR^F = OC(CF₃)₃), the loss of Gibbs lattice energy upon addition of three C₂H₂ molecules ($\Delta\Delta G_{\text{latt}}$) was estimated as only -48 kJ mol^{-1} (and -15 kJ mol^{-1} for the fourth C₂H₂ molecule) in case of the large aluminate anion, but as -173 kJ mol^{-1} (for three) and -46 kJ mol^{-1} (for the fourth) for the tetrafluoroborate (details deposited in the Supporting Information). Only for the large aluminate, the Gibbs energies for the complexation of three or four C₂H₂ molecules are comparable in solid state and gas phase (gas: -260 and -272 kJ mol^{-1} vs. solid: -212 and -209 kJ mol^{-1}). The smaller [BF₄][–] ion is much less suited to provide pseudo-gas phase conditions; the Gibbs energies for the solid state are greatly reduced to -86 and -55 kJ mol^{-1} and show no resemblance to the gas-phase values. According to the calculated cycle, the gas-phase equilibrium between [Ag(C₂H₂)₄]⁺, [Ag(C₂H₂)₃]⁺ and C₂H₂ can easily be calculated [Eq. (1)]; $p(\text{X})$ is the partial pressure of X; $a(\text{X})$ = activity of X = $p(\text{X})/(1 \text{ atm})$ for an ideal gas].

$$K_{(\text{g})} = \frac{a(\text{AgL}_4^+)}{a(\text{AgL}_3^+)a(\text{L})} = \frac{p(\text{AgL}_4^+) 1 \text{ atm}}{p(\text{AgL}_3^+)p(\text{L})} = e^{-\Delta G_{(\text{g})}^0/RT} = 126 \quad (1)$$

with $\Delta G_{(\text{g})}^0 = -12 \text{ kJ mol}^{-1}$ at 298.15 K

Also, the acetylene pressure of solid **2** can be obtained from the equilibrium constant derived from the solid state Gibbs reaction energy ($\Delta G_{(\text{s})}^0$) [Eq. (2)].

[*] Dr. D. Himmel, Dipl.-Chem. N. Trapp, Prof. Dr. I. Krossing
Institut für Anorganische und Analytische Chemie, Albert-Ludwigs-Universität Freiburg, Albertstrasse 21, 79104 Freiburg (Germany)
E-mail: krossing@uni-freiburg.de
Dipl.-Chem. S. Altmannshofer, Dipl.-Phys. V. Herz, Dr. G. Eickerling, Prof. Dr. W. Scherer
Institut für Physik, Universität Augsburg, Universitätsstrasse 1, 86159 Augsburg (Germany)
E-mail: wolfgang.scherer@physik.uni-augsburg.de

[**] This work was supported by Swiss National Fond, the EPFL, the Albert-Ludwigs Universität Freiburg, the Universität Augsburg, the Fond der Chemischen Industrie, the Deutsche Forschungsgemeinschaft (SPP1178 and Normalverfahren) as well as NanoCat, an International Graduate Program within the Elitenetzwerk Bayern. We thank Krapp and Frenking for their suggestions, hope we could address their issues and that this statement helps to gain a deeper insight into our work.



Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200802616>.




Figure 1. Born-Haber cycle with ΔG^0 values [kJ mol^{-1}] at 298.15 K. Gas-phase reaction energy calculated at MP2(FC)/TZVPP, thermal and entropic corrections at the BP86/SV(P) level (ref. [19]). For solid-state reactions and lattice energies, upper values (red) count for $[\text{A}]^- = [\text{BF}_4]^-$, lower values (blue) for $[\text{Al}(\text{OC}(\text{CF}_3)_3)_4]^-$.

$$K_{\text{(s)}} = \frac{a([\text{AgL}_4]^+[\text{A}]^-(\text{s}))}{a([\text{AgL}_3]^+[\text{A}]^-(\text{s}))a(\text{L}_{\text{(g)}})} = \frac{1}{1a(\text{L}_{\text{(g)}})}$$

$$= \frac{1 \text{ atm}}{p(\text{L})} = e^{-\Delta G^0_{\text{(s)}}/RT} = 0.29$$

$$p(\text{L}) = \frac{1 \text{ atm}}{K_{\text{s}}} = 3.4 \text{ atm}$$

with $\Delta G^0_{\text{(s)}} = +3 \text{ kJ mol}^{-1}$ at 298.15 K
for $[\text{A}]^- = [\text{Al}(\text{OR}^{\text{F}})_4]^-$

The calculated acetylene pressure of 3.4 atm for **2** at 25°C correlates quite well with the observed decomposition above −20°C in an atmosphere of acetylene and the need for handling the compounds for sample preparation at temperatures between −50 and −100°C.

KF calculated the free complexation energy of the fourth acetylene ligand as −11.9 kJ mol^{-1} at −20°C. While this reproduces our own results quite accurately,^[18] we disagree with the interpretation: “This means that free $[\text{Ag}(\eta^2\text{-C}_2\text{H}_2)_4]^+$ hardly exists in the gas phase”. As the *negative* value implies, the reaction is *exergonic* at standard pressures. Using KF’s value, the gas-phase equilibrium constant $K_{\text{(g)}}$ [see Eq. (1)] is 285 at −20°C and the ratio between the trisethyne and the tetraethyne complex can be calculated with Equation (3) as 285:1 in C_2H_2 at 1 atm and still 1:1 at 3.5 mbar acetylene pressure.

$$\frac{p([\text{Ag}(\text{C}_2\text{H}_2)_4]^+)}{p([\text{Ag}(\text{C}_2\text{H}_2)_3]^+)} = K_{\text{(g)}} \frac{p(\text{C}_2\text{H}_2)}{1 \text{ atm}} \quad (3)$$

KF also raise the question whether the experimentally accessible $[\text{Ag}(\text{C}_2\text{H}_2)]^+[\text{A}]^-$ (**3**) model complex^[2]

($[\text{A}]^- = [\text{Al}(\text{OC}(\text{CH}_3)_3(\text{CF}_3)_2)_4]^-$) represents a suitable system to analyze the nature of the acetylene coordination at silver(I) cations under pseudo-gas phase conditions. In fact, we never used the term pseudo-gas phase conditions for **3** and described it explicitly as a molecular model compound in which the anion is clearly coordinating.

KF found on the basis of their own DFT calculations that the free $[\text{Ag}(\text{C}_2\text{H}_2)]^+$ ion displays by 6 pm longer Ag–C bond lengths relative to the corresponding ones calculated in our model system $[\text{Ag}(\text{C}_2\text{H}_2)]^+[\text{A}]^-$ (**3**). They describe the shortening of the Ag–C bond in **3** as “influence of a single counterion”. However, the same shortening is calculated^[22] in the case of coordinating two neutral water molecules giving $[\text{Ag}(\text{H}_2\text{O})_2(\text{C}_2\text{H}_2)]^+$.

Furthermore, KF criticize the agreement between the B3LYP structure calculation of $[\text{Ag}(\text{C}_2\text{H}_2)]^+[\text{A}]^-$ (**3**) and the experimental values as a fortuitous error cancellation. This may well be the case. However, this is a general phenomenon and it is very questionable if intermolecular interactions really contribute to this error cancellation as suggested by KF. In fact, we examined several density functionals and basis sets to find the method that reproduces our experimental results as accurately as possible.^[2] Indeed, a broad variation of Ag–C bond lengths was found (221–251 pm), which scatter around the experimental values of 230.7(1) and 230.9(1) pm. The agreement of the “best” method (B3LYP/def2-TZVPP)

showing the smallest discrepancies of less than 1 % with regard to the Ag–C, Ag–O, and C–C bond lengths in comparison with the multipolar model, may well be fortuitous. This only accentuates the need for precise structural data from high-resolution X-ray diffraction.

This brings us to the next point—the interpretation of the electronic structure displayed by silver(I) acetylene complexes. According to KF, “the silver-acetylene bond in **3** should be considered as electrostatic”. This is a rather strange remark in the light of recent statements published by Nechaev, Rayon, and Frenking (NRF) in 2004: “the nature of the metal–ligand interactions is not very different from each other in the donor–acceptor complexes $[(\text{CO})_5\text{TM}-\text{C}_2\text{H}_x]$ (TM=Cr, Mo, W), $[(\text{CO})_4\text{TM}-\text{C}_2\text{H}_x]$ (TM=Fe, Ru, Os), and $[\text{TM}^+-\text{C}_2\text{H}_x]$ (TM=Cu, Ag, Au)”.^[23] Furthermore, NRF stress that “the covalent contributions play an important role” which are characterized by computed stabilizing orbital interaction terms of 41.7 % and 40.6 % (of the attractive interactions) for $[\text{Ag}(\text{C}_2\text{H}_4)]^+$ and $[\text{Ag}(\text{C}_2\text{H}_2)]^+$, respectively.^[23] Consequently, NRF revised earlier work of Böhme, Wagener, and Frenking (BWF) in 1996^[24a] and of other groups^[24b,c] in which “the bonding situation of $[(\text{C}_2\text{H}_4)\text{TM}]^+$ and $[(\text{C}_2\text{H}_2)\text{TM}]^+$ complexes suggested that the interaction is *mainly electrostatic*”.^[23] It remains unclear which scientific facts led Frenking to change his opinion twice between 1996 and 2008. Indeed, Frenking’s present arguments are actually the ones

published earlier in 1996 (BWF) which in turn have been criticized by Hertwig et al.^[25] in the same year. However, we will show again in the following, that KF's line of argumentation is misleading and in conflict with their own published results and the present literature.

According to KF's calculations the electrostatic nature of the silver–acetylene bond in **3** is reflected by 1) the T-shape of the bond paths, 2) the positive value of the Laplacian ($\nabla^2\rho(\mathbf{r}) = 5.51 \text{ e } \text{\AA}^{-5}$), and the small negative value of the energy density ($H(\mathbf{r}) = -0.06 \text{ Hartree } \text{\AA}^{-3}$) at the TCP. The TCP represents the bond critical point of the bond path between Ag and the midpoint of the C≡C bond. We note that the same argumentation has been used 1996 by BWF to classify the bonding in $[\text{Cu}(\text{C}_2\text{H}_x)]^+$ ($x=2, 4$) as electrostatic “Coulombic interactions”. However, on the basis of a personal communication with Frenking, Hertwig et al. could demonstrate in the same year that “increasing the basis set on the ligand to TZP quality, however, also led to a ring structure”.^[25] Hence, subtle changes in the level of approximation employed might render a T-shape structure of transition-metal π complexes into a metallacyclic one and vice versa. Accordingly, the finding of a T-shape bond path does not rule out the presence of covalency in these type of π complexes. As a natural consequence of both σ donor and π back bonding in these transition-metal π complexes, only small charge-density differences are typically present between the ring and bond critical points in d^{10} configured $[\text{M}(\text{C}_2\text{H}_2)]^{(+)}$ π complexes ($\text{M} = \text{Group 10 or 11 metal}$) which render the bond-path graphs rather flexible (see Supporting Information and ref. ^[26a]). These findings are in line with a recent experimental charge-density work on transition-metal acetylides,^[27] where both cyclic and T-shape graphs for ScC_2 moieties were found to co-exist in the solid despite the lack of any significant geometrical differences in the individual ScC_2 moieties.

Also the next two arguments (positive Laplacian and slightly negative $H(\mathbf{r})$ values at the TCP) used by Frenking are misleading. Why should a positive sign of the Laplacian rule out the presence of covalent silver(I) acetylene interactions? KF should more carefully

reconcile their arguments with results published earlier by the same group. For example, $[\text{W}(\text{CO})_5(\text{C}_2\text{H}_4)]$, one of Frenking's theoretical prototypes of a Dewar–Chatt–Duncanson (DCD) type olefin complex, displays a positive Laplacian at the W–C bond critical point (BCP) and a slightly negative $H(\mathbf{r}_c)$ value of $-0.09 \text{ Hartree } \text{\AA}^{-3}$ (Figure 3 in ref. ^[28a]). Furthermore, experimental charge-density studies of related textbook examples of DCD-type complexes such as $[\text{Ni}(\text{cod})_2]$ ($\text{cod} = \text{cyclooctadiene}$)^[28b] or $[\text{Ni}(\text{C}_2\text{H}_4)\text{dbpe}]$ ($\text{dbpe} = t\text{Bu}_2\text{PCH}_2\text{CH}_2\text{PtBu}_2$)^[28c] are characterized by positive values at the M–C BCPs ($\text{M} = \text{transition metal}$) of $+5.6$ and $+8.4 \text{ e } \text{\AA}^{-5}$, respectively. We therefore wish to end this controversy once and for all by pointing out that hardly any chemist would classify the carbon monoxide molecule as an electrostatic $[\text{C}\cdots\text{O}]$ complex lacking covalent bonding because the Laplacian displays a positive value of $+23.6 \text{ e } \text{\AA}^{-5}$ at the BCP.^[29]

Bader^[29] has clearly addressed and carefully investigated these issues over many years. It is therefore not understandable why KF quote in their comment Bader's pioneering work for the wrong reasons. In this respect KF should fully consider the statements given by Bader (page 293 of ref. ^[29]): “Thus for closed shell interactions, $\rho(\mathbf{r}_c)$ (the charge density at the BCP) is relatively low in value”. This clear and simple condition is not fulfilled in the case of **3** and $[\text{Ag}(\text{C}_2\text{H}_2)]^+$ (**3a**). The experimen-

tal charge density at the TCP of the $4d^{10}$ Ag species **3** ($\rho(\mathbf{r}_c) = 0.47 \text{ e } \text{\AA}^{-3}$) is somewhat smaller compared to the corresponding value of the DCD-type $4d^{10}$ Pd complex $[\text{Pd}(\text{C}_2\text{H}_2)]$ (**4**; $\rho(\mathbf{r}_c) = 0.79 \text{ e } \text{\AA}^{-3}$)^[26a], but significantly larger than that of the purely electrostatic π complex $[\text{Na}(\text{C}_2\text{H}_4)]^+$ (**5**; $\rho(\mathbf{r}_c) < 0.1 \text{ e } \text{\AA}^{-3}$).^[30]

Previous to our work, the electronic structures of $[\text{M}(\text{C}_2\text{H}_x)]^+$ complexes have solely been investigated by theoretical methods. We therefore analyzed the density deformations in the valence shell of the silver atom in **3**^[31] to directly reveal the significant role of covalent metal–ligand interactions in $[\text{M}(\text{C}_2\text{H}_x)]^+$ complexes. Indeed, the negative Laplacian ($L(\mathbf{r}) = -\nabla^2\rho(\mathbf{r})$) in the valence shell of the silver atom of **3** reveals a fine structure which is in conflict with “earlier theoretical studies that assumed an electrostatic bonding scenario for silver acetylene complexes”.^[2] This statement was criticized by KF “because the polarization of the charge distribution at the Ag atom may equally be the result of purely inductive interactions with the anisotropic charge distribution of the acetylene ligand”.^[1] We could hardly disagree more. Figure 2 shows a comparison of the fine structure of $L(\mathbf{r})$ of the covalent $4d^{10}$ species **4** and **3a** with that of the reference system $[\text{Rb}(\text{C}_2\text{H}_2)]^+$ (**6**) representing a purely electrostatic $4d^0$ complex employing relativistic all-electron calculations. As a result of our analysis, the electrostatic complex




Figure 2. Contour map of the negative Laplacian of the electron density ($L(\mathbf{r}) = -\nabla^2\rho(\mathbf{r})$) of a) **4**, b) **3a**, and c) **6** at the ZORA-BP86/QZV4P level of approximation in the AgC_2 plane. Contour levels are drawn at $0, \pm 2.0 \times 10^n, \pm 4.0 \times 10^n, \pm 8.0 \times 10^n \text{ e } \text{\AA}^{-5}$, where $n=0, \pm 3, \pm 2, \pm 1$. In the case of **3a** an extra level at $280 \text{ e } \text{\AA}^{-5}$ has been used while omitting the $200 \text{ e } \text{\AA}^{-5}$ contour line; positive and negative values are marked by red and blue lines, respectively. RCPs and BCPs are marked by filled squares and circles, respectively, while the bond path is shown by a black line; $\rho(\mathbf{r})$ values at the CPs are specified in $[\text{e } \text{\AA}^{-3}]$. The locations of the charge concentrations, the BCPs and TCPs are marked by arrows.

6 shows solely a weak, axial polarization. In contrast, **4** and **3a** reveal a more complex fine structure of the Laplacian showing pronounced charge-depletion zones in the molecular yz plane. This is in accordance with the expected charge transfer resulting from the π^* -(C_2H_2) $\leftarrow$ M(d_{yz}) (M = Pd, Ag) back donation in the framework of the DCD model. Our experimental charge-density study of **3** displays similar charge-depletion zones in the yz plane. We further note that the $L(r)$ topology of **3** and model **4** is in good agreement with fully relativistic four-component calculations employing basis sets close to the basis set limit (e.g. Ag/Pd: (33s, 25p, 17d, 9f, 6g, 3h; see Supporting Information). However, as pointed out in our original Communication^[2], the π^* -(C_2H_2) $\leftarrow$ M(d_{yz}) back donation is less pronounced in the silver complex than in its Group 10 analogues. Accordingly, the acetylene ligand in **4** displays a bonded charge concentration at the carbon atom pointing along the Pd–C directrix which is lacking in the case of **3a** (Figure 2). If we use the magnitude of the bonded charge concentration at the carbon atom to classify the metal atoms with respect to their π^* -(C_2H_2) $\leftarrow$ M(d_{yz}) back donation capabilities,^[28c] in the $[M(C_2H_2)]^{(+)}$ complexes (M = Ni, Pd, Cu⁺, Ag⁺) the following order can be derived $Ni > Pd > Cu^+ > Ag^+$. This result is in good agreement with the results of the energy decomposition analysis by NRF.^[23] Hence, a detailed description of the nature of the M–L bonding in **3** and **3a** has to consider orbital interactions in contrast to the over-simplifying electrostatic picture drawn by KF.

Online veröffentlicht am September 9, 2008

- [1] A. Krapp, G. Frenking, *Angew. Chem.* **2008**, DOI: 10.1002/ange.200705423; *Angew. Chem. Int. Ed.* **2008**, DOI: 10.1002/anie.200705423.
- [2] A. Reisinger, N. Trapp, I. Krossing, S. Altmannshofer, V. Herz, M. Presnitz, W. Scherer, *Angew. Chem.* **2007**, *119*, 8445; *Angew. Chem. Int. Ed.* **2007**, *46*, 8295.
- [3] It is well known that the level of approximation (MP2/def2-SVP) applied can not be classified as “high level”

approach. The selected split-valence basis set is too small for the method and a qualitative trend can be expected at best. “In case of MP2 (and other post-HF methods), def2-SVP bases may be used only for explorative calculations, results that can be considered as quantitatively satisfactory are obtained with the def2-TZVPP bases.” (see F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297).

- [4] H. Braunschweig, I. Fernández, G. Frenking, T. Kupfer, *Angew. Chem.* **2008**, *120*, 1977; *Angew. Chem. Int. Ed.* **2008**, *47*, 1951.
- [5] I. Krossing, I. Raabe, *Angew. Chem.* **2004**, *116*, 2116; *Angew. Chem. Int. Ed.* **2004**, *43*, 2066.
- [6] M. Lin, C. Liu, L. Zheng, *Wuli Huaxue Xuebao* **1995**, *11*, 266.
- [7] I. Krossing, *J. Am. Chem. Soc.* **2001**, *123*, 4603.
- [8] I. Krossing, L. Van Wüllen, *Chem. Eur. J.* **2002**, *8*, 700.
- [9] I. G. Dance, K. J. Fisher, G. D. Willett, *Inorg. Chem.* **1996**, *35*, 4177.
- [10] T. S. Cameron, A. Decken, I. Dionne, M. Fang, I. Krossing, J. Passmore, *Chem. Eur. J.* **2002**, *8*, 3386.
- [11] F. Meyer, Y.-M. Chen, P. B. Armentrout, *J. Am. Chem. Soc.* **1995**, *117*, 4071.
- [12] J. L. M. Abboud, O. Castano, M. Hereros, J. Elguero, N. Jagerovic, R. Notario, K. Sak, *Int. J. Mass Spectrom. Ion Proc.* **1998**, *175*, 35.
- [13] I. Krossing, A. Bihlmeier, I. Raabe, N. Trapp, *Angew. Chem.* **2003**, *115*, 1569; *Angew. Chem. Int. Ed.* **2003**, *42*, 1531.
- [14] I. Raabe, D. Himmel, S. Müller, N. Trapp, M. Kaupp, I. Krossing, *Dalton Trans.* **2008**, 946.
- [15] B. C. Guo, A. W. Castleman, Jr., *Chem. Phys. Lett.* **1991**, *181*, 16.
- [16] Z. Tian, Z. Tang, *Rapid Commun. Mass Spectrom.* **2005**, *19*, 2893.
- [17] I. Krossing, A. Reisinger, *Angew. Chem.* **2003**, *115*, 5903; *Angew. Chem. Int. Ed.* **2003**, *42*, 5725.
- [18] A. Reisinger, Dissertation, Cuvillier, Göttingen **2006**.
- [19] In our experience, unscaled BP86/SV(P) vibrational frequencies reproduce experimental data remarkably well at low computational cost. Harmonic MP2/TZVPP vibrations usually need scaling by 0.93 to 0.95 and, even if scaled, are in our experience not more accurate, despite their much higher computational effort. Thus we refrained from using scaled MP2 frequencies. The difference in the calculated thermodynamics by KF and us should result from the use of different frequency calculations for statistical thermodynamics.
- [20] H. D. B. Jenkins, H. K. Roobottom, J. Passmore, L. Glasser, *Inorg. Chem.* **1999**, *38*, 3609.
- [21] H. D. B. Jenkins, L. Glasser, *Inorg. Chem.* **2003**, *42*, 8702.
- [22] The DFT calculations were performed at the same level as those of KF (B3LYP/def-ECP(Ag)/def2-TZVPP level of approximation).
- [23] M. S. Nechaev, V. M. Rayón, G. Frenking, *J. Phys. Chem. A* **2004**, *108*, 3134.
- [24] a) M. Böhme, T. Wagener, G. Frenking, *J. Organomet. Chem.* **1996**, *520*, 31; b) J. Miralles-Sabater, M. Merchán, I. Nebot-Gil, P. M. Viruela-Martín, *J. Phys. Chem.* **1988**, *92*, 4853; c) M. Sodupe, C. W. Bauschlicher, *J. Phys. Chem.* **1991**, *95*, 8640. We note, that ref. [25] was erroneously quoted by NRF along with ref. [24b,c] as a case study in favor of mainly electrostatic interactions. However, Hertwig et al. clearly stated in their paper that “for all three metal cations (Cu, Ag, Au), the interaction with ethylene shows large covalent contributions”. Hence, Hertwig et al. came to the same conclusions as NRF—but eight years earlier.
- [25] R. H. Hertwig, W. Koch, D. Schröder, H. Schwarz, J. Hrušák, P. Schwerdtfeger, *J. Phys. Chem.* **1996**, *100*, 12253.
- [26] a) G. Eickerling, R. Mastalerz, V. Herz, W. Scherer, H.-J. Himmel, M. Reiher, *J. Chem. Theory Comput.* **2007**, *3*, 2182; b) N. Hebben, H.-J. Himmel, G. Eickerling, C. Herrmann, M. Reiher, V. Herz, M. Presnitz, W. Scherer, *Chem. Eur. J.* **2007**, *13*, 10078; c) G. Eickerling, M. Reiher, *J. Chem. Theory Comput.* **2008**, *4*, 286.
- [27] B. Rohrmoser, G. Eickerling, M. Presnitz, W. Scherer, V. Eyert, R.-D. Hoffmann, U. C. Rodewald, C. Vogt, R. Pöttgen, *J. Am. Chem. Soc.* **2007**, *129*, 9356.
- [28] a) G. Frenking, U. Pidun, *J. Chem. Soc. Dalton Trans.* **1997**, 1653; b) P. Macchi, D. M. Proserpio, A. Sironi, *J. Am. Chem. Soc.* **1998**, *120*, 1447; c) W. Scherer, G. Eickerling, D. Shorokhov, E. Gullo, G. S. McGrady, P. Sirsch, *New J. Chem.* **2006**, *30*, 309.
- [29] R. F. W. Bader, *Atoms in Molecules. A Quantum Theory*, Oxford University Press, Oxford, **1994**, p. 312.
- [30] C. F. Matta, R. J. Boyd, *The Quantum Theory of Atoms in Molecules*, Wiley-VCH, Weinheim, **2007**.
- [31] For a general discussion of the shell structure of transition metal atoms, see G. S. McGrady, A. Haaland, H. P. Verne, H. V. Volden, A. J. Downs, D. Shorokhov, G. Eickerling, W. Scherer, *Chem. Eur. J.* **2005**, *11*, 4921 and ref. [26].