

Nitrides


C. L. Schmidt, U. Wedig, R. Dinnebier, M. Jansen*

Synthesis, Crystal Structure, Bonding, and Properties of $(\text{Ba}_6\text{O})(\text{OsN}_3)_2$

Chem. Asian J.

DOI: 10.1002/asia.200800232

4+ preferred: The structure, bonding, and physical properties of a novel nitridoosmate $(\text{Ba}_6\text{O})(\text{OsN}_3)_2$ are presented, containing trigonal, virtually planar OsN_3^{5-} anions. The oxidation state 4+ of osmium is stabilized by oxygen in the Ba partial structure.




Natural Products

C. Mehner, D. Müller, S. Kehraus, S. Hautmann, M. Gütschow, G. M. König*


New Peptolides from the Cyanobacterium *Nostoc insulare* as Selective and Potent Inhibitors of Human Leukocyte Elastase

ChemBioChem

DOI: 10.1002/cbic.200800415



- NRPS gene fragments
- MALDI-TOF detection of peptides
- HLE inhibition



Combinatorial chemistry in cyanobacteria. *Nostoc insulare* has the biosynthetic capability to produce an impressive library of nonribosomal peptides. Molec-

ular biological investigations and bioactivity studies led to the isolation of potent and selective inhibitors of the enzyme human leukocyte elastase (HLE).


Electrochemistry

C. Costentin, M. Robert, J.-M. Savéant*, A.-L. Teillout

Concerted and Stepwise Proton-Coupled Electron Transfers in Aquo/Hydroxo Complex Couples in Water: Oxidative Electrochemistry of $[\text{Os}^{\text{II}}(\text{bpy})_2(\text{py})(\text{OH}_2)]^{2+}$

ChemPhysChem

DOI: 10.1002/cphc.200800361



To be or not to be concerted. That is the question. Successive oxidation of transition metal(II) aqua complexes ($\text{M}^{\text{II}}\text{OH}_2$ to $\text{M}^{\text{III}}\text{OH}$) is a domain in which proton-coupled electron-transfer reactions are extremely common (see figure). The concerted (CPET) or stepwise (EPT, PET) character of the mechanism is an important fundamental issue for understanding and designing natural or artificial catalytic systems.



Antitumor Agents

L. G. León, R. M. Carballo, M. C. Vega-Hernández, P. O. Miranda, V. S. Martín, J. I. Padrón, J. M. Padrón*

β' -Hydroxy- α,β -unsaturated ketones: A new pharmacophore for the design of anticancer drugs. Part 2.


ChemMedChem

DOI: 10.1002/cmdc.200800212




An iron(III)-catalyzed multicomponent domino process is the key step in the synthesis of β' -acyloxy- α,β -unsaturated ketones. These novel compounds arrest the cell cycle and trigger apoptosis by

irreparably damaging cancer cells. The biological activity of these compounds seem parallel to those observed with the antitumor natural product, persin.



The different forms of a rhombic $[\text{Mn}_4]$ single-molecule magnet and their assembly into infinite coordination networks are reviewed as a case study of how the rational design of extended arrays of high-spin complexes may afford unprecedented, interesting magnetic properties. Potential synthetic strategies and outcomes are discussed.




Assemblies of High-Spin Complexes

O. Roubeau,* R. Clérac

Rational Assembly of High-Spin Polynuclear Magnetic Complexes into Coordination Networks: the Case of a $[\text{Mn}_4]$ Single-Molecule Magnet Building Block

Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.200800603



A variety of mono- and polytopic 1,2,3-triazole β -CD derivatives have been synthesized by click chemistry. The synthetic procedure is based on the Cu^{I} -catalyzed azide–alkyne cycloaddition reaction between hydroxylated or randomly methylated β -CD monoazides and alkynyl precursors. Easy to use, the reaction is also high-yielding for many molecules.

β -Cyclodextrin Hosts

M. Mourer, F. Hapiot, S. Tilloy, E. Monflier, S. Menuel*

Easily Accessible Mono- and Polytopic β -Cyclodextrin Hosts by Click Chemistry

Eur. J. Org. Chem.
DOI: 10.1002/ejoc.200800728

Room at the anion? Three diindolylureas assemble around PO_4^{3-} in the solid-state, saturating its coordination sphere with twelve hydrogen bonds (see figure).




Anion Binding

C. Caltagirone, J. R. Hiscock, M. B. Hursthouse, M. E. Light, P. A. Gale*

1,3-Diindolylureas and 1,3-Diindolylthioureas: Anion Complexation Studies in Solution and the Solid State

Chem. Eur. J.
DOI: 10.1002/chem.200801639



Repeat performances: Several enlarged ruthenium-based catalysts were prepared and tested in an olefin metathesis reaction coupled to a nanofiltration (NF) process, aimed at recovering and recycling the catalyst. Retention of the catalyst

was as high as 92%, and five catalytic cycles could be performed in a test ring-closing metathesis reaction in dimethyl carbonate (DMC) before a decrease in the catalyst activity was observed.

Olefin Metathesis

A. Keraani, T. Renouard, C. Fischmeister,* C. Bruneau, M. Rabiller-Baudry*

Recovery of Enlarged Olefin Metathesis Catalysts by Nanofiltration in an Eco-Friendly Solvent

ChemSusChem
DOI: 10.1002/cssc.200800152



Auf diesen beiden Seiten weisen wir auf wichtige aktuelle Beiträge in unseren Schwesterzeitschriften hin. Wenn Sie die Seiten online lesen, dann können Sie die

Artikel mit einem Klick direkt aufrufen, ansonsten sind sie durch Eingabe der DOIs über Wiley InterScience leicht online zugänglich.