


Dinitrogen Reduction and C–H Activation by the Divalent Organoneodymium Complex [(C₅H₂tBu₃)₂Nd(μ-I)K([18]crown-6)]**

Florian Jaroschik, Aurélien Momin, François Nief,* Xavier-Frédéric Le Goff, Glen B. Deacon, and Peter C. Junk*

The isolation and characterization of the first molecular complex of divalent thulium, [TmI₂(dme)₃] (dme = 1,2-dimethoxyethane),^[1] ten years ago, followed by the dysprosium^[2] and neodymium^[3] diiodides, [DyI₂(dme)₃] and [NdI₂(thf)₅], opened up a new era in lanthanide chemistry.^[4] Hitherto, divalent lanthanide chemistry had been restricted for decades to the “classical” Eu^{II}, Yb^{II}, and Sm^{II}.^[4] These “new” divalent complexes have since been investigated in numerous reactions, often revealing unexpected behavior.^[5,6] However, the expectation that these precursors would lead to the isolation of divalent organometallic complexes by salt metathesis reactions was fulfilled only in the case of several very crowded Tm^{II} cyclopentadienyl, phospholyl, and scorpionate complexes.^[7] In contrast, all reported attempts with DyI₂ and NdI₂, or the hypothetical [NdCl₂(thf)_x], afforded only trivalent products.^[7a,8] An early report on the identification of an organoneodymium(II) complex by elemental analysis, IR spectroscopy, and reactivity^[8b] has been questioned.^[8c] Lappert and co-workers attempted the synthesis of La^{II}, Ce^{II}, Nd^{II}, and Tm^{II} complexes by reduction of trivalent precursors of the type [Cp^R₃Ln] (Cp^R = C₅H₃tBu₂, C₅H₃(SiMe₃)₂) using alkali metals^[9] and achieved the isolation of the first crystallographically characterized La^{II} derivatives^[9b,e] and recently a mixed Ce^{II}/Ce^{III} complex.^[9e] A similar approach used by Cloke and co-workers afforded a mixed Sc^I/Sc^{III} compound.^[10] The potential of the “new” divalent lanthanide chemistry has recently been highlighted.^[11] We extended the reductive approach to obtain stable, solvent-free divalent organothulium complexes carrying cyclopentadienyl or phospholyl ligands by reduction of trivalent precursors of the L₂LnX type with potassium graphite in nonpolar solvents.^[12] This

method also gave access to the first divalent organodysprosium ate complexes [Cp^{III}Dy(μ-X)K([18]crown-6)] (Cp^{III} = C₅H₂tBu₃ and X = I (**1**), Br, BH₄), which are powerful reducing agents.^[13] Despite several attempts at reducing trivalent Nd precursors, no organoneodymium(II) complexes could be isolated, owing to decomposition of any Nd^{II} species in situ by interaction with solvent molecules, such as DME, THF, or benzene.^[9a,c-e] We herein report on the synthesis and characterization of the first divalent organoneodymium complex, its behavior towards molecular dinitrogen, and its decomposition by C–H activation to form a structurally characterized “tuck-in” complex, the first such example for a 4f element.^[14]

Reduction of the trivalent precursor [Cp^{III}NdI] (**2**), prepared from NdI₃ and [KCp^{III}], with potassium graphite under argon in the presence of [18]crown-6 led to an immediate color change of the solution from green-blue to dark red. After 24 h, dark red crystals of the divalent ate complex [Cp^{II}Nd(μ-I)K([18]crown-6)]·C₆H₁₂ (**3**), suitable for X-ray crystallography, had formed. (Scheme 1 and Figure 1)



Scheme 1. Synthesis of divalent Nd ate complex **3** and trivalent Nd “tuck-in” complex **4**.

The crystal structure of **3** shows a seven-coordinate neodymium center in an open sandwich array (Cp–Nd–Cp 149.4°) with an iodine bridge to the potassium–crown ether complex. The balance of charges indicates that **3** is indeed a Nd^{II} complex that is best described as a [K([18]crown-6)I] adduct of the neutral [Cp^{II}Nd] fragment. The geometrical arrangement is similar to the previously reported divalent organodysprosium ate complex **1**.^[13] The longer detected Ln–I (3.16 Å for Ln = Nd, 3.06 Å for Ln = Dy) and Ln–Cp(centroid) (2.53 Å for Ln = Nd, 2.44 Å for

[*] A. Momin, Dr. F. Nief, Dr. X.-F. Le Goff
Laboratoire “Hétéroéléments et Coordination” (DCPH), Ecole Polytechnique
91128 Palaiseau (France)
Fax: (+33) 1-6933-4440
E-mail: francois.nief@polytechnique.edu
Dr. F. Jaroschik, Prof. G. B. Deacon, Prof. P. C. Junk
School of Chemistry, Monash University
3800 Clayton, Victoria (Australia)
E-mail: peter.junk@sci.monash.edu.au

[**] We acknowledge financial support from the Ecole Polytechnique, the CNRS and the Australian Research Council. We thank Mr. Finlay Shanks (Monash University) for the Raman measurements, Dr. Boujemaa Moubaraki (Monash University) for the magnetic moment measurement, and Dr. Roger Mulder (CSIRO) for the ¹⁵N NMR measurement.

Supporting Information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200804934>.




Figure 1. Molecular structure of $[\text{Cp}_2^{\text{III}}\text{Nd}(\mu\text{-I})\text{K}([18]\text{crown-6})]$ (**3**). Thermal ellipsoids are set at 50% probability and hydrogen atoms are omitted for clarity. Cyclohexane molecules present in the lattice are not represented.

$\text{Ln}=\text{Dy}$) distances in **3** compared with **1** are in good agreement with the corresponding lanthanide radii.^[15] Compared to the trivalent ate complex $[\text{Cp}_2^{\text{III}}\text{Nd}(\mu\text{-Cl})\text{AlMe}_3]$ ($\text{Nd}\text{--}\text{Cp}$ 2.48 Å),^[16] the $\text{Nd}\text{--}\text{Cp}(\text{centroid})$ distance is significantly longer in **3**.

The reduction of **2** was monitored by ^1H NMR spectroscopy in deuterated cyclohexane. Initially, the *t*Bu resonances of the trivalent precursor **2** were detected at $\delta = -0.9$, -11.6 , and -18.1 ppm, with an integration of 1:1:1, in agreement with reported $[\text{Cp}_2^{\text{III}}\text{LnX}]$ complexes.^[12,13,16,17] Upon reduction to the divalent compound **3**, the *t*Bu resonances shift to a signal consisting of two broad overlapping peaks at $\delta = -4.8$ and -6.7 ppm and another signal at $\delta = -19.4$ ppm, in a 2:1 ratio. In addition, there is a sharp peak at $\delta = -0.05$ ppm, arising from the crown ether. These data are consistent with the solid-state structure of **3**.

Compound **3** was isolated as a dark red powder, which was stable under argon at low temperature for several days, but degraded rapidly at room temperature. Once isolated, **3** was only very slightly soluble in cyclohexane, even after heating up to 60 °C, whereupon decomposition occurred. Attempts to use other solvents, such as benzene, toluene, diethyl ether, THF, or DME, led to the immediate decomposition of **3**, as indicated by a color change to the typical blue-green color of trivalent Nd compounds, in stark contrast to the divalent Dy complexes, which are stable in toluene for several days.^[13]

The room-temperature magnetic moment of solid **3** ($2.8\mu_{\text{B}}$), obtained by SQUID susceptibility measurements, corresponds very well to the theoretical value for Nd^{II} in the $4f^{[4]}$ configuration ($2.83\mu_{\text{B}}$) and to that for $[\text{NdI}_2(\text{thf})_5]$ ($2.7\mu_{\text{B}}$),^[3b] and is markedly different from that for Nd^{III} ($3.68\mu_{\text{B}}$).^[18] This confirmation of the oxidation state by magnetic moment is in contrast to the failure of the technique for the divalent Dy complex **1**, owing to the similarity of the values for Dy^{II} and Dy^{III} ($10.6\mu_{\text{B}}$).

A UV/Vis/NIR spectrum was recorded for a fresh, deep red solution of **3** generated in situ in cyclohexane (ruling out

extinction coefficient data). Absorption bands at 400 (m), 578 (br, sh), and 1500–2200 nm (br) were attributed to Nd^{II} , as they were not present in the spectrum of the solution after exposure to air, or in the spectrum of **2**. Furthermore, each of these bands occurs in regions corresponding to one or more $f\leftarrow f$ bands of Pm^{3+} ,^[19] which is isoelectronic with Nd^{2+} . The relative intensity of the band at 400 nm is greater than normal for an $f\leftarrow f$ absorption, suggesting a contribution from a $d\leftarrow f$ transition, as are featured in the visible spectra of Sm^{2+} , Eu^{2+} , and Yb^{2+} .^[20]

The red solution of **3** in $[\text{D}_2]$ cyclohexane decomposed over several days to a green-blue solution, which afforded a complex ^1H NMR spectrum. No product could be identified from this solution because freezing of cyclohexane prevents crystallization at low temperatures. On reduction of **2** in hexane under the usual conditions, the initial dark red Nd^{III} solution turned yellow-green within 30 min. Cooling this solution to -30°C led to the formation of a small quantity of blue crystals. X-ray crystallographic analysis of these crystals revealed the first crystallographically identified “tuck-in” complex in the lanthanide series, $[\text{Cp}^{\text{III}}\{\mu\text{-C}_5\text{H}_2\text{tBu}_2\text{C}(\text{CH}_3)_2\text{-(CH}_2)_2\}\text{Nd}(\mu\text{-I})\text{K}([18]\text{crown-6})]\cdot\text{C}_6\text{H}_{14}$ (**4**, Scheme 1, Figure 2).




Figure 2. Molecular structure of the “tuck-in” complex $[\text{Cp}^{\text{III}}\{\text{C}_5\text{H}_2\text{tBu}_2\text{C}(\text{CH}_3)_2(\text{CH}_2)_2\}\text{Nd}(\mu\text{-I})\text{K}([18]\text{crown-6})]$ (**4**). Thermal ellipsoids are set at 50% probability and hydrogen atoms are omitted for clarity (except for the methylene group). Hexane molecules present in the lattice are not represented.

In this sandwich complex, the trivalent eight-coordinate Nd center forms a metallacycle with a methylene group from the Cp^{III} ligand and is bound to $[\text{K}([18]\text{crown-6})\text{I}]$ to form an ate complex, analogous to **3**. The poor precision of the structure does not allow a detailed analysis but the bond length between Nd and the inserted methylene carbon (2.5 Å) is in the typical range of $\text{Nd}\text{--}\text{C}$ σ bonds, for example 2.515 Å in $[(\text{C}_5\text{Me}_5)_2\text{NdCH}(\text{SiMe}_3)_2]$.^[21] The formation of **4** may result from the activation of one C–H bond of a *t*Bu group in the coordination sphere of the strongly reactive Nd^{II} center. Andersen and co-workers prepared the related complex

[Cp^{III}[(μ-C₅H₂tBu₂C(CH₃)₂(CH₂))Ce] by thermolysis of [Cp^{III}Ce(CH₂Ph)], but it was not characterized by X-ray crystallography.^[17] Recently, Evans et al. reported the structure of a uranium “tuck-in”–“tuck-over” complex with the C₅Me₅ ligand.^[14]

Dinitrogen activation by well-defined divalent organolanthanide complexes has to date only been achieved with [SmCp₂^{*}] and in the reaction of a Tm^{II} alkyl complex with an amine under dinitrogen.^[6,7e] Other dinitrogen-containing complexes were obtained either by salt metathesis under a dinitrogen atmosphere using the divalent lanthanide diiodides LnI₂ (Ln = Sm, Tm, Dy, Nd) or by the LnZ₃/K/N₂ system (Z = substituted Cp, phenoxide, or amide ligands) and other in situ reduction systems.^[6,22]

Reduction of **2** with KC₈ under a dinitrogen atmosphere in hexane at room temperature gave a yellow solution. Single crystals of [(Cp^{III}Nd)₂(μ-η²:η²-N₂)]·C₆H₁₄ (**5**) with a distinct orange-green dichroism and which were suitable for X-ray diffraction studies were isolated after cooling the solution at –30 °C for several days. The crystal structure of **5** (Figure 3)





Figure 3. Molecular structure of [(Cp^{III}Nd)₂(μ-η²:η²-N₂)] (**5**). Thermal ellipsoids are set at 50% probability and hydrogen atoms are omitted for clarity. Hexane molecules present in the lattice are not represented.

incorporates two {Cp^{III}Nd} sandwich moieties bridged by a reduced dinitrogen molecule. The formation and stability of this binuclear complex is surprising given the steric bulk of the ligand: the Cp^{III}–Nd–Cp^{III} angle is reduced to 131.8° compared with 149.4° in **3**, leading to an elongation of the Cp^{III}–(centroid)–Nd distance in trivalent **5** which is nearly identical to that in divalent **3**. The N=N bond length in **5** (1.23 Å) is in the range of other reported Nd dinitrogen complexes^[23] and significantly shorter than the O–O bond length in peroxo-bridged complexes, for example in [Yb₂{N(SiMe₃)₂}₄(μ-O₂)(thf)₂] (1.543 Å).^[24] A Raman spectrum of crystalline **5** with a very low laser power showed a sharp strong peak at 1622 cm^{–1}, which was attributed to the symmetric stretching of the N=N double bond. This assignment was confirmed by the shift of this signal to 1569 cm^{–1} for the ¹⁵N-labeled complex ¹⁵N-**5** (expected value: 1567 cm^{–1}). This significant

weakening of the N=N bond ($\bar{\nu}[\text{free N}_2] = 2331 \text{ cm}^{-1}$) is in agreement with other organometallic dinitrogen-containing complexes, such as LFeNNFeL (L = β-diketiminato).^[25] To our knowledge, this is the first N=N double bond in lanthanide complexes detected by Raman spectroscopy.^[26] The ¹H NMR spectrum of **5** in [D₆]benzene shows three sharp resonances at δ = 10.61, –1.77, and –23.48 ppm, which indicates that, as in **2**, free rotation of the Cp^{III} ligand is hampered. In contrast to **2**, which shows coalescence at 50 °C, no coalescence occurs for **5** at 60 °C. When the reduction was carried out in [D₁₂]cyclohexane, the same resonances were detected. Despite several attempts, no clearly defined signal could be obtained in the ¹⁵N NMR spectrum of ¹⁵N-**5**. This contrasts the resonances for the slightly less paramagnetic Ce and Pr dinitrogen complexes.^[27]

When a solution of **3** in [D₁₂]cyclohexane was exposed to dinitrogen, an immediate color change from dark red to yellow occurred, and a white precipitate formed. The ¹H NMR spectrum of the yellow solution showed as main product the three sharp peaks identical to the solution of isolated **5**, in addition to several broad peaks that could not be assigned. The white precipitate was found to contain KI, KCp^{III}, and [18]crown-6 (Scheme 2).



Scheme 2. Synthesis of dinitrogen-containing dinuclear Nd complex **5**.

In summary, a careful choice of ligand and solvent system allowed the isolation of the first divalent organoneodymium complex. The highly reactive Nd^{II} center can activate C–H bonds to form a “tuck-in” complex. The reactivity of the divalent complex with dinitrogen led to the formation of the same product as in the [Cp^{III}NdI]/K/N₂ system. Further reactivity studies on the three isolated products are currently in progress.

Experimental Section

Divalent **3** is exceptionally reactive to air, moisture, and solvents, much more so than Tm^{II} and Dy^{II} complexes,^[7a–d,12] and solid **3** is also thermally unstable. Compound **4** is thermally unstable and must also be handled under high-purity argon.

3: a) In an argon atmosphere, KC₈ (5.0 mg, 0.037 mmol) was added to a solution of **2** (10 mg, 0.014 mmol), and [18]crown-6 (3.6 mg, 0.014 mmol) in [D₁₂]cyclohexane (1 mL) in an NMR tube fitted with a J. Young valve, and the reaction mixture was shaken immediately to give quantitatively a dark red solution of **3**. Crystals of **3** suitable for X-ray diffraction formed over 24 h from this solution at room temperature. ¹H NMR (300 MHz, [D₁₂]cyclohexane): δ = –0.05 (s, 24H, [18]crown-6), –4.8 and –6.7 (br s, 36H, ω_{1/2} ≈ 1 kHz, *t*Bu), –19.4 ppm (br s, 18H, ω_{1/2} ≈ 0.5 kHz, *t*Bu).

b) In a similar experiment, the reaction of **2** (20 mg, 0.027 mmol), [18]crown-6 (7.2 mg, 0.027 mmol), and KC₈ (10 mg, 0.074 mmol) was carried out in cyclohexane (3 mL) in a centrifugation tube. The solids

were removed by centrifugation and the solvent was carefully removed under vacuum. The residue was washed with fresh cyclohexane (2 mL) to afford **3** as a dark red powder (14 mg, 0.014 mmol, 51 %). $\mu_{\text{eff}} = 2.8 \mu_{\text{B}}$. IR (Nujol): $\tilde{\nu} = 1580$ (w), 1352 (m), 1260 (w), 1239 (w), 1111 (s), 1019 (m), 963 (m), 798 cm^{-1} (m). UV/Vis/NIR (C_6H_{12}): $\nu = 400$ (m), 578 (br, sh), 1500–2200 nm (vvbr).

4: In a similar synthesis to **3** (b), the reaction of **2** (10 mg, 0.013 mmol), [18]crown-6 (3.6 mg, 0.014 mmol), and KC_8 (5.0 mg, 0.037 mmol) was carried out in hexanes (2 mL) in a centrifugation tube. After centrifugation for 5 min at 3900 rpm the solution was cooled to -30°C , yielding **4** as blue crystals suitable for X-ray analysis (1.5 mg, 0.002 mmol, 15 %). ^1H NMR (300 MHz, $[\text{D}_6]\text{benzene}$): $\delta = 24.1$ (s, 3H, CMe_2), 14.0 and 12.4 (br s, 12H, $\omega_{1/2} \approx 0.3$ kHz, $t\text{Bu}$ and CMe_2), 2.8 (s, 9H, $t\text{Bu}$), 0.73 (s, 24H, [18]crown-6), -0.7 (s, 9H, $t\text{Bu}$), -10.1 (s, 9H, $t\text{Bu}$), -15.4 ppm (br s, 9H, $\omega_{1/2} \approx 0.2$ kHz, $t\text{Bu}$). No signal for the Cp ring and methylene protons was detected.

5: a) In a dinitrogen atmosphere, a mixture of **1** (20 mg, 0.027 mmol), KC_8 (36 mg, 0.270 mmol), and $[\text{D}_{12}]\text{cyclohexane}$ (1 mL), in an NMR tube fitted with a J. Young valve, was kept at 50°C for 24 h. After removal of the solvent under reduced pressure, recrystallization from hexanes at -30°C afforded **5** as orange-green dichroic crystals suitable for X-ray analysis (22 mg, 0.017 mmol, 62 %). A similar procedure was applied for the synthesis of the ^{15}N -labeled compound ^{15}N -**5**. ^1H NMR (300 MHz, $[\text{D}_{12}]\text{cyclohexane}$): $\delta = 24.6$ (s, 8H, Cp-H), 10.6 (s, 36H, $t\text{Bu}$), -1.7 (br s, 36H, $t\text{Bu}$), -23.4 ppm (s, 36H, $t\text{Bu}$). Raman: 2963 (m), 2939 (s), 2875 (s), 1742 (w), 1622 (s) [$(\tilde{\nu}_s(^{14}\text{N}_2))$], 1448 (m), 816 cm^{-1} (w). For ^{15}N -**5** [$(\tilde{\nu}_s(^{15}\text{N}_2))$] 1569 cm^{-1} (s).

b) A dark red solution of **3** (14 mg, 0.014 mmol) in $[\text{D}_{12}]\text{cyclohexane}$ (1 mL) was exposed to dinitrogen, leading to an immediate color change to yellow and the slow formation of a white precipitate. After 24 h, the white solid was removed by centrifugation and identified as a mixture of KI, [18]crown-6, and KCp^{III} , as confirmed by ^1H NMR and IR spectroscopy and by qualitative analysis. The ^1H NMR spectrum of the remaining solution showed compound **5** as the major product. The solvent was removed under reduced pressure and the crude product recrystallized from hexane at -30°C to afford **5** (6 mg, 0.004 mmol, 34 %).

X-ray Crystallographic Data

3: Space group $P\bar{1}$; $a = 13.593(1)$, $b = 14.136(1)$, $c = 16.101(1)$ Å; $\alpha = 81.546(1)$, $\beta = 74.412(1)$, $\gamma = 70.613(1)^\circ$; $V = 2805.4(3)$ Å 3 ; $Z = 2$; $\rho_{\text{calcd}} = 1.332$ Mg m^{-3} ; $R1 = 0.0607$ [$I > 2\sigma(I)$]; $wR2 = 0.1587$; GOF = 1.025.

4: Space group $P\bar{1}$; $a = 13.561(3)$, $b = 13.850(3)$, $c = 15.982(3)$ Å; $\alpha = 88.92(3)$, $\beta = 68.66(3)$, $\gamma = 85.59(3)^\circ$; $V = 2787.7(1)$ Å 3 ; $Z = 2$; $\rho_{\text{calcd}} = 1.342$ Mg m^{-3} ; $R1 = 0.0882$ [$I > 2\sigma(I)$]; $wR2 = 0.1514$; GOF = 1.119.

5: Space group $P2_1/n$; $a = 18.342(4)$, $b = 10.403(2)$, $c = 19.369(4)$ Å; $\beta = 108.77(3)^\circ$; $V = 3499.4(1)$ Å 3 ; $Z = 2$; $\rho_{\text{calcd}} = 1.268$ Mg m^{-3} ; $R1 = 0.0728$ [$I > 2\sigma(I)$]; $wR2 = 0.1341$; GOF = 1.209.

The synthesis of **2** is presented in the Supporting Information together with X-ray collection and processing details. CCDC 704850, 704850, 704852 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Received: October 9, 2008

Published online: December 23, 2008

Keywords: C–H activation · dinitrogen activation · dysprosium · neodymium · sandwich complexes

Angew. Chem. **1997**, *109*, 123–124; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 133–135.

- [2] W. J. Evans, N. T. Allen, J. W. Ziller, *J. Am. Chem. Soc.* **2000**, *122*, 11749–11750.
- [3] a) M. N. Bochkarev, I. L. Fedushkin, S. Dechert, A. A. Fagin, H. Schumann, *Angew. Chem.* **2001**, *113*, 3268–3270; *Angew. Chem. Int. Ed.* **2001**, *40*, 3176–3178; b) M. N. Bochkarev, A. A. Fagin, *Chem. Eur. J.* **1999**, *5*, 2990–2992.
- [4] W. J. Evans, *Inorg. Chem.* **2007**, *46*, 3435–3449.
- [5] a) M. N. Bochkarev, *Coord. Chem. Rev.* **2004**, *248*, 835–851; b) Z. Zhu, C. Wang, X. Xiang, C. Pi, X. Zhou, *Chem. Commun.* **2006**, 2066–2068; c) X. Xiang, Q. Shen, J. Wang, Z. Zhu, W. Huang, X. Zhou, *Organometallics* **2008**, *27*, 1959–1962.
- [6] a) W. J. Evans, *Can. J. Chem.* **2005**, *83*, 375–384.
- [7] a) W. J. Evans, N. T. Allen, J. W. Ziller, *Angew. Chem.* **2002**, *114*, 369–371; *Angew. Chem. Int. Ed.* **2002**, *41*, 359–361; b) F. Nief, D. Turcitu, L. Ricard, *Chem. Commun.* **2002**, 1646–1647; c) D. Turcitu, F. Nief, L. Ricard, *Chem. Eur. J.* **2003**, *9*, 4916–4923; d) F. Nief, B. T. de Borms, L. Ricard, D. Carmichael, *Eur. J. Inorg. Chem.* **2005**, 637–643; e) J. Cheng, J. Takats, M. J. Ferguson, R. McDonald, *J. Am. Chem. Soc.* **2008**, *130*, 1544–1545.
- [8] a) L. J. Bowman, K. Izod, W. Clegg, R. W. Harrington, *Organometallics* **2007**, *26*, 2646–2651; b) M. Wedler, A. Recknagel, F. T. Edelmann, *J. Organomet. Chem.* **1990**, *395*, C26–C29; c) E. N. Kirillov, A. A. Trifonov, S. E. Nefedov, I. L. Eremenko, F. T. Edelmann, M. N. Bochkarev, *Z. Naturforsch. B* **1999**, *54*, 1379–1384.
- [9] a) Y. K. Gun'ko, P. B. Hitchcock, M. F. Lappert, *J. Organomet. Chem.* **1995**, *499*, 213–219; b) M. C. Cassani, D. J. Duncalf, M. F. Lappert, *J. Am. Chem. Soc.* **1998**, *120*, 12958–12959; c) S. Al-Juaid, Y. K. Gun'ko, P. B. Hitchcock, M. F. Lappert, S. Tian, *J. Organomet. Chem.* **1999**, *582*, 143–152; d) P. B. Hitchcock, M. F. Lappert, S. Tian, *Organometallics* **2000**, *19*, 3420–3428; e) P. B. Hitchcock, M. F. Lappert, L. Maron, A. V. Protchenko, *Angew. Chem.* **2008**, *120*, 1510–1513; *Angew. Chem. Int. Ed.* **2008**, *47*, 1488–1491.
- [10] G. K. B. Clentsmith, F. G. N. Cloke, J. C. Green, J. Hanks, P. B. Hitchcock, J. F. Nixon, *Angew. Chem.* **2003**, *115*, 1068–1071; *Angew. Chem. Int. Ed.* **2003**, *42*, 1038–1041.
- [11] G. Meyer, *Angew. Chem.* **2008**, *120*, 5040–5042; *Angew. Chem. Int. Ed.* **2008**, *47*, 4962–4964.
- [12] a) F. Jaroschik, F. Nief, L. Ricard, *Chem. Commun.* **2006**, 426–428; b) F. Jaroschik, F. Nief, X.-F. Le Goff, L. Ricard, *Organometallics* **2007**, *26*, 3552–3558.
- [13] F. Jaroschik, F. Nief, L. Ricard, X.-F. Le Goff, *Organometallics* **2007**, *26*, 1123–1125.
- [14] W. J. Evans, K. A. Miller, A. G. DiPasquale, A. L. Rheingold, T. J. Stewart, R. Bau, *Angew. Chem.* **2008**, *120*, 5153–5156; *Angew. Chem. Int. Ed.* **2008**, *47*, 5075–5078. The “tuck-in” complexes contrast with dinuclear “tuck-over” complexes, in which the metalated ligand is bonded to two metals. Such complexes are known for rare earths: W. J. Evans, T. A. Ullbarri, J. W. Ziller, *Organometallics* **1991**, *10*, 134–142; M. Booi, B. J. Deelman, R. Duchateau, D. J. Postma, A. Meetsma, J. H. Teuben, *Organometallics* **1993**, *12*, 3531–3540; W. J. Evans, J. M. Perotti, J. W. Ziller, *Inorg. Chem.* **2005**, *44*, 5820–5825; W. J. Evans, T. M. Champagne, J. W. Ziller, *J. Am. Chem. Soc.* **2006**, *128*, 14270–14271. There are several group 4 and group 5 “tuck-in” complexes featuring a metalated $t\text{Bu}$ group on a Cp ring: X. Yang, C. L. Stern, T. J. Marks, *J. Am. Chem. Soc.* **1994**, *116*, 10015–10031; C. Legrand, P. Meunier, J. L. Petersen, P. Tavares, J. Bodiguel, B. Gautheron, G. Dousse, *Organometallics* **1995**, *14*, 162–169; D. J. Duncalf, R. J. Harrison, A. McCamley, B. W. Royan, *J. Chem. Soc. Chem. Commun.* **1995**, 2421–2422.
- [15] R. D. Shannon, *Acta Crystallogr. Sect. A* **1976**, *32*, 751–767.

[1] M. N. Bochkarev, I. L. Fedushkin, A. A. Fagin, T. V. Petrovskaya, J. W. Ziller, R. N. R. Broomhill-Dillard, W. J. Evans,

- [16] M. D. Walter, D. Bentz, F. Weber, O. Schmitt, G. Wolmershäuser, H. Sitzmann, *New J. Chem.* **2007**, 31, 305–318.
- [17] L. Maron, E. L. Werkema, L. Perrin, O. Eisenstein, R. A. Andersen, *J. Am. Chem. Soc.* **2005**, 127, 279–292.
- [18] W. J. Evans, M. A. Horzbor, *J. Organomet. Chem.* **1987**, 326, 299–306.
- [19] K. B. Yatsimirskii, N. K. Davidenko, *Coord. Chem. Rev.* **1979**, 27, 223–273.
- [20] D. S. McClure, Z. Kiss, *J. Chem. Phys.* **1963**, 39, 3251–3257.
- [21] H. Mauermann, P. N. Swepston, T. J. Marks, *Organometallics* **1985**, 4, 200–202.
- [22] a) G. Guillemot, B. Castellano, T. Prange, E. Solari, C. Floriani, *Inorg. Chem.* **2007**, 46, 5152–5154; b) C. D. Berube, M. Yazdanbakhsh, S. Gambarotta, G. P. A. Yap, *Organometallics* **2003**, 22, 3742–3747.
- [23] a) W. J. Evans, G. Zucchi, J. W. Ziller, *J. Am. Chem. Soc.* **2003**, 125, 10–11; b) W. J. Evans, D. S. Lee, D. B. Rego, J. M. Perotti, S. A. Kozimor, E. K. Moore, J. W. Ziller, *J. Am. Chem. Soc.* **2004**, 126, 14574–14582; c) W. J. Evans, D. S. Lee, C. Lie, J. W. Ziller, *Angew. Chem.* **2004**, 116, 5633–5635; *Angew. Chem. Int. Ed.* **2004**, 43, 5517–5519.
- [24] M. Niemeyer, *Z. Anorg. Allg. Chem.* **2002**, 628, 647–657.
- [25] J. M. Smith, A. R. Sadique, T. R. Cundari, K. R. Rodgers, G. Lukat-Rodgers, R. J. Lachicotte, C. J. Flaschenriem, J. Vela, P. L. Holland, *J. Am. Chem. Soc.* **2006**, 128, 756–769.
- [26] Raman spectroscopy has to date received little attention in organolanthanide chemistry. See for example: H.-D. Amberger, H. Reddmann, *Z. Anorg. Allg. Chem.* **2008**, 634, 1542–1554.
- [27] W. J. Evans, D. B. Rego, J. W. Ziller, *Inorg. Chem.* **2006**, 45, 10790–10798.