


Palladium-Catalyzed Three-Component Cascade Cyclization Reaction of Bisallenenes with Propargylic Carbonates and Organoboronic Acids: Efficient Construction of *cis*-Fused Bicyclo[4.3.0]nonenes**

Wei Shu, Guochen Jia, and Shengming Ma*

The palladium-catalyzed coupling reaction involving propargylic/allenic metallic reagents provides one of the most straightforward pathways for the formation of allenes.^[1–3] In this area, we have noticed that the palladium-catalyzed coupling reaction of propargylic carbonates with boronic acids or organozinc reagents affords alkynes or allenes depending on the steric hindrance of the starting materials.^[4] Moreover, multicomponent cascade reactions (MCRs) are among the most efficient synthetic methods for the construction of complex organic molecules.^[5] Such processes avoid the time-consuming and costly protection–deprotection steps as well as purification processes and thus are inherently environmentally benign and atom economical.^[6] We envisioned that a three-component reaction^[7] of 2-alkynyl carbonates, bisallenenes, and organoboronic acids would provide an efficient route for the synthesis of vinyl-substituted allenes **4A** or **4B** via the intermediate allylic palladium species **M3-A**^[8] or the vinylic palladium species **M3-B**^[9] formed from the two consecutive carbopalladations (Scheme 1). Herein, we report our observation that this three-component reaction afforded *cis*-fused bicyclo[4.3.0]nonenes highly regio- and stereoselectively with high efficiency, in which only the vinylic **M3-B** intermediate was involved in forming the first five-membered ring, and the newly formed allene moiety was further carbopalladated subsequently to form the second six-membered ring.



Scheme 1. Possible pathways for the palladium(0)-catalyzed reaction of bisallenenes with propargylic carbonates and boronic acids.

Our initial efforts were based on the reaction of propargylic carbonate **1a**, phenyl boronic acid **2a**, and bisallene **3a**. When the reaction was carried out in dichloroethane (DCE) at 80 °C catalyzed by 5 mol % Pd(OAc)₂ and 10 mol % tri(2'-furyl)phosphine (TFP) in the presence of two equivalents Na₂CO₃, the bicyclic conjugate triene *cis*-**5a** was obtained in 58 % yield (entry 1, Table 1). The structure and relative configuration of this product was confirmed unambiguously by X-ray diffraction (Figure 1).^[10] The product could not be obtained in the absence of K₂CO₃ (entry 2, Table 1). The reaction proceeded better at 90 °C, affording *cis*-**5a** in 64 % yield (entry 3, Table 1). The catalyst is also important for this reaction: it was noted that [Pd(dba)₂]/TFP (dba = *trans,trans*-dibenzylidenacetone) is better than the Pd(OAc)₂/TFP combination (compare entries 3 and 4, Table 1). In addition, the reaction did not proceed when 1,2-bis(diphenylphosphanyl)ethane (dppe) was used as the ligand (entry 5, Table 1). After screening the effect of solvent (entries 6–10, Table 1), we defined **3a**, 1.2 equivalents **1a**, and 2.0 equivalents **2a** catalyzed by 5 mol % [Pd(dba)₂] and 10 mol % TFP in the presence of 2.0 equivalents Na₂CO₃ in DCE at 90 °C as the standard conditions. Under the standard reaction conditions, only the *cis* diastereoisomer was detected, and the formation of 4-type allenes **4A** and **4B** or bicyclic products **5a'** and **5a''** did not occur, as judged by ¹H NMR spectroscopic analysis of the crude reaction mixture.

With the optimized conditions in hand, we turned to examine the substrate scope of the reaction. The reaction is

[*] W. Shu, Prof. Dr. S. Ma

State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences 354 Fenglin Lu, Shanghai 200032 (P.R. China)
Fax: (+86) 21-6416-7510
E-mail: masm@mail.sioc.ac.cn


Prof. Dr. G. Jia

Department of Chemistry
The Hong Kong University of Science and Technology
Clear Water Bay, Kowloon, Hong Kong (P.R. China)

[**] Financial support from the National Natural Science Foundation of China (20429201 and 20732005), the State Basic and Research Development Program (Grant No. 2006CB806105), and the Chinese Academy of Sciences is greatly appreciated. We also thank Xiongdong Lian for reproducing the results of entries 1, 2, and 8 in Table 2.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200805422>.

Table 1: Optimization of the reaction conditions for the Pd-catalyzed three-component cyclization of **3a** with **1a** and **2a**.^[a]



Entry	Catalyst	Base	Solvent	T [°C]	Yield [%] ^[b]
1	Pd(OAc) ₂ /TFP	Na ₂ CO ₃	DCE	80	58
2	Pd(OAc) ₂ /TFP	—	DCE	80	trace ^[c]
3	Pd(OAc) ₂ /TFP	Na ₂ CO ₃	DCE	90	64
4	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	DCE	90	70
5	[Pd(dba) ₂]/dppe	Na ₂ CO ₃	DCE	90	n.r. ^[d]
6	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	DMSO	90	trace ^[c]
7	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	CB	90	49
8	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	dioxane	90	19
9	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	toluene	90	24
10	[Pd(dba) ₂]/TFP	Na ₂ CO ₃	CH ₃ NO ₂	90	trace ^[c]

[a] The reaction was carried out with 0.2 mmol **3a**, 0.24 mmol **1a**, and 0.4 mmol **2a** at the indicated temperature. CB = chlorobenzene, DCE = dichloroethane, DMSO = dimethyl sulfoxide, dppe = 1,2-bis(diphenylphosphanyl)ethane, TFP = tri(2'-furyl)phosphine, Ts = toluene-4-sulfonyl. [b] Yield of isolated **5a**. [c] The starting material **3a** was completely consumed. [d] n.r. = no reaction.





Figure 1. ORTEP plot of *cis*-**5a** shown with ellipsoids at the 30% probability level.

very general for bisallenes **3**, propargylic carbonates **1**, and boronic acids **2**, affording *cis*-fused bicyclo[4.3.0]nonenes **5** in good yields (Tables 2 and 3). Phenyl boronic acids and phenyl boronic acids with electron-withdrawing and electron-donating substituents are all good substrates (entries 1–7, Table 2); polysubstituted phenylboronic acids are also suitable for this MCR process (entries 8 and 9, Table 2). However, the

Table 2: The coupling–cyclization of **3a** and **1a** with various boronic acids **2**.^[a]




Entry	R	t [h]	Yield of 5 [%] ^[b]
1	C ₆ H ₅	3	70 (5a)
2	4-ClC ₆ H ₄	5	63 (5b)
3	4-MeC ₆ H ₄	3	70 (5c)
4	4-MeCOC ₆ H ₄	5	64 (5d)
5	3-MeOC ₆ H ₄	6	66 (5e)
6	4-C ₆ H ₅ C ₆ H ₄	3	65 (5f)
7	2-MeOC ₆ H ₄	4	65 (5o)
8	benzo[d][1,3]dioxol-5-yl	3	65 (5g)
9	3,5-Me ₂ C ₆ H ₃	3	77 (5h)

[a] The reaction was carried out with 0.2 mmol **3a**, 0.24 mmol **1a**, 0.4 mmol **2**. [b] Yield of isolated product.

reaction with terminal methyl 2-methylbut-3-yn-2-yl carbonate does not work, which may be because the presence of Na₂CO₃ is problematic for the acidic terminal alkynic proton.

Internal propargylic carbonates with various substituents in the α and γ positions worked equally well under the standard conditions (entries 1, 4, and 5, Table 3). Moreover,

Table 3: The coupling–cyclization reaction of various propargylic carbonates **1**, phenyl boronic acid **2a**, and bisallenes **3**.^[a]




Entry	R ¹	R ²	X	t [h]	Yield of 5 [%] ^[b]
1	Me	<i>n</i> Bu	NTs	3	73 (5i)
2	Me	<i>n</i> Bu	C(E ¹) ₂ ^[c]	4	67 (5j)
3	Me	<i>n</i> Bu	C(E ²) ₂ ^[c]	4	61 (5k)
4	Me	4-BrC ₆ H ₄	NTs	5	68 (5l)
5	–(CH ₂) ₄ –	Ph	NTs	3	67 (5m)

[a] The reaction was carried out with 0.2 mmol **3**, 0.24 mmol **1**, and 0.4 mmol **2a**. [b] Yield of isolated product. [c] E¹ = CO₂Bn, E² = SO₂Ph.


with differently tethered bisallenes as the substrates, bicyclic compounds with a five-membered carbo- or azacyclic ring with conjugate trienes were obtained in good yields (Table 3).

The most notable feature is that when secondary propargylic carbonate **1e** was used as the substrate, only the *cis-E* diastereoisomer referring to the substituted *exo* C=C bond, that is, *cis*-(*E*)-**5n**, was formed (Scheme 2), which was confirmed by ¹H NMR spectroscopy of the crude reaction mixture and a NOE study of this product.

A rationale that accounts for the observed results is shown in Scheme 3. First, the oxidative addition of palladium(0) with the propargylic carbonate **1** would generate the 1,2-allenyl



Scheme 2. The reaction of **3a** with **2b** and **1e**.



Scheme 3. A plausible mechanism for the formation of **5**.

palladium species **M1**. Subsequent carbopalladation of 1,5-bisallene **3** with **M1** would form the delocalized π -allylic species *syn*-**M2**,^[8] which intramolecularly undergoes carbometallation to afford the vinyl palladium species *cis*-**M3-B** highly stereoselectively.^[9] The *cis* stereoselectivity may be explained by the coordination of the palladium center with the allene moiety in *cis*-**M3-B**. Intramolecular carbopalladation of *cis*-**M3-B** involving the in situ formed allene moiety yields again π -allylic species **M4** or **M4'**. The stereoselectivity observed for the formation of *cis*-(*E*)-**5n** may be explained by the fact that π -allylic intermediate **M4'** would easily isomerize to the more stable **M4** to avoid the steric repulsion between the Pd center and the R^1 group ($R^1 \gg R^2$) by a π - σ - π process.^[11] **M4** would further isomerize to **M5** because of the presence of the "upper" *exo* C=C bond. The final product **5** is subsequently formed by the Suzuki-type coupling of **M5** with the organoboronic acid **2**. It should be noted that the direct coupling product of propargylic carbonate and arylboronic acid is not serious (less than 5%), which may be explained by the rapid reaction between the bisallene and 1,2-allenylpalladium.

In conclusion, we have developed an efficient palladium(0)-catalyzed three-component coupling cyclization reaction of 1,5-bisallene with propargylic carbonate in the presence of organoboronic acid. This reaction may involve three carbopalladation reactions to form sequentially a π -allylic palladium, a vinylic palladium, and a π -allylic palladium intermediate. The interesting *cis*-bicyclo[4.3.0]nonene skeleton in the final product exists in some bioactive molecules, such as the analogue of asparvenone.^[12] Further studies in this area are continuing in our laboratory.

Experimental Section

Typical procedure for the preparation of *cis*-5a: To a Schlenk tube containing Na_2CO_3 (42 mg, 0.40 mmol), $[Pd(dba)_3]$ (6 mg, 0.01 mmol, 5 mol %), and tri(2'-furyl)phosphine (5 mg, 0.02 mmol, 10 mol %) were added sequentially **3a** (56 mg, 0.20 mmol), **2a** (49 mg, 0.4 mmol), **1a** (52 mg, 0.24 mmol), and dichloroethane (2 mL). Then, the resulting mixture was stirred at 90 °C. When the reaction was complete, as monitored by TLC, the solvent was evaporated under reduced pressure. Flash chromatography on silica gel (eluent: petroleum ether/ethyl acetate = 10:1) afforded *cis*-**5a** (71 mg, 70%) as a colorless solid; m.p. 126–127 °C (ethyl acetate/petroleum ether); 1H NMR (300 MHz, $CDCl_3$): δ = 7.63 (d, J = 7.8 Hz, 2H), 7.32–7.14 (m, 10H), 7.00 (d, J = 7.8 Hz, 2H), 4.91 (s, 1H), 4.76 (s, 1H), 3.69–3.61 (m, 2H), 3.50 (t, J = 8.7 Hz, 1H), 3.29 (d, J = 15.0 Hz, 1H), 3.06–2.97 (m, 1H), 2.77–2.63 (m, 2H), 2.52 (t, J = 9.6 Hz, 1H), 2.41 (s, 3H), 1.52 (s, 3H), 1.02 ppm (s, 3H); ^{13}C NMR (75.4 MHz, $CDCl_3$): δ = 148.8, 143.4, 141.4, 140.1, 139.5, 134.9, 133.2, 132.0, 131.1, 129.3, 128.9, 128.6, 128.4, 128.0, 127.9, 126.4, 126.2, 110.0, 54.6, 52.8, 45.0, 42.9, 38.6, 22.0, 21.44, 21.42 ppm; MS(EI): m/z (%) 495 (M^+ , 3.24), 91 (100); IR (neat): $\tilde{\nu}$ = 1640, 1598, 1492, 1473, 1452, 1348, 1304, 1163, 1094, 1029 cm^{-1} ; elemental analysis calcd (%) for $C_{32}H_{33}NO_2S$: C 77.54, H 6.71, N 2.83; found: C 77.34, H 6.80, N 2.99.

Received: November 6, 2008

Published online: March 11, 2009

Keywords: allenes · cyclization · multicomponent reactions · palladium · tandem reactions

- [1] For reviews, see: a) J. Tsuji, T. Mandai, *Angew. Chem.* **1995**, *107*, 2830; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2589; b) J. Tsuji, T. Mandai, *Metal-Catalyzed Cross-Coupling Reactions* (Eds.: F. Diedrich, P. J. Stang), Wiley-VCH, New York, **1998**, p. 455.
- [2] K. M. Brummond, J. E. DeForrest, *Synthesis* **2007**, 795.
- [3] For the most recent examples, see: a) K. Lee, P. H. Lee, *Org. Lett.* **2008**, *10*, 2441; b) M. J. Campbell, P. D. Pohlhaus, G. Min, K. Ohmatsu, J. S. Johnson, *J. Am. Chem. Soc.* **2008**, *130*, 9180; c) T. Miura, M. Shimada, S. Y. Ku, T. Tamai, M. Murakami, *Angew. Chem.* **2007**, *119*, 7231; *Angew. Chem. Int. Ed.* **2007**, *46*, 7101; d) X. Pu, J. M. Ready, *J. Am. Chem. Soc.* **2008**, *130*, 10874.
- [4] S. Ma, A. Zhang, *J. Org. Chem.* **2002**, *67*, 2287.
- [5] For reviews on multicomponent reactions, see: a) J. Zhu, H. Bienaymé, *Multicomponent Reactions*, Wiley-VCH, Weinheim, **2005**; b) D. J. Ramón, M. Yus, *Angew. Chem.* **2005**, *117*, 1628; *Angew. Chem. Int. Ed.* **2005**, *44*, 1602; c) V. Nair, C. Rajesh, A. U. Vinod, S. Bindu, A. R. Sreekanth, J. S. Mathen, L. Balagopal, *Acc. Chem. Res.* **2003**, *36*, 899; d) I. Ugi, *Pure Appl. Chem.* **2001**, *73*, 187; e) A. Dömling, I. Ugi, *Angew. Chem.* **2000**, *112*, 3300; *Angew. Chem. Int. Ed.* **2000**, *39*, 3168; f) A. Dömling,

- Curr. Opin. Chem. Biol.* **2000**, *4*, 318; g) A. Dömling, *Chem. Rev.* **2006**, *106*, 17.
- [6] a) B. M. Trost, *Angew. Chem.* **1995**, *107*, 285; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 259; b) P. A. Wender, S. T. Handy, D. L. Wright, *Chem. Ind.* **1997**, 765.
- [7] Reports on palladium-catalyzed three-component reactions of allenes with: a) acyl chlorides and diborons: F. Y. Yang, M. Y. Wu, C. H. Cheng, *J. Am. Chem. Soc.* **2000**, *122*, 7122; b) organic halides and arylboronic acids: T. H. Huang, H. M. Chang, M. Y. Wu, C. H. Cheng, *J. Org. Chem.* **2002**, *67*, 99.
- [8] The most commonly observed pathway for carbopalladation of allenes is the formation of the delocalized π -allyl palladium species; see: a) "Carbopalladation of Allenes": S. Ma in *Handbook of Organopalladium Chemistry for Organic Synthesis* (Eds.: E. Negishi, A. de Meijere), Wiley-Interscience, New York, **2002**, p. 1491; b) S. Ma, *Pure Appl. Chem.* **2006**, *78*, 197.
- [9] Reports on the formation of vinylic palladium intermediates from the carbopalladation of allenes are very limited; see: a) W. Oppolzer, A. Pimm, B. Stammen, W. E. Hume, *Helv. Chim. Acta* **1997**, *80*, 623; b) R. Grigg, R. Rasul, J. Redpath, D. Wilson, *Tetrahedron Lett.* **1996**, *37*, 4609; c) J. Böhmer, R. Grigg, J. D. Marchbank, *Chem. Commun.* **2002**, 768.
- [10] Crystal data for **5a**: $C_{32}H_{33}NO_2S$, MW=495.65, monoclinic, space group $P2_1/n$, final R indices [$I > 2\sigma(I)$], $R1=0.0617$, $wR2=0.1460$; R indices (all data): $R1=0.1035$, $wR2=0.1637$; $a=8.3858(8)$, $b=10.7904(11)$, $c=30.287(3)$ Å, $\beta=92.252(2)^\circ$, $V=2738.4(5)$ Å³, $T=293(2)$ K, wavelength: 0.71073 Å, $Z=4$, reflections collected/unique: 15 811/5960 ($R_{int}=0.1179$); number of observations [$> 2\sigma(I)$] 3478, parameters: 329. CCDC 703891 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [11] a) C. G. Frost, J. Howarth, J. M. J. Williams, *Tetrahedron: Asymmetry* **1992**, *3*, 1089; b) S. Ma, Z. Gu, *J. Am. Chem. Soc.* **2005**, *127*, 6182.
- [12] M. Bös, H. Stadler, J. Wichmann, F. Jenck, J. R. Martin, J.-L. Moreau, A. J. Sleight, *Helv. Chim. Acta* **1998**, *81*, 525.