


Nonmetal-Mediated Fragmentation of P₄: Isolation of P₁ and P₂ Bis(carbene) Adducts**

Olivier Back, Glenn Kuchenbeiser, Bruno Donnadieu, and Guy Bertrand*

Transition metals are well-known for activating small molecules and for stabilizing highly reactive species. A recent trend in the accomplishment of either of these goals is the use of nonmetals,^[1] especially stable singlet carbenes. Robinson and co-workers have reported that N-heterocyclic carbenes (NHCs) give rise to stable adducts with HBBH^[2] and Si₂,^[3] which are otherwise not isolable.^[4] We have shown that cyclic (alkyl)(amino)carbenes (CAACs) can activate CO,^[5] H₂,^[6] and even NH₃,^[7] which is difficult when attempted with transition-metal centers.^[7,8]

White phosphorus (P₄) is a small molecule that is of industrial interest as it is the classical starting material for the large-scale preparation of organophosphorus derivatives.^[9] The reactivity of P₄ with transition metals has been widely studied,^[10–12] and is therefore an excellent model to further test if carbenes can undergo reactions in the same manner as transition metals. Our research group has already shown that the bulky rigid CAAC **1** opens P₄ and simultaneously stabilizes the resulting acyclic P₄ species (**A**,^[13a] Scheme 1),




Scheme 1. Activation and aggregation of white phosphorus by singlet carbenes. Dipp = 2,6-diisopropylphenyl.

which is otherwise highly reactive. Moreover, when a bulky NHC **2** was used, the NHC-stabilized P₁₂ cluster **B** was isolated in high yield.^[13b] Therefore, singlet carbenes can activate and induce the aggregation of white phosphorus and stabilize the resulting species, in a similar manner to transition metals. However, the most synthetically useful organophosphorus derivatives contain only one or two phosphorus atoms, and therefore it is of primary importance to induce the

fragmentation of P₄, which can be accomplished by using transition metals.^[10] Cummins et al. have even shown that the resulting P₁– and P₂–niobium complexes can be used as phosphorus transfer agents.^[14]

Herein we report that carbenes can induce the fragmentation of white phosphorus in the same manner as transition metals. Depending on the nature of the activator, carbene-stabilized P₂ and P₁ species can be isolated. Preliminary mechanistic studies of the reaction that leads to **A** and **B**, in particular trapping experiments, showed that both carbenes **1** and **2** first formed an unstable monocarbene adduct of type **C**. A second carbene molecule subsequently induces a ring-opening reaction and the formation of bis(carbene) adducts of type **A**.^[13] The different outcomes of the reactions shown in Scheme 1 can be rationalized by the different electronic properties of CAACs and NHCs. CAACs are more electrophilic (π acceptor) and strengthen the PC bonds of **A**,^[5,6] while NHCs are less basic and are therefore better leaving groups, and favor the formation of clusters such as **B**. This analysis indicates that strongly basic but electrophilic carbenes should be the best candidates to induce the fragmentation of white phosphorus, provided that they are small enough to further attack the P₄ fragments of adducts of type **A** or **C**.

According to our previous investigations, the acyclic (alkyl)(amino)carbene **3**^[15] (Scheme 2) is one of the most



Scheme 2. Reaction of white phosphorus with acyclic (alkyl)(amino)carbene **3**.

electrophilic stable carbenes known. An excess of carbene **3** (3.5 equivalents) was added to a suspension of white phosphorus in ether. After two hours at room temperature and subsequent workup, the bis(carbene) P₄ adduct **D** was isolated as light-yellow crystals in 66% yield (based on P₄), and its bis(carbene) P₄ adduct structure was unambiguously assigned by a single crystal X-ray diffraction study^[16] (Figure 1). This bicyclic compound clearly results from the cycloaddition of carbene **3** onto the adduct of type **C**.

This result indicates that the acyclic (alkyl)(amino)carbene **3** is sufficiently nucleophilic to open the P₄ tetrahedron, but so electrophilic that it undergoes a cyclopropanation reaction, rather than inducing the ring-opening of the initially formed adduct **C**. Therefore, we chose the cyclohexyl CAAC **4**,^[17] in the hope that such a small CAAC would be able to induce the formation of an adduct of type **A**, but then react again and induce the fragmentation of the P₄ chain. Upon addition of three equivalents of CAAC **4** to an

[*] O. Back, G. Kuchenbeiser, B. Donnadieu, Prof. G. Bertrand
UCR-CNRS Joint Research Chemistry Laboratory (UMI 2957),
Department of Chemistry, University of California
Riverside, CA 92521-0403 (USA)
Fax: (+1) 951-827-2725
E-mail: guy.bertrand@ucr.edu
Homepage: <http://research.chem.ucr.edu/groups/bertrand/guybertrandwebpage/>

[**] Financial support from the NSF (CHE 0808825) and Rhodia Inc. are gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200902344>.





Figure 1. Thermal ellipsoids plot (40% probability surface) for **D** (hydrogen atoms omitted for clarity). Selected bond distances [Å] and angles [°]: P(1)–P(2) 2.2320(19), P(2)–P(3) 2.220(2), P(2)–P(4) 2.2342(17), P(3)–P(4) 2.1700(19), C(1)–P(1) 1.747(2), C(18)–P(3) 1.912(2), C(18)–P(4) 1.925(2), N(1)–C(1) 1.3893(19), N(2)–C(18) 1.4472(18); P(1)–P(2)–P(3) 93.29(3), P(1)–P(2)–P(4) 91.99(6), P(3)–P(2)–P(4) 58.31(7), P(2)–P(4)–P(3) 60.51(5), P(3)–P(4)–P(2) 60.51(5).

suspension of P_4 in ether, two new products **E** and **F** were isolated in moderate yields (67% and 12%, respectively, based on P_4 ; Scheme 3).



Scheme 3. Reaction of white phosphorus with the nonhindered CAAC **4**.

The ^{31}P NMR spectrum of the major product **E** shows a doublet signal at $\delta = +68.1$ ppm and a quartet signal at $\delta = -66.2$ ppm, with a large P–P coupling constant (227 Hz). These data indicated the presence of three magnetically equivalent P nuclei that were directly connected to another P center. This hypothesis was confirmed by a single crystal X-ray diffraction study (Figure 2). Very surprisingly, the for-





Figure 2. Thermal ellipsoids plot (40% probability surface) for **E** (hydrogen atoms omitted for clarity). Selected bond distances [Å] and angles [°]: P(1a)–P(2a) 2.2168(6), P(1a)–P(3a) 2.2270(6), P(1a)–P(4a) 2.2192(7), C(1a)–P(2a) 1.7328(19), C(24a)–P(3a) 1.7343(18), C(47a)–P(4a) 1.7324(18), N(1a)–C(1a) 1.374(2), N(2a)–C(24a) 1.370(2), N(3a)–C(47a) 1.368(2); P(2a)–P(1a)–P(4a) 90.15(2), P(3a)–P(1a)–P(4a) 90.15(2), P(2a)–P(1a)–P(4a) 90.15(2), C(1a)–P(2a)–P(1a) 108.22(6), C(24a)–P(3a)–P(1a) 106.62(6), C(47a)–P(4a)–P(1a) 108.45(7).

mation of tris(carbene) P_4 adduct **E** implies that two CAACs **4** react on the initially formed adduct of type **C**.

The minor product **F** was determined to be the desired bis(carbene) P_2 adduct, and therefore the reaction that leads to **F** represents the first example of fragmentation of P_4 with a neutral organic species. Interestingly, compound **F** is reminiscent of compounds **F'**^[18] and **F''**,^[19] which were prepared as shown in Scheme 4, and characterized as 2,3-diphosphabutadiene and NHC-stabilized bis(phosphinidene), respectively.



Scheme 4. Reported syntheses^[18,19] and electronic structure of compounds related to **F**.

The ^{31}P NMR chemical shift for **F** ($\delta = +59.4$ ppm) is comparable to that of **F'** ($\delta = 54.2$ ppm), and is at a lower field than **F''** ($\delta = -52.4$ ppm); similarly, the P–C bond length (Figure 3) in **F** (1.719(7) Å) is shorter than in **F''** (1.750(2) Å). Both of these data indicate that **F** has a 2,3-diphosphabutadiene structure, which is expected because of the higher electrophilicity of CAACs than NHCs.





Figure 3. Thermal ellipsoids plot (40% probability surface) for **F** (hydrogen atoms omitted for clarity). Selected bond distances [Å] and angles [°]: Selected bond distances [Å] and angles [°]: P(1a)–P(2a) 2.184(3), C(1a)–P(1a) 1.719(7), N(1a)–C(1a) 1.387(9); C(1a)–P(1a)–P(2a) 105.1(2); C(1a)–P(1a)–P(2a)–C(24a) 149.22.

Compound **F** most probably results from the attack of the carbene to the β -phosphorus centers of a bis(carbene) adduct of type **A**. In order to obtain a P_1 adduct by attack at a phosphorus center in a position that is α to the carbene(s) of adducts **A** or **C**, we have used the least sterically demanding stable carbene known to date, namely bis(diisopropylamino)cyclopropenylidene (**5**).^[20] The ^{31}P NMR spectrum of the reaction mixture of P_4 and three equivalents of carbene **5** after 12 hours at room temperature revealed an ABX system ($\delta_A = 242.8$, $\delta_B = 237.0$, $\delta_X = 157.0$ ppm, $J_{AX} = -484.7$ Hz, $J_{AB} =$

+ 38.8 Hz, $J_{\text{BX}} = -481.1$ Hz) and a singlet signal at $\delta = -93.2$ ppm. All attempts to purify the mixture led to the disappearance of the second-order system, but, in the presence of chloroform or any salts, the high-field singlet remained unchanged. When chloroform was used, single crystals suitable for X-ray diffraction readily formed upon standing at room temperature. The isolated product was the bis(carbene) P_1 cation **G** with Cl^- as counterion (obtained in 74% yield based on P_4 ; Scheme 5, Figure 4). Analogous



Scheme 5. Reaction of P_4 with cyclopropenylidene **5**.




Figure 4. Thermal ellipsoids plot (40% probability surface) for **GCl**. H atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: P(1)–C(1) 1.787(2), P(1)–C(16) 1.788(2), N(1)–C(2) 1.323(3), N(2)–C(3) 1.328(3), C(1)–C(3) 1.397(3), C(1)–C(2) 1.404(3), C(2)–C(3) 1.388(3); C(1)–P(1)–C(16) 104.68(9).

(NHC) P^+ systems that used (R_3P) P^+ , (RNPCl) $_2$, or PCl_3 as P^+ source have previously been reported by Schmidpeter et al.^[21] and MacDonald et al.^[22] On the other hand, although we have not yet been able to identify the P_3 fragment, its anionic nature is clear, and therefore it is likely to be an allylic triphosphorus anion that is unsymmetrically substituted by cyclopropenylidenes.

The results reported herein demonstrate that reactions of P_4 , namely, activation, aggregation, and importantly, fragmentation, which have previously been carried out by using transition metals, can also be achieved by using stable singlet carbenes. The next challenge is to use the resulting adducts, especially **F** and **G**, to prepare useful organophosphorus derivatives. This procedure would avoid the use of Cl_2 gas, which is important to meet the growing demand for phosphorus derivatives that are produced by using environmentally friendly processes.

Experimental Section

All manipulations were performed under an atmosphere of dry argon by using standard Schlenk techniques.

Synthesis of D: Ether (40 mL) was added to a mixture of carbene **3** (4.07 mmol) and P_4 (1.16 mmol) at room temperature and the

resulting suspension stirred for 2 h. The solvent was removed under vacuum, and the solid residue washed with hexane (6 mL) at -35°C to afford **D** (480 mg, 66.4% based on P_4). Single crystals of **D** were grown by layering acetonitrile over a THF solution. m.p. 150°C ; $^{31}\text{P}\{^1\text{H}\}$ NMR (202.5 MHz, $[\text{D}_8]\text{THF}$, 25°C): $\delta = 238.8$ (dt, $^1J_{\text{PP}} = 220$ Hz and $^2J_{\text{PP}} = 87$ Hz), -105.8 (dt, $^1J_{\text{PP}} = 220$ Hz and $^2J_{\text{PP}} = 167$ Hz), -168.2 ppm (dd, $^1J_{\text{PP}} = 167$ Hz and $^2J_{\text{PP}} = 87$ Hz); ^{13}C NMR (125.75 MHz, $[\text{D}_8]\text{THF}$, 25°C): $\delta = 227.6$ (d, $^1J_{\text{PC}} = 89$ Hz, C_{carb}), 101.5 ppm (br s, C_{cyclo}).

Synthesis of E and F: A solution of carbene **4** (3.25 g, 10.01 mmol) in ether (20 mL) was added at room temperature to a suspension of P_4 (0.41 g, 3.34 mmol) in ether (20 mL). The mixture was stirred at room temperature for 2 h and then half of the solvent was removed under vacuum. The remaining solution was cooled to -30°C and the resulting precipitate filtered to give **F** as a bright yellow powder (580 mg, 12.1% yield based on P_4). Evaporation of the filtrate gave a dark red powder, which was washed three times with acetonitrile (3×30 mL), and dried under vacuum to give **E** as a bright-yellow powder (2.5 g, 67.4% yield based on P_4). Single crystals were grown from a saturated solution of **F** in diethyl ether at room temperature, and single crystals of **E** were obtained by slow evaporation of a solution of **E** in hexane at room temperature. **E**: m.p. 142°C ; ^{13}C NMR (125.75 MHz, $[\text{D}_6]\text{benzene}$, 25°C): $\delta = 207.4$ ppm (m, C_{carb}). **F**: m.p. 216°C ; $^{13}\text{C}\{^1\text{H}\}$ NMR (125.75 MHz, $[\text{D}_6]\text{benzene}$, 25°C): $\delta = 202.2$ ppm (dd, $^1J_{\text{PC}} = 32$ Hz, $^2J_{\text{PC}} = 26$ Hz, C_{carb}).

Synthesis of GCl: White phosphorus (P_4 ; 0.17 g, 1.40 mmol) was added to a solution of carbene **5** (1.00 g, 4.24 mmol) in THF (5 mL). After stirring for 12 h at room temperature, chloroform (5 mL) was added to the crude reaction mixture. After stirring at room temperature for 30 min, the volatile components were removed under vacuum, and the resulting red solid washed with Et_2O (3×20 mL) to afford a dark-orange powder. Yield: 564 mg (74% based on P_4). Single crystals of **GCl** were readily obtained by slow evaporation of a solution of **GCl** in chloroform at room temperature. m.p. 174°C (dec.); ^1H NMR (300 MHz, CDCl_3 , 25°C): $\delta = 3.79$ (sept, $J = 6.5$ Hz, 4H), 1.25 ppm (d, $J = 6.5$ Hz, 24H); $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3 , 25°C): $\delta = 136.5$ (C_{ring}), 123.7 (d, $^1J_{\text{C-P}} = 104.6$ Hz, C_{carb}), 51.3, 21.5 ppm. For compounds **D–G**, only selected NMR data are given. For full details, see the Supporting Information.

Received: May 2, 2009

Published online: June 19, 2009

Keywords: carbenes · cyclopropenylidenes · fragmentation · phosphorus · P–P activation

- [1] For recent examples of small molecule activation by nonmetals other than carbenes, see: a) S. J. Geier, D. W. Stephan, *J. Am. Chem. Soc.* **2009**, *131*, 3476–3477; b) M. Ullrich, A. J. Lough, D. W. Stephan, *J. Am. Chem. Soc.* **2009**, *131*, 52–53; c) S. N. Riduan, Y. Zhang, J. Y. Ying, *Angew. Chem.* **2009**, *121*, 3372–3375; *Angew. Chem. Int. Ed.* **2009**, *48*, 3322–3325; d) A. Jana, C. Schulzke, H. W. Roesky, *J. Am. Chem. Soc.* **2009**, *131*, 4600–4601; e) Z. L. Zhu, X. P. Wang, Y. Peng, H. Lei, J. C. Fetting, E. Rivard, P. P. Power, *Angew. Chem.* **2009**, *121*, 2065–2068; *Angew. Chem. Int. Ed.* **2009**, *48*, 2031–2034; f) Y. Peng, M. Brinda, B. D. Ellis, J. C. Fetting, E. Rivard, P. P. Power, *Chem. Commun.* **2009**, 6042–6044; g) Y. Peng, B. D. Ellis, X. P. Wang, J. C. Fetting, P. P. Power, *J. Am. Chem. Soc.* **2008**, *130*, 12268–12269; h) M. A. Dureen, D. W. Stephan, *Chem. Commun.* **2008**, 4303–4305; i) P. A. Chase, D. W. Stephan, *Angew. Chem.* **2008**, *120*, 7543–7547; *Angew. Chem. Int. Ed.* **2008**, *47*, 7433–7437; j) D. Holschumacher, T. Bannenberg, C. G. Hrib, P. G. Jones, M. Tamm, *Angew. Chem.* **2008**, *120*, 7538–7542; *Angew. Chem. Int. Ed.* **2008**, *47*, 7428–7432; k) P. Spies, S. Schewendemann, S. Lange, G. Kehr, R. Fröhlich, G. Erker, *Angew. Chem.* **2008**, *120*,

- 7654–7657; *Angew. Chem. Int. Ed.* **2008**, *47*, 7543–7546; l) V. Sumerin, F. Schulz, M. Atsumi, C. Wang, M. Nieger, M. Leskela, T. Repo, P. Pyykko, B. Rieger, *J. Am. Chem. Soc.* **2008**, *130*, 14117–14119; m) V. Sumerin, F. Schulz, M. Nieger, M. T. R. Leskela, B. Rieger, *Angew. Chem.* **2008**, *120*, 6090–6092; *Angew. Chem. Int. Ed.* **2008**, *47*, 6001–6003; n) G. C. Welch, R. R. S. Juan, J. D. Masuda, D. W. Stephan, *Science* **2006**, *314*, 1124–1126.
- [2] a) Y. Z. Wang, B. Quillian, P. R. Wei, C. S. Wannere, Y. Xie, R. B. King, H. F. Schaefer, P. v. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.* **2007**, *129*, 12412–12413; b) Y. Z. Wang, B. Quillian, P. R. Wei, Y. Xie, C. S. Wannere, R. B. King, H. F. Schaefer, P. v. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.* **2008**, *130*, 3298–3299.
- [3] Y. Z. Wang, Y. M. Xie, P. R. Wei, R. B. King, H. F. Schaefer, P. von R. Schleyer, *Science* **2008**, *321*, 1069–1071; See also: C. A. Dyker, G. Bertrand, *Science* **2008**, *321*, 1050–1051.
- [4] a) L. B. Knight, K. Kerr, P. K. Miller, C. A. Arrington, *J. Phys. Chem.* **1995**, *99*, 16842–16848; b) A. H. Cowley, *ACS Symp. Ser.* **2006**, *917*, 2–19; c) R. West, *ACS Symp. Ser.* **2006**, *917*, 166–178; d) P. P. Power, *Chem. Rev.* **1999**, *99*, 3463–3503.
- [5] V. Lavallo, Y. Canac, B. Donnadieu, W. W. Schoeller, G. Bertrand, *Angew. Chem.* **2006**, *118*, 3568–3571; *Angew. Chem. Int. Ed.* **2006**, *45*, 3488–3491.
- [6] G. D. Frey, V. Lavallo, B. Donnadieu, W. W. Schoeller, G. Bertrand, *Science* **2007**, *316*, 439–441.
- [7] A few examples of NH_3 activation by a transition metal have been reported, see: a) C. M. Fafard, D. Adhikari, B. M. Foxman, D. J. Mindiola, O. V. Ozerov, *J. Am. Chem. Soc.* **2007**, *129*, 10318–10319; b) T. E. Hanna, E. Loblovski, P. J. Chirik, *Eur. J. Inorg. Chem.* **2007**, 2677–2685; c) Y. Nakajima, H. Kameo, H. Suzuki, *Angew. Chem.* **2006**, *118*, 964–966; *Angew. Chem. Int. Ed.* **2006**, *45*, 950–952; d) J. Zhao, A. S. Goldman, J. F. Hartwig, *Science* **2005**, *307*, 1080–1082.
- [8] Examples of metal-catalyzed hydroamination of alkynes with NH_3 have recently been reported: V. Lavallo, G. D. Frey, B. Donnadieu, M. Soleilhavoup, G. Bertrand, *Angew. Chem.* **2008**, *120*, 5302–5306; *Angew. Chem. Int. Ed.* **2008**, *47*, 5224–5228.
- [9] a) D. E. C. Corbridge, *Phosphorus—An Outline of its Chemistry, Biochemistry and Technology*, 5th ed., Elsevier, Amsterdam, **1995**; b) J. Emsley, *The 13th Element: The Sordid Tale of Murder, Fire, and Phosphorus*, Wiley, New York, **2002**.
- [10] For recent reviews, see: a) J. S. Figueroa, C. C. Cummins, *Dalton Trans.* **2006**, 2161–2168; b) C. C. Cummins, *Angew. Chem.* **2006**, *118*, 876–884; *Angew. Chem. Int. Ed.* **2006**, *45*, 862–870; c) M. Peruzzini, L. Gonsalvi, A. Romero, *Chem. Soc. Rev.* **2005**, *34*, 1038–1047; d) M. Peruzzini, R. R. Abdrimova, Y. Budnikova, A. Romero, O. J. Scherer, H. Sitzmann, *J. Organomet. Chem.* **2004**, *689*, 4319–4331; e) M. Ehse, A. Romero, M. Peruzzini, *Top. Curr. Chem.* **2002**, *220*, 107–140.
- [11] For recent reports, see: a) P. Barbaro, M. Di Vaira, M. Peruzzini, S. S. Costantini, P. Stoppioni, *Inorg. Chem.* **2009**, *48*, 1091–1096; b) W. W. Seidel, O. T. Summerscales, B. O. Patrick, M. D. Fryzuk, *Angew. Chem.* **2009**, *121*, 121–123; *Angew. Chem. Int. Ed.* **2009**, *48*, 115–117; c) P. Barbaro, M. Di Vaira, M. Peruzzini, S. S. Costantini, P. Stoppioni, *Angew. Chem.* **2008**, *120*, 4497–4499; *Angew. Chem. Int. Ed.* **2008**, *47*, 4425–4427; d) M. Caporali, P. Barbaro, L. Gonsalvi, A. Ienco, D. Yakhvarov, M. Peruzzini, *Angew. Chem.* **2008**, *120*, 3826–3828; *Angew. Chem. Int. Ed.* **2008**, *47*, 3766–3768.
- [12] Main-group-mediated activation of P_4 has also been reported; for recent examples, see: a) J. J. Weigand, M. Holthausen, R. Frölich, *Angew. Chem.* **2009**, *121*, 301–304; *Angew. Chem. Int. Ed.* **2009**, *48*, 295–298; b) W. T. K. Chan, F. Garcia, A. D. Hopkins, L. C. Martin, M. McPartlin, D. S. Wright, *Angew. Chem.* **2007**, *119*, 3144–3146; *Angew. Chem. Int. Ed.* **2007**, *46*, 3084–3086; c) Y. Xiong, S. Yao, M. Brym, M. Driess, *Angew. Chem.* **2007**, *119*, 4595–4597; *Angew. Chem. Int. Ed.* **2007**, *46*, 4511–4513; d) A. R. Fox, R. J. Wright, E. Rivard, P. P. Power, *Angew. Chem.* **2005**, *117*, 7907–7911; *Angew. Chem. Int. Ed.* **2005**, *44*, 7729–7733; e) H. W. Lerner, M. Bolte, K. Karaghiosoff, M. Wagner, *Organometallics* **2004**, *23*, 6073–6076.
- [13] a) J. D. Masuda, W. W. Schoeller, B. Donnadieu, G. Bertrand, *Angew. Chem.* **2007**, *119*, 7182–7185; *Angew. Chem. Int. Ed.* **2007**, *46*, 7052–7055; b) J. D. Masuda, W. W. Schoeller, B. Donnadieu, G. Bertrand, *J. Am. Chem. Soc.* **2007**, *129*, 14180–14181. See also: J. M. Lynam, *Angew. Chem.* **2008**, *120*, 843–845; *Angew. Chem. Int. Ed.* **2008**, *47*, 831–833.
- [14] a) B. M. Cossairt, M. C. Diawara, C. C. Cummins, *Science* **2009**, *323*, 602–602; b) B. M. Cossairt, C. C. Cummins, *Angew. Chem.* **2008**, *120*, 8995–8998; *Angew. Chem. Int. Ed.* **2008**, *47*, 8863–8866; c) N. A. Piro, C. C. Cummins, *J. Am. Chem. Soc.* **2008**, *130*, 9524–9535; d) A. R. Fox, C. R. Clough, N. A. Piro, C. C. Cummins, *Angew. Chem.* **2007**, *119*, 991–994; *Angew. Chem. Int. Ed.* **2007**, *46*, 973–976; e) N. A. Piro, J. S. Figueroa, J. T. McKellar, C. C. Cummins, *Science* **2006**, *313*, 1276–1279.
- [15] V. Lavallo, J. Mafhouz, Y. Canac, B. Donnadieu, W. W. Schoeller, G. Bertrand, *J. Am. Chem. Soc.* **2004**, *126*, 8670–8671.
- [16] CCDC 730165 (D), 730166 (E), 730167 (F), and 730168 (G) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [17] a) V. Lavallo, Y. Canac, C. Prasang, B. Donnadieu, G. Bertrand, *Angew. Chem.* **2005**, *117*, 5851–5855; *Angew. Chem. Int. Ed.* **2005**, *44*, 5705–5709; b) R. Jazsar, R. D. Dewhurst, J. B. Bourg, B. Donnadieu, Y. Canac, G. Bertrand, *Angew. Chem.* **2007**, *119*, 2957–2960; *Angew. Chem. Int. Ed.* **2007**, *46*, 2899–2902.
- [18] V. D. Romanenko, L. S. Kachkovskaya, L. N. Markovskii, *Zh. Obshch. Khim.* **1985**, *55*, 2140–2141.
- [19] Y. Z. Wang, Y. M. Xie, P. R. Wei, R. B. King, H. F. Schaefer, P. von R. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.* **2008**, *130*, 14970–14971.
- [20] V. Lavallo, Y. Canac, B. Donnadieu, W. W. Schoeller, G. Bertrand, *Science* **2006**, *312*, 722–724.
- [21] A. Schmidpeter, S. Lochschmidt, A. Willhalm, *Angew. Chem.* **1983**, *95*, 561–562; *Angew. Chem. Int. Ed. Engl.* **1983**, *22*, 545–546.
- [22] B. D. Ellis, C. A. Dyker, A. Decken, C. L. B. Macdonald, *Chem. Commun.* **2005**, 1965–1967.