


## Expedient Synthesis of (–)-Amphidinolide X\*\*

Jae Hoon Jung and Eun Lee\*


(–)-Amphidinolide X (**1**, Scheme 1) is one of the cytotoxic macrolides isolated by Kobayashi and co-workers from the laboratory-cultured dinoflagellates *Amphidinium* sp., which are symbionts of the Okinawan marine flatworms *Amphisco-*



**Scheme 1.** Retrosynthetic analysis. Tol: tolyl.

*lops* sp.<sup>[1]</sup> Amphidinolide X (**1**) is known to possess cytotoxic activity with IC<sub>50</sub> values of 0.6 and 7.5 μg mL<sup>−1</sup> against murine lymphoma L1210 and human epidermoid carcinoma KB cells, respectively, in vitro. It is the only naturally occurring macrodiolide known to date that consists of a diacid and a diol unit, and it has been a popular target for synthetic studies because of its scarcity and unique structure.<sup>[2]</sup> The most characteristic feature in the structure of **1** is the substituted 3-hydroxyoxolane structure, which is derived from a tertiary alcohol. Construction of this type of motif<sup>[3]</sup> normally utilizes the nucleophilicity of hydroxy groups, and indeed, hydroxy addition to allenes,<sup>[2a,b]</sup> epoxides,<sup>[2c,g]</sup> or olefins<sup>[2f]</sup> is a key step in the synthesis of **1**. We intended to obtain the key structural element **A** through 5-*exo* cyclization of aldehyde β-alkoxyvinyl sulfoxide **B** (Scheme 1). It is known<sup>[4]</sup> that 5-*exo* cyclizations of aldehyde β-alkoxyvinyl sulfoxides derived from secondary alcohols proceed under carbinol chirality control, and *cis*-2,5-disubstituted oxolanes are formed regardless of the sulfoxide chirality. For (*E*)- and (*Z*)-β-alkoxyvinyl sulfoxides derived from tertiary alcohols, will the reactions proceed under sulfoxide or carbinol chirality control?

In practice, the known (3*R*)-epoxide *p*-methoxyphenyl (PMP) ether **2**<sup>[5]</sup> was converted into the (3*R*)-3-methylhexan-1,3-diol PMP ether **3** (Scheme 2). Reaction of **3** with alkynyl



**Scheme 2.** Synthesis of the β-alkoxyvinyl sulfoxides. a) EtMgBr, CuI, THF; b) **4**, EtMgBr, LiCl, THF; c) **5**, EtMgBr, LiCl, THF; d) CAN, MeCN/H<sub>2</sub>O (9:1); e) I<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, THF; tetrahydrofuran; CAN: ceric ammonium nitrate.

sulfoxides **4** or **5** in the presence of ethylmagnesium bromide and lithium chloride led to (*Z*)-β-alkoxyvinyl sulfoxides **6** or **7**, respectively. The crucial vinyl ether formation did not proceed in the presence of other bases.<sup>[6]</sup> PMP deprotection of **6** through CAN oxidation provided (*Z*)-β-alkoxyvinyl sulfoxide **10**. When **6** was treated with iodine before the CAN oxidation, (*E*)-β-alkoxyvinyl sulfoxide **8** was obtained. By the same sequence of reactions, (*Z*)- and (*E*)-β-alkoxyvinyl sulfoxides **9** and **11** were prepared from **7**.


Dess–Martin oxidation of (*E*),(*S*)-β-alkoxyvinyl sulfoxide **8** and then reaction with SmI<sub>2</sub> in the presence of methanol yielded a 9.2:1 mixture of the 3-hydroxyoxolane products, with the major product being **12**, which was isolated in 67% yield (Scheme 3). Through the same reaction sequence, a 15.4:1 mixture with **13** as the favored product (62% yield) was obtained from the *Z*,*R* isomer **9**. Likewise, the *Z*,*S* isomer **10** was converted into a 5.7:1 mixture with **14** as the favored product (62% yield). When the *E*,*R* isomer **11** was subjected to the same reaction conditions, a 3.3:1 mixture of products was obtained with **15** as the major product, which was isolated in 73% yield. The minor products were epimeric carbinols.<sup>[7]</sup> *m*-CPBA oxidation of sulfoxides **12–15** produced four different sulfones **16–19**.<sup>[8]</sup>

The SmI<sub>2</sub>-mediated 5-*exo* cyclization reactions of aldehyde β-alkoxyvinyl sulfoxides **8–11** derived from tertiary

[\*] J. H. Jung, Prof. Dr. E. Lee  
Department of Chemistry, College of Natural Sciences  
Seoul National University, Seoul 151-747 (Korea)  
Fax: (+82) 2-889-1568  
E-mail: eunlee@snu.ac.kr

[\*\*] This work was supported by a grant from the Marine Biotechnology Program funded by the Ministry of Land, Transport and Maritime Affairs, Republic of Korea, and a grant from the Center for Bioactive Molecular Hybrids (Yonsei University and KOSEF). A BK21 graduate fellowship grant and a Seoul Science Fellowship grant to J.H.J. are gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200900865>.




**Scheme 3.**  $\text{SmI}_2$ -mediated cyclization of aldehyde  $\beta$ -alkoxyvinyl sulfoxides. a) DMP,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$ ; b)  $\text{SmI}_2$ , MeOH, THF,  $0^\circ\text{C}$ ; c)  $m$ -CPBA,  $\text{CH}_2\text{Cl}_2$ . DMP: Dess–Martin periodinane;  $m$ -CPBA:  $m$ -chloroperoxybenzoic acid.


alcohols were stereospecific, and a useful degree of stereoselectivity was achieved. The structures of the major products may be predicted on the basis of the double-bond stereochemistry and the sulfoxide configuration (sulfoxide chirality control). The observed stereospecificity appears to be totally unrelated to the data previously obtained for substrates derived from secondary alcohols.<sup>[4]</sup> The results of  $\text{SmI}_2$ -mediated 5-*exo* cyclizations of aldehyde  $\beta$ -alkoxyvinyl sulfoxides prepared from secondary and tertiary alcohols are presented in Scheme 4.

The results may be explained by proposing “eclipsed lone pair” transition states **C–F** for the formation of the major product in each case.<sup>[9]</sup> In these structures, the samarium ketyl group coordinated to the sulfoxide oxygen atom necessarily approaches the double bond opposite from the bulky aryl group (Scheme 5).

After *p*-nitrobenzoate protection of the hydroxy group, the Pummerer rearrangement of **12** proceeded smoothly to yield aldehyde **20** (Scheme 6). *p*-Nitrobenzoate protection is advantageous, because the minor product produced along with **12** (and **13**) may also be converted into the same ester by a Mitsunobu reaction. *p*-Aminobenzoate ketone **21** was obtained through Horner–Emmons reaction of **20** and hydrogenation. Wittig homologation of ketone **21** and LAH reduction led to the hydroxy olefin **22**. A cross-olefin metathesis reaction between olefin **22** and TBS ether **23**, prepared from a known alcohol,<sup>[24]</sup> proceeded smoothly to produce *E* olefin **24**. The corresponding *Z* olefin was not detected in the product mixture. Reaction of **24** with acryloyl chloride, TBS deprotection, and careful acylation with the acid chloride prepared from the known carboxylic acid **25**<sup>[10]</sup>




**Scheme 4.** Comparison of the cyclization products from  $\beta$ -alkoxyvinyl sulfoxides derived from secondary (the previous study)<sup>[4]</sup> and tertiary alcohols (the present work). Bn: benzyl.



**Scheme 5.** Transition-state structures for the reactions of **8–11**.

produced triene **26**. The key ring-closing olefin metathesis reaction of **26** in the presence of the second-generation Grubbs catalyst was successful and provided (–)-amphidinolide **X** (**1**) in 74% yield, accompanied by the corresponding *Z* isomer in 11% yield.

In summary, the key 3-hydroxyoxolane fragment of (–)-amphidinolide **X** was prepared through  $\text{SmI}_2$ -mediated 5-*exo* cyclization of an aldehyde  $\beta$ -alkoxyvinyl sulfoxide derived from (*R*)-3-hydroxy-3-methylhexanal. Three other possible stereoisomers were also obtained by changing the double-bond stereochemistry and the sulfoxide chirality. In comparison with the results obtained previously,<sup>[4]</sup> a subtle change in the structure of the substrate (from a hydrogen atom to a methyl group) triggered a dramatic change of stereospecificity. The method described in this communication opens up new and rational ways for the preparation of functionalized oxacycles, particularly those derived from



**Scheme 6.** Synthesis of **1**. a) PNBA, DCC, DMAP,  $\text{CH}_2\text{Cl}_2$ ; b) TFAA, pyridine, MeCN,  $0^\circ\text{C}$ ; KOAc,  $\text{H}_2\text{O}$ ; c)  $\text{MeCOCH}_2\text{PO}(\text{OMe})_2$ , DIPEA, MeCN; d)  $\text{H}_2$ , Pd/C, MeOH; e)  $\text{Ph}_3\text{PMe}^+\text{Br}^-$ ,  $n\text{BuLi}$ ,  $-20^\circ\text{C}$ ; f) LAH, diethyl ether; g) **23**, 20 mol %  $(\text{H}_2\text{IMes}_2)(\text{Cy}_3\text{P})\text{Cl}_2\text{RuCHPh}$ ,  $\text{CH}_2\text{Cl}_2$ , reflux; h)  $\text{CH}_2\text{CHCOCl}$ , pyridine,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$ ; i) AcOH,  $\text{H}_2\text{O}$ ,  $60^\circ\text{C}$ ; j)  $(\text{COCl})_2$ , **25**; alcohol, DMAP,  $\text{CH}_2\text{Cl}_2$ ; k) 7 mol %  $(\text{H}_2\text{IMes}_2)(\text{Cy}_3\text{P})\text{Cl}_2\text{RuCHPh}$ ,  $\text{CH}_2\text{Cl}_2$ , reflux. TBS: *tert*-butyldimethylsilyl; PNB: *p*-nitrobenzoyl; PAB: *p*-aminobenzoyl; PNBA: *p*-nitrobenzoic acid; DCC: *N,N'*-dicyclohexylcarbodiimide; DMAP: 4-dimethylaminopyridine; TFAA: trifluoroacetic anhydride; DIPEA: diisopropylethylamine; LAH: lithium aluminum hydride; Mes: mesyl; Cy: cyclohexyl.

tertiary alcohols. Future studies will focus on these bioactive oxacyclic natural products.

Received: February 13, 2009  
Published online: May 7, 2009

**Keywords:** natural products · ring-closing metathesis ·  $\text{Sml}_2$ -mediated cyclization · sulfoxides · total synthesis

- [1] M. Tsuda, N. Izui, K. Shimbo, M. Sato, E. Fukushi, J. Kawabata, K. Katsumata, T. Horiguchi, J. Kobayashi, *J. Org. Chem.* **2003**, 68, 5339–5345.
- [2] For reports on the total synthesis, see: a) O. Lepage, E. Kattnig, A. Fürstner, *J. Am. Chem. Soc.* **2004**, 126, 15970–15971; b) A. Fürstner, E. Kattnig, O. Lepage, *J. Am. Chem. Soc.* **2006**, 128, 9194–9204; c) W.-M. Dai, Y. Chen, J. Jin, J. Wu, J. Lou, Q. He, *Synlett* **2008**, 1737–1741; d) C. Rodríguez-Esrich, F. Urpi, J. Vilarrasa, *Org. Lett.* **2008**, 10, 5191–5194. For related oxacycle syntheses, see: e) W.-M. Dai, Y. Chen, J. Jin, J. Wu, *Synlett* **2006**, 1177–1180; f) C. Rodríguez-Esrich, A. Olivella, F. Urpi, J. Vilarrasa, *Org. Lett.* **2007**, 9, 989–992; g) H. D. Doan, J. Gallon, A. Piou, J.-M. Vatlè, *Synlett* **2007**, 983–985.
- [3] For some of the most recent examples, see: a) I. Calis, A. A. Dönmez, A. Perrone, C. Pizza, S. Piacente, *Phytochemistry* **2008**, 69, 2634–2638; b) N.-Y. Ji, X.-M. Li, H. Xie, J. Ding, K. Li, L.-P. Ding, B.-G. Wang, *Helv. Chim. Acta* **2008**, 91, 1940–1946; c) P. G. Williams, E. D. Miller, R. N. Asolkar, P. R. Jensen, W. Fenical, *J. Org. Chem.* **2007**, 72, 5025–5034. For recent examples of a large number of marine norditerpenoids featuring 3-oxoxolanes derived from tertiary alcohols, see: d) A. F. Ahmed, J.-H. Su, Y.-H. Kuo, J.-H. Sheu, *J. Nat. Prod.* **2004**, 67, 2079–2082.
- [4] J. H. Jung, Y. W. Kim, M. A. Kim, S. Y. Choi, Y. K. Chung, T.-R. Kim, S. Shin, E. Lee, *Org. Lett.* **2007**, 9, 3225–3228.
- [5] C.-L. Chen, S. M. Sparks, S. F. Martin, *J. Am. Chem. Soc.* **2006**, 128, 13696–13697.
- [6] The reaction did not proceed in the presence of *N*-methylmorpholine, lithium hexamethyldisilazide, lithium diisopropylamide, calcium hydride, or cesium carbonate.
- [7] For example, the minor product accompanying **12** was converted into sulfone **17** upon *m*-CPBA oxidation.
- [8] NOESY correlations were used in the structural assignment of **16–19**.
- [9] In the cyclization studies of  $\beta$ -alkoxyvinyl sulfoxides derived from secondary alcohols, transition states were proposed with the R group “equatorial” at the flap of the envelope, which resemble the stable conformation of methylcyclopentane. For discussions on conformations of methylcyclopentanes, see: E. L. Eliel, S. H. Wilen, *Stereochemistry of Organic Compounds*, Wiley, New York, **1994**, pp. 758–759. In the case of transition states **C–F**, it may be proposed that the tertiary carbinol center avoids the flap side of the envelope due to steric reasons.
- [10] K. Tomioka, T. Suenaga, K. Koga, *Tetrahedron Lett.* **1986**, 27, 369–372.