


Efficient Synthesis of Benzothiophenes by an Unusual Palladium-Catalyzed Vinylic C–S Coupling**

Christopher S. Bryan, Julia A. Braunger, and Mark Lautens*

Transition-metal-catalyzed cross-coupling reactions have become indispensable tools for the synthetic chemist.^[1,2] The construction of carbon–heteroatom bonds is of particular interest. Consequently, a large number of general and efficient systems for the catalytic amination and etherification of aryl and vinyl halides have been described.^[3,4] In contrast, research in the field of carbon–sulfur coupling reactions has lagged behind, largely because of sulfur's long-standing reputation as a catalyst poison.^[5–7] Despite this difficulty, coupling reactions of thiols with aryl halides under the catalysis of Pd,^[8] Cu,^[9] Ni,^[10] Co,^[11] Fe,^[12] and In^[13] have been reported. Reactions of vinyl halides are comparatively rare: only a handful of publications describe the vinylation of thiols,^[14,15] and the palladium-catalyzed intramolecular coupling is unknown.^[16]

We anticipated that the use of Pd could facilitate access to a number of tandem coupling reactions in which an S-vinylation reaction could be combined with orthogonal carbon–carbon bond formation in a single synthetic operation. Tandem/domino reactions are an area of intense research, as they enable streamlined synthetic sequences and a consequent reduction of waste.^[17] Herein, we disclose a tandem catalytic reaction of a *gem*-dihalovinyl thiophenol system in which an intramolecular S-vinylation is paired with an intermolecular C–C bond-forming reaction (a Suzuki–Miyaura, Heck, or Sonogashira reaction) to yield 2-substituted benzothiophenes (Scheme 1). These heterocycles form the core of a number of medically important molecules, such as raloxifene^[18] and zileuton.^[19] However, catalytic




routes to these targets have only recently been developed.^[20] To the best of our knowledge, the reaction described herein is the first example of a tandem catalytic process that incorporates a C–S coupling, as well as the first example of the palladium-catalyzed vinylation of a thiol.

We and others have previously reported the synthesis of indoles substituted at the 2-position by tandem coupling reactions of *gem*-dihalovinyl anilines.^[21–27] In these systems, an intramolecular amination or amidation was combined with an inter- or intramolecular Suzuki–Miyaura, Heck, or Sonogashira coupling, amidation, direct arylation, or carbonylation. Similar transformations involving C–S coupling would provide rapid access to benzothiophenes and expand the range of available tandem coupling reactions.

Our initial attempts were aimed at combining the carbon–sulfur bond-forming process with a Suzuki–Miyaura coupling (Table 1). We found that the conditions developed previously for the aniline system^[22b] were less effective for the thiophenol substrate **1a**: the desired product was isolated in only 48% yield (Table 1, entry 1).^[28] The screening of various bases (Table 1, entries 1–4, 8), precatalysts (entries 5–7), and ligand/catalyst ratios (entries 9 and 10) revealed that the best results were obtained when the reaction was carried out with PdCl₂, SPhos,^[29] and anhydrous K₃PO₄/Et₃N in dioxane at 110 °C. Under these conditions, **2a** was formed in 89% yield (Table 1, entry 10). Lowering of the catalyst loading to 1 mol% led to enhanced turnover with minimal impact on the yield (Table 1, entry 12).

With these optimized conditions in hand, we investigated the effect of varying the boronic acid on the tandem process (Table 2). The reaction was found to tolerate changes in the electronic nature of the boronic acid. Electron-poor boronic acids displayed similar reactivity to that of electron-rich 3,4-dimethoxyphenylboronic acid: the boronic acid used for optimization of the reaction (Table 2, entries 1–4). The coupling proceeded equally well when a sterically congested boronic acid was used (Table 2, entry 5). The use of hetero-aromatic boronic acids gave the best results: the corresponding heterocyclic products were obtained in up to 99% yield



Scheme 1. Retrosynthetic strategy.

[*] C. S. Bryan, J. A. Braunger, Prof. Dr. M. Lautens
Davenport Laboratories, Department of Chemistry
University of Toronto
80 St. George Street, Toronto, Ontario (Canada)
Fax: (+1) 416-946-8185
E-mail: mlautens@chem.utoronto.ca
Homepage: <http://www.chem.utoronto.ca/staff/ML/>

[**] This research was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), the Merck Frosst Centre for Therapeutic Research, and the University of Toronto. We thank Dr. Valentina Aureggi for her intellectual contributions.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200902843>.

solution was degassed with argon for an additional 5 min. The septum was then replaced with a Teflon microwave cap, the vial was sealed, and the reaction mixture was stirred for 10 min and then heated in an oil bath for 16 h at 110°C. The flask was then cooled to room temperature, and the solids were removed by vacuum filtration through a plug of silica gel and washed with EtOAc. The solution was concentrated, and the crude product was purified by column chromatography over silica gel (eluent: Et₂O/pentane 0–2%).

Received: May 27, 2009

Published online: August 18, 2009

Keywords: cross-coupling · palladium · sulfur heterocycles · synthetic methods · tandem catalysis

- [1] *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. 1 (Ed.: E.-i. Negishi), Wiley-Interscience, New York, **2002**.
- [2] For reviews on metal-catalyzed cross-coupling reactions, see: a) J. F. Hartwig, *Nature* **2008**, *455*, 314; b) G. Fu, *Acc. Chem. Res.* **2008**, *41*, 1555; c) K. C. Nicolaou, P. G. Bulger, D. Sarlah, *Angew. Chem.* **2005**, *117*, 4516; *Angew. Chem. Int. Ed.* **2005**, *44*, 4442; d) *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed. (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, **2004**.
- [3] For reviews on catalytic amination, see: a) J. F. Hartwig, *Acc. Chem. Res.* **2008**, *41*, 1534; b) D. S. Surry, S. L. Buchwald, *Angew. Chem.* **2008**, *120*, 6438; *Angew. Chem. Int. Ed.* **2008**, *47*, 6338; c) J. Barluenga, C. Valdés, *Chem. Commun.* **2006**, 4891; d) S. V. Ley, A. Thomas, *Angew. Chem.* **2003**, *115*, 5558; *Angew. Chem. Int. Ed.* **2003**, *42*, 5400; Corrigendum: S. V. Ley, A. Thomas, *Angew. Chem.* **2004**, *116*, 1061; *Angew. Chem. Int. Ed.* **2004**, *43*, 1043.
- [4] For catalytic etherification, see: a) C. H. Burgos, T. E. Barder, X. Huang, S. L. Buchwald, *Angew. Chem.* **2006**, *118*, 4427; *Angew. Chem. Int. Ed.* **2006**, *45*, 4321; b) A. V. Vorogushin, X. Huang, S. L. Buchwald, *J. Am. Chem. Soc.* **2005**, *127*, 8146; c) K. E. Torraca, X. Huang, C. A. Parrish, S. L. Buchwald, *J. Am. Chem. Soc.* **2001**, *123*, 10770; d) Q. Shelby, N. Kataoka, G. Mann, J. F. Hartwig, *J. Am. Chem. Soc.* **2000**, *122*, 10718; e) R. A. Altman, A. Shafir, A. Choi, P. A. Lichtor, S. L. Buchwald, *J. Org. Chem.* **2008**, *73*, 284; f) R. Gujadhur, C. G. Bates, D. Venkataraman, *Org. Lett.* **2001**, *3*, 4315.
- [5] Sulfur-containing compounds display strong coordinating and adsorptive properties; see: a) E. Alvaro, J. F. Hartwig, *J. Am. Chem. Soc.* **2009**, *131*, 7858; b) G. Mann, D. Barañano, J. F. Hartwig, A. L. Rheingold, I. A. Guzei, *J. Am. Chem. Soc.* **1998**, *120*, 9205; c) J. Louie, J. F. Hartwig, *J. Am. Chem. Soc.* **1995**, *117*, 11598; d) L. L. Hegedus, R. W. McCabe in *Catalyst Poisoning*, Marcel Dekker, New York, **1984**; e) A. T. Hutton in *Comprehensive Coordination Chemistry*, Vol. 5 (Eds.: G. Wilkinson, R. D. Gillard, J. A. McCleverty), Pergamon, Oxford, **1984**, p. 1131.
- [6] During the course of this study, an excellent publication describing the use of Zn salts to prevent catalyst poisoning in a palladium-catalyzed C–S coupling reaction appeared: C. C. Eichman, J. P. Stambuli, *J. Org. Chem.* **2009**, *74*, 4005.
- [7] For reviews on metal-catalyzed C–S bond formation, see: a) I. P. Beletskaya, V. P. Ananikov, *Eur. J. Org. Chem.* **2007**, 3431; b) T. Kondo, T.-a. Mitsudo, *Chem. Rev.* **2000**, *100*, 3205.
- [8] a) J.-Y. Lee, P. H. Lee, *J. Org. Chem.* **2008**, *73*, 7413; b) M. A. Fernández-Rodríguez, Q. Shen, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, *128*, 2180; c) C. Mispelaere-Canivet, J. F. Spindler, S. Perrio, P. Beslin, *Tetrahedron* **2005**, *61*, 5253; d) M. Murata, S. L. Buchwald, *Tetrahedron* **2004**, *60*, 7397; e) T. Itoh, T. Mase, *Org. Lett.* **2004**, *6*, 4587; f) G. Y. Li, *Angew. Chem.* **2001**, *113*, 1561; *Angew. Chem. Int. Ed.* **2001**, *40*, 1513; g) U. Schopfer, A. Schlapbach, *Tetrahedron* **2001**, *57*, 3069; h) N. Zheng, J. C. McWilliams, F. J. Fleitz, J. D. Armstrong III, R. P. Volante, *J. Org. Chem.* **1998**, *63*, 9606; i) T. Migita, T. Shimizu, Y. Asami, J.-i. Shiobara, Y. Kato, M. Kosugi, *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385; j) M. Kosugi, T. Shimizu, T. Migita, *Chem. Lett.* **1978**, 13.
- [9] a) D. Ma, S. Xie, P. Xue, X. Zhang, J. Dong, Y. Jiang, *Angew. Chem.* **2009**, *121*, 4289; *Angew. Chem. Int. Ed.* **2009**, *48*, 4222; b) M. Carril, R. SanMartin, E. Domínguez, I. Tellitu, *Chem. Eur. J.* **2007**, *13*, 5100; c) A. K. Verma, J. Singh, R. Chaudhary, *Tetrahedron Lett.* **2007**, *48*, 7199; d) X. Lv, W. Bao, *J. Org. Chem.* **2007**, *72*, 3863; e) C. G. Bates, R. K. Gujadhur, D. Venkataraman, *Org. Lett.* **2002**, *4*, 2803.
- [10] a) Y. Zhang, K. C. Ngeow, J. Y. Ying, *Org. Lett.* **2007**, *9*, 3495; b) H. J. Cristau, B. Chabaud, A. Chene, H. Christol, *Synthesis* **1981**, 892.
- [11] Y.-C. Wong, T. T. Jayanth, C.-H. Cheng, *Org. Lett.* **2006**, *8*, 5613.
- [12] A. Correa, M. Carril, C. Bolm, *Angew. Chem.* **2008**, *120*, 2922; *Angew. Chem. Int. Ed.* **2008**, *47*, 2880.
- [13] V. P. Reddy, A. V. Kumar, K. Swapna, K. R. Rao, *Org. Lett.* **2009**, *11*, 1697.
- [14] For examples of intermolecular copper-catalyzed C–S coupling reactions of vinyl halides, see: a) M. S. Kabir, M. L. Van Linn, A. Monte, J. M. Cook, *Org. Lett.* **2008**, *10*, 3363; b) B. Jiang, H. Tian, A.-G. Huang, M. Xu, *Org. Lett.* **2008**, *10*, 2737; c) Y. Zheng, X. Du, W. Bao, *Tetrahedron Lett.* **2006**, *47*, 1217; d) C. G. Bates, P. Saejueng, M. Q. Doherty, D. Venkataraman, *Org. Lett.* **2004**, *6*, 5005.
- [15] For an intramolecular copper-catalyzed S-vinylation, see: Q. Zhao, L. Li, Y. Fang, D. Sun, C. Li, *J. Org. Chem.* **2009**, *74*, 459.
- [16] For examples of intermolecular palladium-catalyzed S-vinylation, see: a) X. Moreau, J.-M. Campagne, *J. Organomet. Chem.* **2003**, *687*, 322; b) I. G. Trostyanskaya, D. Y. Tititskiy, E. A. Anufrieva, A. A. Borisenko, M. A. Kazankova, I. P. Beletskaya, *Russ. Chem. Bull.* **2001**, *50*, 2095; c) A. M. Rane, E. I. Miranda, J. A. Soderquist, *Tetrahedron Lett.* **1994**, *35*, 3225; d) S.-I. Murahashi, M. Yamamura, K.-i. Yanagisawa, N. Mita, K. Kondo, *J. Org. Chem.* **1979**, *44*, 2408.
- [17] a) L.-F. Tietze, G. Brasche, K. Gericke, *Domino Reactions in Organic Synthesis*, Wiley-VCH, Weinheim, **2006**; b) D. E. Fogg, E. S. dos Santos, *Coord. Chem. Rev.* **2004**, *248*, 2365.
- [18] Z. Qin, I. Kasrati, E. P. Chandrasena, H. Liu, P. Yao, P. A. Petukhov, J. L. Bolton, G. R. J. Thatcher, *J. Med. Chem.* **2007**, *50*, 2682.
- [19] C.-N. Hsiao, T. Kolasa, *Tetrahedron Lett.* **1992**, *33*, 2629.
- [20] For catalytic approaches to benzothiophenes, see: a) K. Inamoto, Y. Arai, K. Hiroya, T. Doi, *Chem. Commun.* **2008**, 5529; b) I. Nakamura, T. Sato, Y. Yamamoto, *Angew. Chem.* **2006**, *118*, 4585; *Angew. Chem. Int. Ed.* **2006**, *45*, 4473; c) M. C. Willis, D. Taylor, A. T. Gillmore, *Tetrahedron* **2006**, *62*, 11513.
- [21] C. Bryan, V. Auregi, M. Lautens, *Org. Synth.* **2009**, *86*, 36.
- [22] a) Y.-Q. Fang, M. Lautens, *J. Org. Chem.* **2008**, *73*, 538; b) Y.-Q. Fang, M. Lautens, *Org. Lett.* **2005**, *7*, 3549; c) Y.-Q. Fang, J. Yuen, M. Lautens, *J. Org. Chem.* **2007**, *72*, 5152; d) Y.-Q. Fang, R. Karisch, M. Lautens, *J. Org. Chem.* **2007**, *72*, 1341; e) S. Thielges, E. Meddah, P. Bissere, J. Eustache, *Tetrahedron Lett.* **2004**, *45*, 907.
- [23] A. Fayol, Y.-Q. Fang, M. Lautens, *Org. Lett.* **2006**, *8*, 4203.
- [24] M. Nagamochi, Y.-Q. Fang, M. Lautens, *Org. Lett.* **2007**, *9*, 2955.
- [25] J. Yuen, Y.-Q. Fang, M. Lautens, *Org. Lett.* **2006**, *8*, 653–656.
- [26] a) C. S. Bryan, M. Lautens, *Org. Lett.* **2008**, *10*, 4633; for an example of direct arylation paired with Suzuki–Miyaura coupling, see: b) D. Chai, M. Lautens, *J. Org. Chem.* **2009**, *74*, 3054.
- [27] T. O. Viera, L. A. Meaney, Y.-L. Shi, H. Alper, *Org. Lett.* **2008**, *10*, 4899.
- [28] The starting materials, such as **1a**, can be prepared readily from the corresponding 2-halobenzaldehydes; see the Supporting Information for details.

- [29] T. E. Barder, S. D. Walker, J. R. Martinelli, S. L. Buchwald, *J. Am. Chem. Soc.* **2005**, *127*, 4685.
- [30] For the cross-coupling of trifluoroborates, see: a) G. A. Molander, D. L. Sandroock, *J. Am. Chem. Soc.* **2008**, *130*, 15792; b) Y. Bolshan, R. A. Batey, *Angew. Chem.* **2008**, *120*, 2139; *Angew. Chem. Int. Ed.* **2008**, *47*, 2109; c) S. Darses, J.-P. Genêt, J.-L. Brayer, J.-P. Demoute, *Tetrahedron Lett.* **1997**, *38*, 4393.
- [31] For N-heterocycle synthesis through the palladium-catalyzed intramolecular addition of a nitrogen nucleophile to an alkyne, see: a) L. Kaspar, L. Ackermann, *Tetrahedron Lett.* **2005**, *61*, 11311; b) S. Cacchi, V. Carnicelli, F. Marinelli, *J. Organomet. Chem.* **1994**, *475*, 289; c) K. Utimoto, H. Miwa, H. Nozaki, *Tetrahedron Lett.* **1981**, *22*, 4277.
- [32] T. Jeffery, *Tetrahedron* **1996**, *52*, 10113–10130.
-