


Coupling Reactions of Zirconate Complexes Induced by Carbonyl Compounds**

Chanjuan Xi,* Xiaoyu Yan, Wei You, and Tamotsu Takahashi*

During the past few decades the chemistry of 16-electron organozirconocene complexes has been extensively explored and a tremendous number of applications in synthetic chemistry have been found.^[1] However, organozirconate complexes, which usually have an 18-electron configuration with two cyclopentadienyl (Cp) rings, three Zr–C bonds, and alkali counter ions, have rarely been explored in spite of being the supposed intermediates in a number of stoichiometric and catalytic reactions, and able to form C–C bonds through a successful 1,2-migration of organozirconates.^[2–4] Our research group has an interest in the above mentioned subjects, and we have recently reported^[5] that by adding quinone as an oxidant to a mixture of alkynylzirconates then zirconocenendiyne compounds are afforded: these compounds can be converted into various endiynes through coupling reaction with electrophiles. These results prompted us to further study the chemistry of organozirconate complexes. Herein, we report the coupling reactions of three-membered alkynylzirconates induced by carbonyl compounds. The unique features of this reaction involve the carbonyl compound acting as an electrophile, participating in the reaction with zirconates, and also inducing the coupling of an alkynyl group and an sp²-hybridized carbon atom of zirconacyclopentene. To the best of our knowledge, this type of reaction has not been reported. When acyl compounds were used as electrophiles, then functionalized allenes were formed. When aldehydes were used as electrophiles, then dienols were formed. These outcomes are illustrated in Scheme 1.


The reaction of zirconate **1a**, which was generated by the treatment of *n*BuLi with bis(phenylethynyl)zirconocene at room temperature,^[6] with chloroformate at 0 °C for five hours afforded allene **3a** in 73 % yield after hydrolysis. Quenching



Scheme 1. Diverse reactivity of zirconates.

the reaction mixture with DCl in D₂O provided allene **5a** in a comparable yield with 91 % deuterium incorporation. This result indicates that the zirconium-containing complex **6a** is formed (Scheme 2).

Oxazirconacycle **6a** was formed in 74 % yield (based on NMR spectroscopy). The ¹H NMR spectrum of **6a** showed a singlet resonance at $\delta = 5.96$ ppm, which was assigned to the protons on the Cp ring. In the ¹³C NMR spectrum, the signal



Scheme 2. The reaction with chloroformate.

for the carbon atoms in the Cp ring appeared at $\delta = 104.5$ ppm, and the signal for the allenic-type sp-hybridized carbon atom appeared at $\delta = 198.5$ ppm. The carbon signals in Zr–C(Ph) and –C(OEt)O appeared at $\delta = 149.5$ and 169.6 ppm, respectively.

The allene group is a versatile functionality because it is useful as either a nucleophile or an electrophile as well as a substrate for cycloaddition reactions. This multiple reactivity make allenes as excellent candidates for synthetic manipulation.^[7] The reaction described here allowed the efficient synthesis of various substituted allenes. A variety of acyl compounds were subjected to this novel reaction and all reactions afforded analogous products in good to high yields after hydrolysis. Notably, the acyl compounds induced

[*] Prof. C. Xi, X. Yan, W. You

Key Laboratory of Bioorganic Phosphorus Chemistry & Chemical Biology (Ministry of Education), Department of Chemistry Tsinghua University, Beijing 100084 (China)

and

State Key Laboratory of Elemento-Organic Chemistry

Nankai University, Tianjin 300071 (China)

Fax: (+86) 10-6277-1194

E-mail: cxixi@tsinghua.edu.cn

Prof. T. Takahashi

Catalysis Research Center and Graduate School of Life Science

Hokkaido University, Sapporo 001-0021 (Japan)


[**] We thank the National Natural Science Foundation of China (20872076), and the Doctoral Program of Higher Education (200800030072) for financial support. We also thank Dr. Haifang Li for her kind help with measurement of HRMS data.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.200904255>.

coupling between an alkynyl group and an sp^2 -hybridized carbon atom with high regioselectivity. This result is due to the higher reactivity of the carbon atom attached to the alkyl group as compared to the carbon atom attached to the aryl group. The representative results are summarized in Table 1. Chloroformates afforded the corresponding allenes in 47–61% yields (Table 1, entries 1–5). The addition of esters similarly gave allenes in good yields (Table 1, entries 6–8). The diethylcarbamyl chloride afforded the corresponding product in 55% yield (Table 1, entry 9). When acyl chloride and anhydride were treated with zirconates under similar reaction conditions the desired products were not observed.

The remaining Zr–C bond of oxazirconacycle **6** was further treated with allyl bromide in the presence of one equivalent of CuCl^[8] and afforded fully substituted allenes **7a** and **7b** in moderate yields, respectively (Scheme 3).

Based on the above results, we propose the following mechanism for this reaction (Scheme 4). The zirconate [LiCp₂Zr(C≡CR³)(R¹C=CR²)] **1** undergoes a coupling reaction to give intermediate **2** in the presence of an acyl



7a: R = Ph, yield = 59% (43%)

7b: R = *p*BuC₆H₄, yield = 58% (45%)

Scheme 3. The reaction of oxazirconacycle **6** with allyl bromide (yield of isolated product given in parentheses).

compound. The intermediate **2** reacts with an acyl compound through nucleophilic substitution to form **8** with elimination of a lithium salt and then undergoes cyclic oxidation/rearrangement (path a) to give **6**. Alternatively, intermediate **8** undergoes oxidative coupling of the carbonyl and alkenyl groups to give oxazirconacycle **9** with an alkynyl moiety at the α position (path b),^[9] which subsequently rearranges to **6**. Hydrolysis of **6** affords the product **3**.

In light of the unusual reactivity of the organozirconate,

we further investigated the reaction of zirconate **1a** with benzaldehyde (Scheme 5). Interestingly, dienol **4a** was formed in good yield after hydrolysis. Deuteriolysis instead of hydrolysis afforded the dideuterated compound **10a** in 66% yield with 93% and 97% deuterium incorporation. The structure of product **4a** was confirmed by ¹H NMR, ¹³C NMR, HSQC, HMBC, and NOESY experiments.^[10]


Intermediate **11a** was also characterized by NMR spectroscopy. The ¹H NMR spectrum of **11a** showed two signals at δ = 5.72 and 6.00 ppm which are assigned to the protons of two Cp rings, and a resonance at δ = 5.39 ppm is assigned to the proton in –OCHPh. In the ¹³C NMR spectrum, the signals for the carbon atoms of the Cp ring appear at δ = 111.0 and 111.6 ppm, and the resonance for the carbon atoms of the diene appear at δ = 146.9, 150.4, 154.8, and 183.4 ppm.

A variety of aromatic aldehydes were subjected to the reaction. The results were summarized in Table 2. Aromatic aldehydes bearing a chlorine, fluorine, or phenyl group were treated with zirconates and led to the corresponding products in good yields (Table 2, entries 3–6, and 8). The *p*-methylbenzaldehyde was treated with zirconates and led to the


Table 1: Formation of functionalized allenes by the reaction of zirconates with acyl compounds.

Entry	Zirconate	Acyl compound	Product ^[a]	Yield [%] ^[b]
1			3a	73(57)
2	1a		3b	65(47)
3			3c	76(61)
4			3d	69(53)
5			3e	67(52)
6	1a		3f	46(32)
7	1a		3g	44(28)
8			3h	40(31)
9	1c		3i	67(55)

[a] Single isomer, the stereoselectivity was not defined. [b] Yield is based on ¹H NMR spectroscopy, yield of isolated product given in parentheses. Hex = hexyl, Tol = tolyl.




Scheme 4. Proposed mechanism for the reaction with acyl compounds.



Scheme 5. The reaction with benzaldehyde.

corresponding products in moderate yields (Table 2, entries 2 and 7). When aliphatic aldehydes were treated with zirconates under similar reaction conditions, the desired products were not observed, but aldol products were obtained.

These results indicate that aldehydes behaved differently to acyl compounds. A plausible reaction mechanism is shown in Scheme 6. Firstly, the oxygen atom of the carbonyl group coordinates to zirconate **1** and induces coupling of the alkynyl and an sp^2 -hybridized carbon atom of zirconacyclopentadiene to give intermediate **12**, which forms an equilibrium between **13**. In this case, the reaction pathway differs from the reaction with acyl compound. A possible explanation is that the nucleophilicity of the carbon atom attached to zirconium is not high enough to react with an aldehyde. On the other hand,



Scheme 6. Plausible mechanism for the formation of dienols.

Table 2: Formation of dienols by the reaction of zirconates with aldehydes.

Entry	Zirconate	Aldehyde	Product	Yield [%] ^[a]
1	1a	PhCHO	4a	66(54)
2	1a	<i>p</i> -MeC ₆ H ₄ CHO	4b	51(45)
3	1a	<i>p</i> -FC ₆ H ₄ CHO	4c	69(57)
4	1a	<i>p</i> -ClC ₆ H ₄ CHO	4d	66(50)
5	1c	<i>p</i> -PhC ₆ H ₄ CHO	4e	64(52)
6	1c	<i>p</i> -ClC ₆ H ₄ CHO	4f	70(59)
7	1e	<i>p</i> -MeC ₆ H ₄ CHO	4g	46(37)
8	1e	<i>p</i> -ClC ₆ H ₄ CHO	4h	65(50)

[a] Yield is based on ¹H NMR spectroscopy; yield of isolated product given in parentheses.

oxidative coupling of an aldehyde and an alkyne on the zirconocene(II) is preferred.^[11] The intermediate **13** undergoes oxidative coupling to form **11**. Hydrolysis of **11** affords the product **4**.^[12]

In summary, we have demonstrated a novel coupling reaction of alkynylzirconate complexes with carbonyl compounds. A unique feature of this reaction is that carbonyl compounds not only participated in the reaction with zirconates but also induced coupling of an alkynyl and alkenyl group of zirconacyclopentadiene with high regioselectivity. In this reaction, two alkyne groups, introduced as to alkynyllithiums and an organolithium, and carbonyl compounds were selectively coupled in one-pot by the reaction of [Cp₂ZrCl₂] to afford highly substituted allene and dienol derivatives. Investigation of the reaction mechanism and further application of this chemistry are in progress.

Experimental Section

Typical procedure for the reaction of alkynylzirconates with acyl compounds: *n*BuLi (4.5 mmol, 1.6 M solution in hexane) was added to a solution of phenylacetylene (3.0 mmol) at -78°C and stirred for 1 h. Then [Cp₂ZrCl₂] (1.5 mmol) was added and the solution was stirred at -78°C for 1 h then warmed to room temperature and kept at this

temperature for 12 h. After cooling the reaction mixture to 0 °C, ethyl chloroformate (1.0 mmol) was added and the reaction mixture was stirred for 5 h. The reaction mixture was quenched with 6 N HCl and extracted with ethyl acetate. The organic extract was dried over MgSO₄, and the filtrate was concentrated under reduced pressure. Purification of the residue by column chromatography on silica gel (hexane/ethyl acetate 50:1) afforded **3a** as yellow liquid (190 mg) in 57 % yield. ¹H NMR (300 MHz, CDCl₃, Me₄Si): δ = 0.82 (t, ³J_{HH} = 7.2 Hz, 3H), 1.12 (t, ³J_{HH} = 7.2 Hz, 3H), 1.21–1.44 (m, 4H), 2.05 (dt, ³J_{HH} = 7.3 Hz, ²J_{HH} = 2.4 Hz, 2H), 4.00 (dq, ³J_{HH} = 7.0 Hz, ²J_{HH} = 10.8 Hz, 1H), 4.04 (dq, ³J_{HH} = 7.0 Hz, ²J_{HH} = 10.8 Hz, 1H), 4.33 (q, ⁴J_{HH} = ⁵J_{HH} = 2.4 Hz, 1H), 6.22 (q, ⁴J_{HH} = ⁵J_{HH} = 2.4 Hz, 1H), 7.27–7.33 ppm (m, 10H); ¹³C NMR (75 MHz, CDCl₃, Me₄Si): δ = 14.0, 14.1, 22.5, 29.8, 31.7, 55.9, 61.2, 98.1, 108.5, 126.7, 126.8, 127.4, 128.2, 128.5, 129.0, 135.0, 137.0, 171.8, 203.4 ppm. HRMS calcd for C₂₃H₂₆O₂ ([M]⁺): 334.1933; found: 334.1925.

Received: July 31, 2009

Published online: September 23, 2009

Keywords: acyl compounds · aldehydes · allenes · dienols · organozirconates

- [1] a) For recent reviews, see: H. G. Alt, A. Köppl, *Chem. Rev.* **2000**, *100*, 1205; b) G. Erker, *Acc. Chem. Res.* **2001**, *34*, 309; c) E. Negishi, *Pure Appl. Chem.* **2001**, *73*, 239; d) P. Wipf, C. Kendall, *Chem. Eur. J.* **2002**, *8*, 1778; e) E. I. Marek, *Titanium and Zirconium in Organic Synthesis*, Wiley-VCH, Weinheim, **2002**; f) K. Suzuki, P. Wipf, *Tetrahedron* **2004**, *60*, 1267; g) T. Takahashi, Z. Xi, R. Hara, *Trends Organomet. Chem.* **1997**, *2*, 117.
- [2] a) E. Negishi, K. Akiyoshi, B. O'Connor, K. Takagi, G. Wu, *J. Am. Chem. Soc.* **1989**, *111*, 3089; b) K. Takagi, C. J. Rousset, E. Negishi, *J. Am. Chem. Soc.* **1991**, *113*, 1440; c) T. Takahashi, T. Seki, Y. Nitto, M. Saburi, C. J. Rousset, E. Negishi, *J. Am. Chem. Soc.* **1991**, *113*, 6266; d) T. Takahashi, S. Huo, R. Hara, Y. Noguchi, K. Nakajima, W.-H. Sun, *J. Am. Chem. Soc.* **1999**, *121*, 1094; e) Y. Dumond, E. Negishi, *J. Am. Chem. Soc.* **1999**, *121*, 11223; f) R. Choukroun, J. Zhao, C. Lorber, P. Cassoux, B. Donnadieu, *Chem. Commun.* **2000**, 1511.
- [3] a) A. H. Hoveyda, J. P. Morken, A. F. Houry, Z. Xu, *J. Am. Chem. Soc.* **1992**, *114*, 6692; b) T. Takahashi, N. Suzuki, M. Kageyama, Y. Nitto, M. Saburi, E. Negishi, *Chem. Lett.* **1991**, 1579; c) K. S. Knight, R. M. Waymouth, *J. Am. Chem. Soc.* **1991**, *113*, 6268; d) D. P. Lewis, P. M. Muller, R. J. Whitby, R. U. H. Jones, *Tetrahedron Lett.* **1991**, *32*, 6797; e) C. J. Rousset, E. Negishi, N. Suzuki, T. Takahashi, *Tetrahedron Lett.* **1992**, *33*, 1965; f) J.-P. Majoral, M. Zablocka, *New J. Chem.* **2005**, *29*, 32.
- [4] a) G. J. Gordon, R. J. Whitby, *Chem. Commun.* **1997**, 1321; b) S. F. Fillery, G. J. Gordon, T. Luker, R. J. Whitby, *Pure Appl. Chem.* **1997**, *69*, 633; c) A. Kasatkin, R. J. Whitby, *Tetrahedron Lett.* **1997**, *38*, 4857; d) A. Kasatkin, R. J. Whitby, *J. Am. Chem. Soc.* **1999**, *121*, 7039; e) A. N. Kasatkin, R. J. Whitby, *Tetrahedron Lett.* **1999**, *40*, 9353; f) A. N. Kasatkin, R. J. Whitby, *Tetrahedron Lett.* **2000**, *41*, 6211; g) A. N. Kasatkin, R. J. Whitby, *Tetrahedron Lett.* **2000**, *41*, 6201; h) A. Kasatkin, R. J. Whitby, *Tetrahedron* **2003**, *59*, 9857.
- [5] C. Xi, Y. Liu, X. Yan, C. Chen, *J. Organomet. Chem.* **2007**, *692*, 4612.
- [6] E. Thomas, S. Dixon, R. J. Whitby, *Angew. Chem.* **2006**, *118*, 7228; *Angew. Chem. Int. Ed.* **2006**, *45*, 7070.
- [7] a) V. Lavallo, G. D. Frey, S. Kousar, B. Donnadieu, G. Bertrand, *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 13569; b) N. Krause, A. S. K. Hashmi, *Modern Allene Chemistry*, Wiley-VCH, Weinheim, **2004**; c) J. C. Aponte, G. B. Hammond, B. Xu, *J. Org. Chem.* **2009**, *74*, 4623.
- [8] For CuCl-catalyzed or -mediated reactions of organozirconium compounds, see: a) L. M. Venanzi, R. Lehman, R. Keil, B. H. Lipshutz, *Tetrahedron Lett.* **1992**, *33*, 5857; b) T. Takahashi, M. Kotora, K. Kasai, N. Suzuki, *Tetrahedron Lett.* **1994**, *35*, 5685; c) T. Honda, M. Mori, *Organometallics* **1996**, *15*, 5464; d) K. Kasai, M. Kotora, N. Suzuki, T. Takahashi, *J. Chem. Soc. Chem. Commun.* **1995**, 109; e) B. H. Lipshutz, M. Segi, *Tetrahedron* **1995**, *51*, 4407; f) T. Takahashi, R. Hara, Y. Nishihara, M. Kotora, *J. Am. Chem. Soc.* **1996**, *118*, 5154; g) F. Dekura, T. Honda, M. Mori, *Chem. Lett.* **1997**, 825; h) T. Takahashi, W.-H. Sun, Y. Liu, K. Nakajima, M. Kotora, *Organometallics* **1998**, *17*, 3841; i) C. Xi, M. Kotora, T. Takahashi, *Tetrahedron Lett.* **1999**, *40*, 2375.
- [9] a) D. P. Hsu, W. M. Davis, S. L. Buchwald, *J. Am. Chem. Soc.* **1993**, *115*, 10394; b) U. Rosenthal, A. Ohff, W. Baumann, R. Kempe, A. Tillack, V. V. Burlakov, *Angew. Chem.* **1994**, *106*, 1678; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1605.
- [10] HSQC, HMBC, and NOESY experiments show cross-peak correlations for **4a**, see the Supporting Information.
- [11] a) T. Takahashi, M. Kageyama, V. Denisov, R. Hara, E. Negishi, *Tetrahedron Lett.* **1993**, *34*, 687; b) T. Takahashi, C. Xi, Z. Xi, M. Kageyama, R. Fischer, K. Nakajima, E. Negishi, *J. Org. Chem.* **1998**, *63*, 6802.
- [12] a) D. Kashinath, C. Mioskowski, J. R. Falck, M. Goli, S. Meunier, R. Baati, A. Wagner, *Org. Biomol. Chem.* **2009**, *7*, 1771; b) P. A. Clarke, G. A. Rolla, A. P. Cridland, A. A. Gill, *Tetrahedron* **2007**, *63*, 9124; c) G. Metha, H. S. P. Rao, *The Chemistry of Dienes and Polyenes, Vol. 1* (Ed.: Z. Rappoport, Wiley, Chichester, **1997**, p. 359.